



**Table 1**

## SCIENTIFIC DENTAL JOURNAL







About

Articles

Authors

Search

Subscribe

Contact Us

Login

## Table of Contents



September-December 2019

Volume 3 | Issue 3

Page Nos. 75-108

Online since Monday, October 14, 2019

Accessed 5,715 times.

### PDF access policy

Journal allows immediate open access to content in HTML + PDF

[View issue as eBook](#)

[Issue statistics](#)

[RSS](#)

[Next Issue](#)

[Previous Issue](#)

[POPULAR ARTICLES](#)


[JOIN AS REVIEWER](#)

[GET EMAIL ALERTS](#)

This website uses cookies. By continuing to use this website you are giving consent to cookies being used. For information on cookies and how you can disable them visit our [Privacy and Cookie Policy](#).

[AGREE & PROCEED](#)

## ORIGINAL ARTICLES




### Effects of brushing and immersion in denture cleanser on the surface roughness of polymethyl methacrylate p. 75

Agnes Victoria Kurniawan, Octarina, Hernindya Dwifulqi

DOI:10.4103/SDJ.SDJ\_19\_19

[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[Sword Plugin for Repository\]](#)<sup>Beta</sup>




### The rational use of antibiotics by dentists for odontogenic infection treatment in Kupang, East Nusa Tenggara p. 81

Manginar Sidabutar, Friska D Simamora, Faber Sidabutar, Yuliana R Dolo Wain, Mira Enjelina Malelak

DOI:10.4103/SDJ.SDJ\_26\_19

[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[Sword Plugin for Repository\]](#)<sup>Beta</sup>




### Efficacy of disinfectants on microbial contaminated toothbrushes p. 85

Angelica Tiara, Armelia Sari Widyarman, Christine Anastasia Rovani

DOI:10.4103/SDJ.SDJ\_23\_19

[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[Sword Plugin for Repository\]](#)<sup>Beta</sup>



### Viability test of ethanol extract of beluntas (*pluchea indica*) leaves on *In vitro* fibroblast cells p. 90

Rani Wulan Sari, Natallia Pranata, Vinna Kurniawati Sugiaman

DOI:10.4103/SDJ.SDJ\_18\_19

[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[Sword Plugin for Repository\]](#)<sup>Beta</sup>

## CASE REPORTS



### Removing a fractured instrument from the root canal using ultrasonic tips p. 95

Aryadi Subrata, Nadia Hardini

DOI:10.4103/SDJ.SDJ\_15\_19

[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[Sword Plugin for Repository\]](#)<sup>Beta</sup>



### Bone regeneration on chronic apical abscess after root canal treatment on left mandibular first molar: A case report p. 100

Diana Puspa Indah, Eko Fibryanto, Ie Elline Istanto

DOI:10.4103/SDJ.SDJ\_17\_19

[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[Sword Plugin for Repository\]](#)<sup>Beta</sup>



### Clinical assessment of a squamous cell carcinoma located in the posterior oral cavity p. 105

Wahyuning Ratnawidya, Endah Ayu Tri Wulandari, Ening Krisnuhoni, Yuniardini Septorini Wimardhani, Anak Iamaroon

DOI:10.4103/SDJ.SDJ\_14\_19

[\[ABSTRACT\]](#) [\[HTML Full text\]](#) [\[PDF\]](#) [\[Mobile Full text\]](#) [\[EPub\]](#) [\[Sword Plugin for Repository\]](#)<sup>Beta</sup>



## Original Article

# Viability Test of Ethanol Extract of *Beluntas* (*Pluchea indica*) Leaves on *In vitro* Fibroblast Cells

Rani Wulan Sari, Natallia Pranata<sup>1</sup>, Vinna Kurniawati Sugiaman<sup>1</sup>

Faculty of Dentistry,  
Maranatha Christian  
University, <sup>1</sup>Department of  
Oral Biology, Maranatha  
Christian University,  
Bandung, Indonesia

Received : 22-04-19  
Revised : 15-08-19  
Accepted : 04-09-19  
Published Online: 14-10-19

### ABSTRACT

**Background:** Tooth extraction is the most frequently conducted dental care procedure. In Indonesia, there is an extremely high utilization of dental and oral health services for tooth extraction, reaching 79.6%. Pain is a side effect of extraction. Pain due to extraction wounds can be treated with analgesic drugs, but alternative drugs with minimal or no side effects are still being researched. An herbal active *beluntas* leaf substance can reduce pain from extraction wounds. The *beluntas* plant not only aids in healing wounds but also exhibits anti-inflammatory and antipyretic effects. **Objectives:** In this study, the aims were to determine the 50% inhibitory concentration (IC<sub>50</sub> value) and examine the viability effect of an ethanol extract of *beluntas* leaves on fibroblast cell cultures *in vitro*. **Methods:** Laboratory experiments were carried out. *Beluntas* leaves were obtained; their leaf extracts were prepared using ethanol as the solvent; phytochemical tests were performed. Triplicate measurements for the viability of 3T3 BALB/c fibroblast cells were carried out using the Methylthiazol sulfophenyl (MTS Assay) method. The extract concentrations were 500, 250, 125, 62.50, 31.25, 15.63, and 7.81 µg/mL. **Results:** Data analysis was carried out by one-way analysis of variance statistical test. Analysis results revealed that extract concentrations of 500, 31.25, 15.63, and 7.81 µg/mL exhibit a significant difference in the effect of cytotoxicity ( $P < 0.05$ ) on fibroblast cells, and the IC<sub>50</sub> value is 265.388 µg/mL. **Conclusion:** A significant difference in the cytotoxicity effect between the concentrations of the ethanol extract of *beluntas* (*P. indica*) leaves on the fibroblast cell cultures *in vitro* was observed. The *beluntas* leaf extract at an IC<sub>50</sub> value of 7.81 µg/mL did not affect cell viability; hence, it is considered safe.

**KEYWORDS:** *Beluntas* leaves, cytotoxicity, ethanol extract, fibroblast cells, MTS assay, viability

## BACKGROUND

Tooth extraction is common in dental practice. In Indonesia, there is extremely high utilization of dental and oral health services for tooth extraction, reaching 79.6%.<sup>1</sup> Dental extraction is a surgical procedure that deals with soft and hard tissues in the oral cavity.<sup>2</sup> Pain is a side effect of extraction. Pain starts to disappear during the proliferation phase in the proliferation process of fibroblast cells. The healing process of extraction occurs in several phases, viz., inflammatory phase, proliferation phase (complete wound closure and epithelial formation occur in this phase), and maturation.

The wound-healing process is strongly affected by the migration and proliferation of fibroblasts in the wound area, and fibroblasts play a key role in the repair process.<sup>3</sup>

Pain due to extraction wounds can be treated using analgesic drugs, but alternative drugs with minimal or no side effects are remained unclear. Flavonoids are herbal

**Address for correspondence:** Dr. Vinna Kurniawati Sugiaman,  
Department of Oral Biology, Maranatha Christian University,  
Bandung, Indonesia.  
E-mail: [vinnakurniawati@yahoo.co.id](mailto:vinnakurniawati@yahoo.co.id)

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**For reprints contact:** [reprints@medknow.com](mailto:reprints@medknow.com)

**How to cite this article:** Sari RW, Pranata N, Sugiaman VK. Viability test of ethanol extract of *beluntas* (*Pluchea indica*) leaves on *in vitro* fibroblast cells. *Sci Dent J* 2019;3:90-4.

### Access this article online

#### Quick Response Code:



**Website:** [www.scidentj.com](http://www.scidentj.com)

**DOI:** 10.4103/SDJ.SDJ\_18\_19

active substances that can reduce pain caused by extraction wounds. Besides flavonoids, *beluntas* leaves also contain other active compounds that can accelerate wound healing, including alkaloids, saponins, polyphenols, tannins, sterols, sodium, citrus oils, amino acids, fats, calcium, magnesium, phosphorus, Vitamin A, and Vitamin C. The *beluntas* plant can not only aid in wound healing but also exhibit anti-inflammatory and antipyretic effects.<sup>4,5</sup>

Flavonoids can stop the production of prostaglandin; therefore, it reduces pain, renders antibacterial effects, functions as an anti-inflammatory agent, and exhibits a working mechanism for the inhibition of lipid peroxidation, which serves to reduce reactive oxygen species; hence, it can slow tissue death, increase vascularity and collagen, prevent cell damage, and increase DNA synthesis.<sup>6</sup>

Tannins also exhibit an effect that can stimulate not only the production of fibroblast cells but also the formation of collagen tissues in wound healing. Tannins exhibit antimicrobial and antioxidant effects that can help prevent infections and fight free radicals; therefore, tannins accelerate the wound-healing process.<sup>7</sup>

*Beluntas* leaves can be used as therapeutic agents for oral mucous tissue damage. To serve as therapeutic agents, *beluntas* leaves must satisfy the requirements for biocompatibility, i.e., must not cause irritation and toxicity to the body. Therefore, it is imperative to conduct a standardized study that can determine the effect of the *beluntas* leaves extract on cell viability involved in tissue healing, including fibroblast cells.

In this study, the 50% inhibitory concentration (IC<sub>50</sub>) value was determined, as well as the cytotoxicity effect of the ethanol extract of *beluntas* (*P. indica*) leaves on fibroblast cell cultures *in vitro* was determined.

## MATERIALS AND METHODS

### Ethanol extract of *beluntas*

A  $\pm$  10-year-old *beluntas* leaf sample was obtained from the experimental garden in Manoko Spice and Medicinal Plant Research Institute (Balitro), which was determined in the laboratory of identification and determination of the Faculty of Life Sciences, Institut Teknologi Bandung (ITB).

A crude *beluntas* leaf extract was prepared by the maceration method using 70% ethanol, which was carried out at the Aretha Medika Utama, Biomolecular and Biomedical Research Center.

The dilution of the ethanol extract of the *beluntas* leaf started with the preparation of a stock solution, which had a concentration of 500  $\mu$ g/mL, followed by

the preparation of a series of working solutions with concentrations of 250, 125, 62.50, 31.25, 15.63, and 7.81  $\mu$ g/mL.

### Phytochemical test

Phytochemical analysis by the Farnsworth method was carried out to identify the chemical groups of alkaloids, saponins, tannins, flavonoids, terpenoids, phenols, and steroids/triterpenoids.<sup>8</sup>

### MTS assay

The MTS assay was performed to measure the viability of fibroblast cells, which was based on the conversion of tetrazolium salts to colored formazan by the mitochondrial activity of living cells. The amount of the produced formazan depends on the number of cells that are feasible in the culture, which is measured in triplicate by a spectrophotometer at 490 nm, with 24 h of incubation.<sup>9</sup>

The viability test of the *beluntas* leaf extract was carried out using confluent 70%–80% 3T3 BALB/c fibroblast cells and planted with a density of 5000 cells/well in 96 wells/plate using the cell culture medium of the Dulbecco's modified Eagle medium, 10% fetal bovine serum, and 1% antibiotic-antimycotic solution, followed by incubation for 24 h in an incubator at a temperature of 37°C and a CO<sub>2</sub> level of 5%.

The old medium was removed and replaced with 200  $\mu$ L of a new medium, and 20  $\mu$ L of the extract with various series of concentrations was added to each well, followed by incubation for 24 h with a temperature of 37°C and a CO<sub>2</sub> content of 5%. After 24 h, 20  $\mu$ L of the MTS reagent was added to each well and incubated for 3 h at the same temperature and CO<sub>2</sub> levels, and the cells were calculated on the basis of their absorbance and curve integration. Based on the number of viable cells, the cytotoxicity of a material can be classified [Table 1].

IC<sub>50</sub> was calculated from the plot of concentration as a function of the percentage of viability.<sup>10</sup>

$$\% \text{ Viability} = \frac{\text{Absorbance of the sample}}{\text{Absorbance of control}} \times 100\%$$

$$\text{IC}_{50} = \frac{\text{Absorbance of control} - \text{absorbance of the sample}}{\text{Absorbance of control}} \times 100\%$$

The obtained data were first processed by a normality test, i.e., the Kolmogorov–Smirnov test. Second, one-way analysis of variance with  $\alpha = 0.05$  and an advanced test, i.e., the *post hoc* test (Tukey), were carried out. Finally, probit analysis was performed to process the IC<sub>50</sub> data.

## RESULTS

Phytochemical test results of the *beluntas* leaf extract revealed the presence of alkaloids, saponins, flavonoids, phenols, terpenoids, steroids, and tannins [Table 2]. These results are in agreement with those reported previously.<sup>11</sup>

Table 2 and Figure 1 summarize the results obtained from the viability test. Based on the classification of cell viability, the table data revealed that concentrations of 500, 250, and 125 µg/mL are classified as extremely toxic. By contrast, lower concentrations of 62.5, 31.25, and 15.63 µg/mL are classified as slightly toxic. A concentration of 7.81 µg/mL is classified as nontoxic.

Then, the probit  $IC_{50}$  test was performed, and the  $IC_{50}$  value (safe concentration) of the ethanol extract of the *beluntas* was 265.388 µg/mL [Table 3 and Figure 1].<sup>[10]</sup>

**Table 1: Classification of material cytotoxicity**

Percentage of viability cells	Classification
>90% of the cells are viable	Nontoxic
60%-90% of the cells are viable	Slightly toxic
30%-59% of the cells are viable	Quite toxic
<30% of cells are viable	Extremely toxic

**Table 2: Phytochemical examination results of *beluntas* leaves**

Phytochemical compounds	Results
Flavonoids	+
Saponins	+
Phenols	+
Tannins	+
Steroids/triterpenoids	+
Terpenoids	+
Alkaloids	+

+: Present active compound

**Table 3: Viability tests for the 3T3 BALB/c fibroblast cells**

Concentration (µg/mL)	Absorbance	Average		Number of viable cells
		Viability (%)	Inhibition (%)	
Cell control	2.185	100	0.00	5000
DMSO control	1.939	97.23	2.77	4.861
7.81	1.934	91.16±0.95	8.84	4.558
15.63	1.843	79.01±1.72	20.99	3.950
31.25	1.945	65.37±2.55	34.63	3.268
62.50	2.025	59.18±1.13	40.82	2.959
125	2.145	57.18±2.38	42.82	2.859
250	2.234	55.68±1.84	44.32	2.784
500	2.348	45.53±2.04	54.47	2.276

DMSO: Dimethyl sulfoxide

## DISCUSSION

Healing occurs in several phases, viz., homeostasis, inflammation, proliferation, and maturation. The wound-healing process is strongly affected by the migration and proliferation of the fibroblasts in the wound area, and fibroblasts play a key role in the repair process.<sup>12,13</sup>


Flavonoids in the *beluntas* leaves play a role in the anti-inflammatory process because it can shorten the inflammation time; therefore, proliferation can occur immediately and inhibit bleeding. The flavonoid activity can accelerate the wound-healing process, and it is supported by antioxidant mechanisms for the inhibition of the free radical activity.

Antioxidants can block the initiation of free radical and trigger the proliferation of fibroblast cells. Besides flavonoids, *beluntas* leaves also contain saponins, which can not only stimulate collagen formation but also increase the fibroblast density via the activation of TGF-β.<sup>4,14,15</sup>

Phenols, alkaloids, and tannins can serve as antibacterial agents, and terpenoids are active ingredients that help to accelerate the formation of collagen fiber produced by fibroblast cells.<sup>14</sup>

Hence, to develop materials for the natural treatment of postextraction wounds, materials must be subjected first to biocompatibility tests in accordance with the material requirements in dentistry, especially those used in the mouth. One test to determine the various properties of a dental material is a cytotoxicity test on tissues. To determine the cytotoxicity of the ethanol extract of the *beluntas* leaves (*P. indica*), a fibroblast cell was tested using the MTS assay.

Parameters for the cytotoxicity test are based on the absorbance value, i.e., if the cell color becomes thicker (purple), the absorbance value is higher, implying that more cells are alive; however, if the cell



**Figure 1: Relationship between the concentrations of the *beluntas* leaf extract on the viability of 3T3 BALB/c fibroblast cell**

color fades, the absorbance value is lower, implying that several cells die.<sup>16</sup>

The cell viability decreased after the administration of the extracts to cells, and significant differences between concentrations were related to the differences in the cell response to the concentration, including the number of active ingredients at different concentrations, indicating that the extract exhibits cytotoxic properties on fibroblast cells. The mechanism and intensity of cell death depend on material content or cell contact. *Beluntas* leaves contain eugenol, which is a phenol derivative.<sup>17</sup> Eugenol can cause cell cytotoxicity, which can damage the protein structure by a number of physical and chemical elements.<sup>18</sup>

Cells exposed to materials or extracts exceeding the peak of exposure cause cell death. It can cause toxicity to cells via different mechanisms, such as the destruction of cell membranes; prevention of protein synthesis; irreversible binding to receptors; inhibition of the polydeoxynucleotide elongation; and other enzymatic reactions.<sup>9,18,19</sup>

In a particular cell, cytotoxic agents can also be metabolized without any observable effect although most cells experience necrosis when confronted with toxic compounds. Necrosis can occur when cells are exposed to conditions that are extremely different from their physiological conditions or when the compound content of the extracts can damage cell membranes. Necrosis starts from the disruption of the cell's ability to maintain homeostasis because it can cause the entry of water and extracellular ions. Intracellular organelles, especially mitochondria and all other cells, can swell and rupture (cell lysis).<sup>20</sup>

As a result of the disruption of the plasma membrane, the cytoplasmic contents, including the lysosome enzyme, are released into the extracellular fluid. The activity of these enzymes in extracellular media can be used to determine the level of necrosis. Besides these factors, the dose, exposure duration, and mechanism of cytotoxic agents are other factors that can cause cell death.<sup>9,21,22</sup>

The IC<sub>50</sub> value for the ethanol extract of *beluntas* leaves on fibroblast cells is 265.388 µg/mL, which is categorized as safe because it does not interfere with the viability of fibroblast cells; however, further research is required to investigate the effect of the *beluntas* leaves extract on the proliferation of fibroblast cells.

## CONCLUSION

A significant difference in the cytotoxicity effect between the concentrations of the ethanol extract of

*beluntas* (*P. indica*) leaves on fibroblast cell cultures *in vitro* was observed. The *beluntas* leaf extract at an IC<sub>50</sub> concentration of 7.81 µg/mL did not affect cell viability; hence, it is considered safe.

## Acknowledgment

We would like to acknowledge the Aretha Medika Utama, Biomolecular and Biomedical Research Center Laboratory.

## Financial support and sponsorship

Nil.

## Conflict of interest

There are no conflicts of interest.

## REFERENCES

1. Data Center and Information Ministry of Health of the Republic of Indonesia. [Dental and Oral Health Situation]. Jakarta; 2014.
2. Widiyastomo W, Andari WK, Permatasari I. Effect of starfruit fruit juice (*Averrhoa carambola* Linn.) On increasing the number of fibroblasts in Wistar mouse socket after tooth socket. *Prodenta J Dent* 2013;1:62-70.
3. Kartikasari ID, Soelistiono, Prihartiningsih., Effect of *Salvadora persica* Stem Extract on the Growth of *Streptococcus α-haemolyticus* Bacteria after Isolation of Mandibular Third Molar Tooth Extract (*in vitro* study). *Kedokt Gigi Univ Gajah Mada*. 2008;1-6. FKG UGM 2008;1-6.
4. Hafsari AR, CahyantoT, Sujarwo T, Lestari RI. Antibacterial activity test of *beluntas* (*Pluchea indica* (L.) Less.) extract against *Propionibacterium acnes* cause of acne. *J ISTEK* 2015;9:141-61.
5. Ibrahim N', Wong SK, Mohamed IN, Mohamed N, Chin KY, Ima-Nirwana S, et al. Wound healing properties of selected natural products. *Int J Environ Res Public Health* 2018;15. pii: E2360.
6. Suranto A. The Enormity of Propolis to Cure Disease. Jakarta: Agro Media Pustaka; 2010.
7. Ahmed KA, Abdulla MA, Mahmoud FM. Leaf extract in experimental rats. *Middle East J Sci Res* 2012;11:1614-8.
8. Widowati W, Mozef T, Risdian C, Yellianty Y. Anticancer and free radical scavenging potency of *Catharanthus roseus*, *Dendrophthoe petandra*, *Piper betle* and *Curcuma mangga* extracts in breast cancer cell lines. *Oxid Antioxid Med Sci* 2013;2:137-42.
9. Miret S, De Groene EM, Klaffke W. Comparison of *in vitro* assays of cellular toxicity in the human hepatic cell line HepG2. *J Biomol Screen* 2006;11:184-93.
10. Heravi F, Ramezani M, Poosti M, Hosseini M, Shajiei A, Ahrari F, et al. *In vitro* cytotoxicity assessment of an orthodontic composite containing titanium-dioxide nano-particles. *J Dent Res Dent Clin Dent Prospects* 2013;7:192-8.
11. Widyawati PS, Wijaya CH, Hardjosworo PS, Sajuthi D. Evaluation of antioxidant activity of *beluntas* (*Pluchea indica*) leaf extract based on differences in leaf segments. *Rekapangan J Teknol Pangan* 2011;5:1-17.
12. Li B, Wang JH. Fibroblasts and myofibroblasts in wound healing: Force generation and measurement. *J Tissue Viability* 2011;20:108-20.
13. Gonzalez AC, Costa TF, Andrade ZA, Medrado AR. Wound healing – A literature review. *An Bras Dermatol* 2016;91:614-20.



14. Muslihah K, Sumono A, Fatmawati DW. Cytotoxicity effect of pectin extract from *Coffea robusta* (*Coffea acenephora*) fruit peels on human dental pulp fibroblast cell. *Pustaka Kesehatan* 2018;6:173-8.
15. Laksmiawati DR, Prasanti AP, Larasinta N, Syauta GA, Hilda R, Ramadaniati HU, *et al.* Anti-Inflammatory potential of *Gendarussa* (*Gendarussa vulgaris* Nees) and *Soursoup* (*Annona muricata* L) extracts in LPS stimulated-macrophage cell (RAW264.7). *J Nat Remedies* 2016;16:73-81.
16. Bahuguna A, Khan I, Bajpai VK, Kang SC. MTT assay to evaluate the cytotoxic potential of drug. *Bangladesh J Pharmacol* 2017;12:115-8.
17. Gunasekaran S, Vinot KT, Lakshamana SS, Suganya P, Rincy Y, Amrutha C. Screening of *in vitro* cytotoxic activity of brown seaweeds against hepatocellular carcinoma. *J Appl Pharm Sci* 2017;7:51-60.
18. Nejad SM, Ozgunes H, Basaran N. Pharmacological and toxicological properties of eugenol. *Turk J Pharm Sci* 2017;14:201-6.
19. Susanty A, Rimayanti R, Sukmanadi M. Antibacterial activity of the ethanol extract *Pluchea indica* less leaves against *Escherichia coli* by *in vitro*. *Veterinaria Medika* 2008;1:29-32.
20. Galluzzi L, Vitale I, Aaronson SA, Abrams JM, Adam D, Agostinis P, *et al.* Molecular mechanisms of cell death: Recommendations of the nomenclature committee on cell death 2018. *Cell Death Differ* 2018;25:486-541.
21. Aslantürk OS. *In vitro* cytotoxicity and cell viability assay: Principles, advantages, and disadvantages. In: Larramendy ML, Soloneski S, editors. *Genotoxicity-A Predictable Risk to Our Actual World*. London: Intech Open; 2018. p. 1-18.
22. Van Tonder JJ. Development of an *In vitro* Mechanistic Toxicity Screening Model Using Cultured Hepatocytes [PhD Thesis]. Pretoria: University of Pretoria; 2011.