


Biomedical Engineering


ISSN 2356-2471

BE

<http://be.ub.ac.id>



Smart Molecule of Natural
SMONAGENS
UB
Genetics Resource





- Home
- About
- Login
- Register
- Search
- Current
- Archives
- Announcements
- Editorial Board
- Focus and Scopes
- Author Guidelines

Home > About the Journal > Editorial Team

Editorial Team

Editors

1. Rista Nikmatu Rohmah, Institut Biosains Universitas Brawijaya Malang
2. mrs Lidwina Faraline, Indonesia
3. Prof Fatchiyah Fatchiyah, Brawijaya University Malang, Indonesia, Indonesia
4. Antonius Christianto

Section Editors

1. Rista Nikmatu Rohmah, Institut Biosains Universitas Brawijaya Malang
2. Mrs. Atika Hanoum Rahasta, University of Brawijaya, Indonesia

Open Journal Systems

Journal Help

User

Username

Password

Remember me

Login

Notifications

- View
- Subscribe

Journal Content

Search

Search Scope All

Search

Browse

- By Issue
- By Author
- By Title

Information

- For Readers
- For Authors
- For Librarians


[Home](#) > [Archives](#) > [Vol 1, No 1 \(2015\)](#)

Vol 1, No 1 (2015)

BIOMEDICAL ENGINEERING

The first volume of Biomedical Engineering journal published in this year. Contains 10 scientific journals, with new discoveries and analysis. Bring to you with a intention and hope to deliver a good science work that hopefully could help people.

Full Issue

View or download the full issue

[Vol1_No1_2015](#)

Table of Contents

Articles

BIOMEDICAL ENGINEERING <i>Biomedical Engineering</i>	57
BIOMEDICAL ENGINEERING Vol.1 No.1 Tahun 2015 <i>Biomedical Engineering</i>	57
Phytochemical assay, potential of antimalarial and antioxidant activities of green tea extract and its fraction <i>Fanny Rahardja</i>	7
The effects of regular exercise on the number of osteoclasts in ovariectomized Sprague dawley rat mandible <i>Rini Maya Puspita</i>	5
Antioxidant and hypoglycemic activities of extract and fractions of Rambutan seeds (<i>Nephelium lappaceum</i> L.) <i>Sylvia Soeng</i>	6
CD35 (DEL C) frameshift mutation in exon 2 of β -globin gene on β -thalassemia carriers <i>Niken Satuli Nur Handayani</i>	5
Antioxidant properties of spice extracts <i>Wahyu Widowati</i>	6
Carbonated hydroxyapatite inflammation's responses on local rabbits: study of neutrophils's cell count, macrophages, and edema volumes on mandible <i>Aryan Marita</i>	5
Investigation of testicular germ cell tumor on Cryptorchidism with Hypospadias and Ambiguous Genitalia <i>Amalia setyawati</i>	5
Effect of Folic Acid and Vitamin B12 Supplementation on Homocysteine Level in association with MTHFR C677T Polymorphisms in Overweight Female Adults <i>Fidella Fiddelia</i>	6
Protein extract of Detam 1 soybean contains β -conglycinin stimulating cholecystokinin signal transduction through activities of PKC, c-RAF and ERK 1/2 in Wistar rats <i>Melinah Hidayat</i>	7
Gene p53 Mutations after the Induction of 7,12-Dimethylbenz(a)anthracene (DMBA) and Administration of Anti-Carcinogenesis Properties of <i>Gynura procumbens</i> in Sprague Dawley Rats <i>abdul Gofur</i>	5
Phytochemical assay, potential of antimalarial and antioxidant activities of green tea extract and its fraction <i>Fanny Rahardja</i>	1-6

Open Journal Systems

Journal Help

User

Username

Password

Remember me

[Login](#)

Notifications

- View
- Subscribe

Journal Content

Search

Search Scope

[Search](#)

- Browse
- By Issue
 - By Author
 - By Title

Information

- For Readers
- For Authors
- For Librarians



Antioxidant properties of spice extracts

Wahyu Widowati^{1†}, Hana Ratnawati¹, Winsa Husin¹, Maesaroh Maesaroh²

¹Faculty of Medicine, Maranatha Christian University, Bandung, West Java, Indonesia

²Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung, West Java, Indonesia

ARTICLE INFO

Article history:

Received: January 2015

Accepted: June 2015

Available online:

August 2015

Keywords:

DPPH

Free radicals

Spice

SOD

Total phenolic content

[†]Corresponding author:

Faculty of Medicine,
Maranatha Christian
University, Jl. Prof. drg.
Suria Sumantri no 65
Bandung 40164, Indonesia
wahyu_w60@yahoo.com

ABSTRACT

Objective : This study was conducted to determine the antioxidant activities of spice extracts including clove (*Syzygium aromaticum* L.), Indonesian cassia (*Cinnamomum burmanni* (C. Nees & T. Ness)), coriander (*Coriandrum sativum* L), nutmeg (*Myristica fragrans* Houtt), java cardamom (*Amomum compactum* Soland. Ex maton).

Methods : This research was to evaluate antioxidant activities including 1,1-diphenyl-2-picrylhydrazyl (DPPH) scavenging activity, the value of superoxide dismutase (SOD), and total phenolic content.

Result : The highest DPPH activity is clove and Indonesian cassia extracts with IC₅₀ value 4.16 µg/ml and 5.46 µg/ml respectively. The highest SOD value are Indonesian cassia extract (9.1432 U/ml) at 500 µg/ml, 7.0045 U/ml at 125 µg/ml and 4.6751 U/ml at 31.25 µg/ml. Clove extract was the highest of phenolic content (188.35 µg/mg eugenol equivalent).

Conclusion : Indonesian cassia extract have high antioxidant activities both DPPH scavenging and SOD activities. Clove extract contain the highest eugenol compared with Indonesian cassia, coriander, nutmeg and java cardamom.

1. Introduction

Non communicable diseases (NCD) mainly cardiovascular diseases (CVD), cancers, diabetes mellitus (DM) and chronic respiratory diseases are leading threat to human health. These four diseases are the world's biggest killers, causing an estimated 35 million deaths each year - 60% of all death globally. WHO projects that, globally, NCD deaths will increase by 17% over the next ten years [1]. The free radicals formation is co-related with the pathological conditions including CVD, cancer, neurological, age-related disorders, degenerative disorder [2, 3]. Free radical will attack the DNA, lipid and proteins thus leading to chronic disease [4, 3]. Oxidative stress is an imbalance condition between free radicals and antioxidants, to reduce oxidative stress or high level of free radicals including Reactive Oxygen Species (ROS) and Reactive Nitrogen

Species(RNS) can be combated with the involvement of antioxidants both exogenous and endogenous [5].

Spices commonly known as food adjuncts, but they can be used as traditional medicines too [6, 7]. Species such as clove, cinnamon, rosemary, paprika, nutmeg, ginger, garlic, mustard, etc., have antioxidant activities [8]. Antioxidant activity is associated with their phenolic compounds activities to scavenge free radical, transition-metal-chelating, have capacity to quenching singlet-oxygen, and their redox properties [9, 10, 11, 12]. Activities of spices to scavenge ROS and RNS are considered as effective remedies for diabetes [13]. Spices have capability to detoxify the liver, to be anti-inflammatory agent, and infectious protector agent [14].

The aims of this study were to determine antioxidant activities of spices extract including clove, Indonesian cassia, coriander, nutmeg, java

cardamom, and to evaluate the total phenolic content.

2. Materials and Methods

2.1 Plant material

Plant material were collected from several plantation in Indonesia : clove (*Syzygium aromaticum* L.) from Cimahi-Bandung-west Java, Indonesian cassia (*Cinnamomum burmanni* (C. Nees& T. Ness) from Lembang-Bandung- West Java, Indonesia, coriander (*Coriandrum sativum* L) from Cipanas-Cianjur- West Java, nutmeg (*Myristica fragrans* Houtt) from Cicurug-Sukabumi- West Java, java cardamom (*Amomum compactum* Soland. Ex maton) from Cicurug-Sukabumi- West Java. The plants were identified by staff at the herbarium, Department of Biology, School of Life Sciences and Technology, Bandung Institute of Technology, Bandung, West Java, Indonesia. Seed of coriander, nutmeg, java cardamom and skin of bark of Indonesian cassia and flower of clove were chopped and dried in a dry tunnel (40-45°C) to achieve a stable water level (11% water content), then chopped finely in a blender, producing a 60 mesh size flour [15, 16].

2.2 Extract preparation

The dried flour of samples (400 g) were immersed in 70% ethanol. After 24 h, the filtrates were collected and the residues were immersed again in 70% ethanol for 24 h. The filtrates were evaporated with a rotary evaporator at 40°C, yielded clove extract 13.29%; java cardamom extract 9.71%, Indonesian cassia extract 17.92%, coriander extract 8.80% and nutmeg extract 11.64% [15, 16]. The extracts were stored at 4°C. The ethanol extracts of clove, Indonesian cassia, coriander, nutmeg, java cardamom were dissolved in HPLC-grade methanol (Merck) for DPPH scavenger and phenolic content assay and were dissolved in 10% dimethylsulfoxide (DMSO; Merck) for SOD assay [15, 16]. Pure compounds were used as comparison of antioxidant activities including limonin (Biophurify Phytochemical Ltd.), coumarin (Biophurify Phytochemicals Ltd.), trans-cinnamic acid (Sigma-Aldrich), thymoquinone (Sigma-Aldrich) and cardamom oil (Sigma-Aldrich),

2.3 DPPH scavenging activity assay

Amount of 50 µl extracts and limonin, coumarin, trans-cinnamic acid, thymoquinone

and cardamom oil were introduced into a microplate followed by 200 µl DPPH (Sigma-Aldrich) solution (0.077 mmol/l in methanol). The mixtures was shaken vigorously and kept in the dark room for 30 min at room temperature; DPPH scavenging activity was determined with a microplate reader at 517 nm [15, 16, 17]. The radical scavenging activity of each sample was measured

$$\text{DPPH scavenging \%} = 1 - \frac{\text{Sample absorbance}}{\text{Negative control absorbance}} \times 100$$

according to the following formula:

2.4 Superoxide Dismutase (SOD) Assay

The SOD assay was performed with SOD assay kit (Cayman) including assay buffer, sample buffer, radical detector, SOD standard, and xanthine oxidase. SOD standards were prepared by introducing 200 µl diluted radical detector and 10 µl SOD standard (7-level standard) per well. Sample wells contained 200 µl of the diluted radical detector and 10 µl of the sample. The reaction was initiated by adding 20 µl of diluted radical detector to all wells. The samples consist of five extracts and five compounds were measured at 3 concentrations in triplicate. The mixtures were shaken carefully for few seconds, incubated for 20 minutes at room temperature, SOD activity was measured on a microplate reader (Multi Go Skan) at 440-460 nm. The linearized SOD standard rate and SOD activity were calculated using the equation which obtained from linear regression of the standard. One unit is defined as the amount of enzyme to yield 50% dismutation of the superoxide radical [16]. Furthermore, standard curves were constructed based on the value of the LR and SOD value be calculated:

The calculation is done by calculating the value of SOD linearized rate/LR (LR Std A= Abs Std A/Abs Std A: LR Std B= Abs Std A/Abs Std B)

$$\text{SOD (U/ml)} = \left[\frac{\text{Sample LR-y intercept}}{\text{Slope}} \right] \times 0,23\text{ml}/0,01\text{ml} \times \text{Sample dilution}$$

2.5 Total phenolic content assay

Total phenolic content was measured according to the Folin-Ciocalteu method. Samples (15 µl) were introduced into microplate; 75 µl of Folin-Ciocalteu's reagent (2.0 M) and 60 µl of sodium carbonate (7.5%) were added. The samples were mixed and incubated at 45°C for 15 min [18,19]. Subsequently, absorbance value was measured at

760 nm using microplate reader (Multi Skan Go). Total phenolic content expressed as Eugenol equivalent was calculated by the following formula:

$$C = \frac{c \times V}{m}$$

C: total content of phenolic compounds ($\mu\text{g}/\text{mg}$) of clove, Indonesian cassia, coriander, nutmeg, java cardamom extract in eugenol equivalent;
c: the concentration of eugenol established from the calibration curve, $\mu\text{g}/\text{ml}$;
V: the volume of extract (ml); m: the weight of spices extract (mg).

Table 1. DPPH scavenging activity (IC_{50}) of spice extracts. The DPPH scavenging activity assay was measured triplicate for each sample. Linear equations, coefficient of regression (R^2), and IC_{50} were calculated.

Samples	Linear equation	R^2	IC_{50} ($\mu\text{g}/\text{ml}$)
Java cardamom extract	$Y=0.2071X+13.157$	0.8797	177.90
Indonesian cassia extract	$Y=6.878X+12.448$	0.918	5.46
Coriander extract	$Y=0.4873X+2.011$	0.9850	98.48
Clove extract	$Y=10.342X+7.006$	0.9825	4.16
Nutmeg extract	$Y=0.861x+16.058$	0.878	39.422
Limonin	$Y=0.1181X+7.4917$	0.8631	359.93
Coumarin	$Y=0.0404X+2.7142$	0.8387	1170.44
Trans-cinammic acid	$Y=0.066X+15.578$	0.8849	521.55
Thymoquinone	$Y=5.7709X+22.53$	0.9471	4.76
Cardamom oil	$Y=0.2183X-20.577$	0.8472	323.30

standard. DPPH accept a proton from an antioxidant becomes 1,1-diphenyl-2-picrylhydrazine which is a stable radical [17]. The DPPH free radical scavenging activity of clove, Indonesian cassia, coriander, nutmeg, java cardamom extracts compared to compounds of spice including limonin, coumarin, trans-cinnamic acid, thymoquinone and cardamom oil. The IC_{50} is the concentration of antioxidant needed to scavenge 50% of the DPPH free radicals [15].

Based on the IC_{50} values of DPPH scavenging activity (Table 1) showed that clove extract was the most active antioxidant based on DPPH scavenging activity and the lowest was coumarin.

3.2 The SOD activity

Total phenol value was obtained from the regression equation for eugenol :

$$Y=0.0024 X + 0.1228, \text{ with } R^2 =0.94$$

3. Result

3.1 DPPH scavenging activity

The antioxidant activities were measured including DPPH scavenging activity, SOD activity and phenolic content using eugenol as

Superoxide dismutase (SOD) as an endogenous antioxidant activity would be increased by changing the superoxide anion ($\text{O}_2^{\cdot-}$) into hydrogen peroxide (H_2O_2) and oxygen (O_2) [20]. The SOD activity of extracts and compounds of spices can be seen in Table 2.

Based on the results (Table 2.) showed that the highest SOD value at a concentration of 500 $\mu\text{g}/\text{ml}$, 31.25 $\mu\text{g}/\text{ml}$ and 31.25 $\mu\text{g}/\text{ml}$ was the Indonesian casia extract 9.1432 U/ml, 7.0045 U/ml and 4.6751 U/ml respectively.

3.3 Total phenol antioxidant activity assay

Phenols have the ideal chemical structure that can scavenge free radicals, so this class of phenolic compounds exhibit higher antioxidant activity than tocopherols and ascorbate [21]. The

total phenolic compound of spice extracts in Eugenol equivalent can be seen in Table 3.

Based on the test results of phenolic compound total (Table. 3) showed that the

highest phenolic compound was clove extract contained eugenol 188.35 $\mu\text{g/g}$.




Figure 1. DPPH scavenging activity of spice extracts diluted in methanol to achieve the final concentrations 100; 50; 25; 12.5; 6.25; 3.125; 1.563; 0.781; 0.391; 0.190 $\mu\text{g/ml}$.

4. Discussion

Based on Table 1,2, and 3 showed that spice extracts have high antioxidant activities and Indonesian cassia extract was the highest SOD value. The most active DPPH scavenging activity was clove extract. The highest phenolic content was clove extract. In previous study, among 30 spices from subtropical country Ireland, clove also had highest antioxidant capacity [8]

Many spices contained high levels of phenolics and demonstrated high antioxidant capacity [21]. The spices and related families with the highest antioxidant capacity were screened, e.g., clove, cinnamon or Indonesian cassia representing potential sources of potent natural antioxidants for commercial exploitation [21]. Clove extract contained very high levels of phenolics (188.35 $\mu\text{g/mg}$ eugenol equivalent). Other spices with high levels of phenolics were cardamom extract

Table 2. Mean, standard deviation and Tukey HSD post hoc test of extracts and compounds of spices SOD activity (U/ml) were measured in triplicate for sample. (Linear equation, coefficient of regression (R^2) of SOD standard and SOD activity of extracts and compounds were calculated.)

Samples	Concentrations ($\mu\text{g/ml}$)		
	500	125	31,25
Java cardamom extract	2.6903±0.1890 ^e	0.9022±0.2242 ^b	0.3178±0.1158 ^a
Indonesian cassia extract	9.1432±0.7559 ^g	7.0045±0.2821 ^f	4.6751±0.1732 ^f
Coriander extract	1.8448±0.0393 ^{cd}	1.4919±0.0729 ^{cd}	0.9747±0.0587 ^b
Clove extract	4.5607±0.2502 ^f	3.6723±0.0644 ^e	2.7509±0.0653 ^e
Nutmeg extract	4.3044±0.1368 ^f	3.5462±0.0287 ^e	2.3532±0.0765 ^d
Limonin	0.5615±0.0619 ^a	0.3930±0.0415 ^a	0.2554±0.0350 ^a
Coumarine	0.6960±0.0677 ^{ab}	0.4251±0.0359 ^a	0.3423±0.0069 ^a
Trans-cinnamic acid	2.3707±0.0253 ^{dc}	1.7197±0.0859 ^d	1.3663±0.1180 ^c
Thymoquinone	1.4034±0.0452 ^{bc}	1.2249±0.0143 ^{bc}	0.9961±0.0181 ^b

Cardamom oil	1.1368±0.0777 ^{abc}	0.9999±0.0479 ^b	0.8883±0.0373 ^b
--------------	------------------------------	----------------------------	----------------------------

Table 3. Mean total phenolic compound in spice extracts (Eugenol equivalent/mg)

Sample	Eugenol/ml			Eugenol µg/mg			Average
Clove extract	94.92	92.04	96.29	188.4	184.08	192.58	188.35
Cardamom extract	67.63	67.54	66.21	135.26	135.08	132.42	134.25
Indonesian cassia extract	55.04	69.54	56.88	110.08	139.08	113.76	120.97
Coriander extract	-	-	-	-	-	-	-
Nutmeg extract	47.08	51.04	48.33	94.16	102.08	96.66	97.63

(134.25 µg/mg eugenol equivalent) and nutmeg contained relatively low phenolics (97.63 µg/mg eugenol equivalent). Previous studies also showed that clove had a very strong antioxidant activity and a high level of phenolics [22, 23]. The various antioxidant mechanisms of clove extracts were attributed to a strong hydrogen-donating ability, a metal chelating ability, and their effectiveness as good scavengers of hydrogen peroxide, superoxide, and free radicals [8]. Our results showed that the clove extract was the most powerful phenolic antioxidant and exhibited the strongest radical scavenging activity among five spice extracts.

Spices, like vegetables, fruits, and medicinal herbs, are known to possess a variety of antioxidant effects and properties [24]. The antioxidant effect of phenolic compounds is mainly due to their redox properties and is the result of various possible mechanisms: free-radical scavenging activity, transition-metal-chelating activity, and/or singlet-oxygen-quenching capacity [11, 21]. These multiple potential mechanisms of antioxidant action make the diverse group of phenolic compounds an interesting target in the search for health-beneficial phytochemicals and also offer a possibility to use phenolic compounds or extracts rich in them in lipid-rich foods to extend shelf life [25]. The presence of antioxidative and antimicrobial of phenolic compounds in many spices gives food-preserving properties [6, 26]. Several studies also showed that black pepper, clove, cinnamon, and coriander exhibited antioxidant properties [22]. In recent decades, a number of phenolic substances were isolated from a variety of spice sources, including phenolic acids (e.g., gallic acid, caffeic acid, etc.), flavonoids (e.g.,

quercetin, rutin, myricetin, luteolin, naringenin, and silybin), phenolic diterpenes, and volatile oils [27].

5. Conclusion

The most active antioxidant is Indonesian cassia in SOD activity. Clove extract is the highest phenolic content and highest DPPH scavenging activity.

Acknowledgment

We gratefully acknowledge the financial support of the Directorate General of Higher Education, National Ministry of Republic Indonesia for research grant of Hibah Bersaing 2012.

Conflict of Interest

The authors report no conflicts of interest

References

1. WHO Library Cataloguing-in-Publication Data. Global status report on noncommunicable diseases. World Health Organization 2011: ISBN 978 92 4 068645 8.
2. Aruoma OI, Spencer JPE, Rossi R, Aeschbach R, Khan A, Mahmood N, Munoz A, Murcia A, Butler J, Halliwell B. An evaluation of the antioxidant and antiviral action of extracts of rosemary and provencal herbs. *Food Chem Toxicol* 1996;34:449–456.
3. Joshan DS, Rawal S. Comparative evaluation of antioxidant and antihemolytic capacities of plants of Indian origin using multiple antioxidant assays. *Int J Phytopharm* 2012; 2(4):107–115
4. Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacogn Rev*; 2010: 118–126.
5. Kumar M, Kumar A, Dandapat S, Sinha MP. Phytochemical screening and antioxidant potency of *Adhatoda Vasica* and *Vitex Negundo*. *The Bioscan* 2013; 8(2): 727-730.

6. Srinivasan K. Role of spices beyond food flavoring: Nutraceuticals with multiple health effects. *Food Rev Int* 2005 21:167-188.
7. Gao, XM, Xu ZM, Li ZW. *Traditional Chinese Medicines*. People's Health Publishing House: Beijing, China, 2000; p 2019.
8. Hossain M, Brunton N, Barry-Ryan C, Martin-Diana A, Wilkinson M. Antioxidant activity of spice extracts and phenolics in comparison to synthetic antioxidants. *Rasayan J Chem* 2008; 1(4):751-756.
9. Rice-Evans CA, Miller J, Paganga G. Antioxidant properties of phenolic compounds. *Trends Plant Sci* 1997; 2:152-159.
10. Rice-Evans C A, Miller NJ, Bolwell PG, Bramley P M, Pridham JB. The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free Radical Res* 1995; 22: 375-383.
11. Luiz M, Biasutti A, Garcia NA. Micellar effect on the scavenging of singlet molecular oxygen by hydroxybenzenes. *Redox Rep* 2002; 7:23-28.
12. Chen XY, Ahn DU. Antioxidant activities of six natural phenolics against lipid oxidation induced by Fe²⁺ or ultraviolet light. *J Am Oil Chem Soc* 1998; 75: 1717-1721.
13. Gutteridge JM, Halliwell B. *Antioxidants in Nutrition, Health and Disease*. New York: Oxford University Press Inc; 1996
14. Surh YJ. Anti-tumor promoting potential of selected spice ingredients with antioxidative and anti-inflammatory activities: a short review. *Food Chem Toxicol* 2002; 40:1097-1100.
15. Widowati W, Mozef T, Risdian C, Yellianty Y. Anticancer and free radical scavenging potency of *Catharanthus roseus*, *Dendrophthoe petandra*, *Piper betle* and *Curcuma mangga* extracts in breast cancer cell lines. *Oxidant Antioxid Med Sci* 2013; 2(2):137-142
16. Widowati W, Wijaya L, Wargasetia TL, Bachtiar I, Yellianty, Laksmitawati DR. Antioxidant, anticancer, and apoptosis-inducing effects of *Piper* extracts in HeLa cells. *J Exp Integr Med* 2013; 3(3):225-230.
17. Unlu GV, Candan F, Sokmen A, Dafefera D, Polissiou M, Sokmen E, Donmez E, Tepe B. Antimicrobial and antioxidant activity of the essential oil and methanol extracts of *Thymus pectinatus* Fisch. et Mey. Var. *pectinatus* (Lamiaceae). *J Agric Food Chem* 2003; 51:63-67.
18. Ivanova D, Gerova D, Chervenkov T and Yankova T. Polyphenols and antioxidant capacity of Bulgarian medicinal plants. *J. Ethnopharmacol* 2005; 96: 145-150.
19. Widowati W, Herlina T, Ratnawati H, Mozef T. Antioxidant activities and platelet aggregation inhibitor of black tea (*Camellia sinensis* L.) extract and fractions. *Med Plants*; 2011. 3(1):21-26.
20. Sairam RK, Srivastava GC, Agarwal S, Meena RC. Differences in antioxidant activity in response to salinity stress in tolerant and susceptible wheat genotypes. *Biologia Plantarum* 2005; 49(1): 85-91.
21. Shan B, Chai YZ, Sun M, Corke H. Antioxidant Capacity of 26 spice extracts and characterization of their phenolic constituents. *J Agric Food Chem* 2005; 53(20):7749-7759.
22. Gulcin I, Sat I G, Beydemir S, Elmastas M, Kufrevioglu O I. Comparison of antioxidant activity of clove (*Eugenia caryophyllata* Thunb.) buds and lavender (*Lavandula stoechas* L.). *Food Chem* 2004; 87:393-400.
23. Lee K G, Shibamoto T. Antioxidant property of aroma extract isolated from clove buds [*Syzygium aromaticum* (L.) Merr. et Perry]. *Food Chem* 2001;74, 443-448.
24. Zheng W, Wang S Y. Antioxidant activity and phenolic compounds in selected herbs. *J. Agric Food Chem* 2001; 49:5165-5170.
25. Yanishlieva NV, Marinova EM. Stabilisation of edible oils with natural antioxidants. *Eur J. Lipid Sci Technol* 2001;103:752-767.
26. Badei AZM, El-Akel ATM, Faheid SMM, Mahmoud BSM. Application of some spices in flavoring and preservation of cookies: 1-Antioxidant properties of cardamom, cinnamon, and clove. *Dtsch Lebensm.-Rundsch* 2002;98:176-183.
27. Melo ED, Mancini J, Guerra NB. Characterization of antioxidant compounds in aqueous coriander extract (*Coriandrum sativum* L.). *Sci Technol* 2005;38:15-19.

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau PEER REVIEW

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Antioxidant properties of Spice Extracts
 Jumlah Penulis : 4 Orang
 Nama-nama Penulis : **Wahyu Widowati**, Hana Ratnawati, Winsa Husin, Macsaroh Macsaroh
 Status Penulis : Penulis Pertama / ~~Penulis ke-....~~ / Penulis Korespondensi **)
 Identitas Jurnal Ilmiah : a. Nama jurnal : Biomedical Engineering
 b. Nomor ISSN : 2356-2471
 c. Vol., No., Bulan, Tahun : Vol 1; No 1; 2015
 d. Penerbit : Universitas Brawijaya
 e. DOI Artikel (jika ada) : -
 f. Alamat Web Jurnal : <https://be.ub.ac.id/index.php/jibe>
 g. Terindeks di : DOAJ

Kategori Publikasi Jurnal Ilmiah: Jurnal Ilmiah Internasional / ~~Internasional Bereputasi **~~
 (beri tanda \checkmark yang dipilih)

Jurnal Ilmiah Nasional Terakreditasi

Jurnal Ilmiah Nasional / Nasional terindeks ***

HASIL PENILAIAN (Peer Review) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional <input checked="" type="checkbox"/>	Nasional Terakreditasi <input type="checkbox"/>	Nasional *** <input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	2			1,5
b.	Ruang lingkup dan kedalaman pembahasan (30%)	6			5,5
c.	Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	6			5,5
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	6			5,6
	Total	100%	20		18,1

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur.....
 Paper ditulis dengan baik, hasil sesuai judul penelitian, mengikuti kaidah penelitian
- b. Ruang lingkup & kedalaman pembahasan.....
 Penelitian tentang potensi ekstrak rempah (cengkeh, kayumanis, tumbak, kapulaga) sebagai antioksidan secara *In vitro*
- c. Kecukupan & kemutakhiran data serta metodologi.....
 Ide dasar penelitian baik membahas, membandingkan dengan senyawa murni yang terdapat pada berbagai ekstrak rempah
- d. Kelengkapan unsur dan kualitas penerbit.....
 Jurnal Biomed Engineering, terindeks DOAJ. Penerbit Universitas Brawijaya
- e. Indikasi plagiasi.....
 tidak terdapat indikasi plagiarisme atau self-plagiarisme

f. Kesesuaian bidang ilmu Paper bidang biokimia, biomedic sesuai dengan bidang ilmu penulis

REVIEWER 1



(Prof. Dr. Chrismis Novalinda Ginting, M.Kes)

NIK: 0115127801

UNIVERSITAS PRIMA INDONESIA

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau PEER REVIEW

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Antioxidant properties of Spice Extracts
 Jumlah Penulis : 4 Orang
 Nama-nama Penulis : **Wahyu Widowati**, Hana Ratnawati, Winsa Husin, Macsaroh Macsaroh
 Status Penulis : Penulis Pertama / ~~Penulis ke-....~~ / Penulis Korespondensi **)
 Identitas Jurnal Ilmiah :
 a. Nama jurnal : Biomedical Engineering
 b. Nomor ISSN : 2356-2471
 c. Vol., No., Bulan, Tahun : Vol 1; No 1; Agustus, 2015
 d. Penerbit : Universitas Brawijaya
 e. DOI Artikel (jika ada) : -
 f. Alamat Web Jurnal : <https://bc.ub.ac.id/index.php/jibe>
 g. Terindeks di : DOAJ

Kategori Publikasi Jurnal Ilmiah: Jurnal Ilmiah Internasional / ~~Internasional Bereputasi **)~~
 (beri tanda \surd yang dipilih)
 Jurnal Ilmiah Nasional Terakreditasi
 Jurnal Ilmiah Nasional / Nasional terindeks ***)

HASIL PENILAIAN (Peer Review) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / <input checked="" type="checkbox"/>	Nasional Terakreditasi <input type="checkbox"/>	Nasional ***) <input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	2			1,6
b.	Ruang lingkup dan kedalaman pembahasan (30%)	6			5,9
c.	Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	6			3,8
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	6			3,2
	Total	100%	20		18,5

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur
Jurnal ini sudah cukup lengkap dan juga terdapat kesesuaian antara unsur dan isinya.
- b. Ruang lingkup & kedalaman pembahasan
Ruang lingkup bahasan sudah memadai dan ada kedalaman dalam setiap isinya.
- c. Kecukupan & kemutakhiran data serta metodologi
Data-data dan sumber referensi yang digunakan cukup mutakhir serta kelengkapan yang dibuat juga cukup baik.
- d. Kelengkapan unsur dan kualitas penerbit
Kualitas penerbit sudah memiliki reputasi yang baik.

e. Indikasi plagiasi
serasa umum belum terlintas adanya unsur plagiat.

f. Kesesuaian bidang ilmu

Jurnal ini sudah sesuai dengan bidang Ilmu yang diteliti penulis.

REVIEWER 2



(Prof. Dr. Ermi Girsang, M. Kes)

NIK : 0117057501

UNIVERSITAS PRIMA INDONESIA

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau PEER REVIEW

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Antioxidant properties of Spice Extracts
 Jumlah Penulis : 4 Orang
 Nama-nama Penulis : Wahyu Widowati, Hana Ratnawati, Winsa Husin, Maesaroh Maesaroh
 Status Penulis : Penulis Pertama / ~~Penulis ke-....~~ / Penulis Korespondensi **)
 Identitas Jurnal Ilmiah :
 a. Nama jurnal : Biomedical Engineering
 b. Nomor ISSN : 2356-2471
 c. Vol., No., Bulan, Tahun : Vol 1; No 1, 2015
 d. Penerbit : Universitas Brawijaya
 e. DOI Artikel (jika ada) : -
 f. Alamat Web Jurnal : <https://be.ub.ac.id/index.php/jibe>
 g. Terindeks di : DOAJ

Kategori Publikasi Jurnal Ilmiah: Jurnal Ilmiah Internasional / ~~Internasional Bereputasi **)~~
 (beri tanda \checkmark yang dipilih)
 Jurnal Ilmiah Nasional Terakreditasi
 Jurnal Ilmiah Nasional / Nasional terindeks ***)

HASIL PENILAIAN (Peer Review) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi	Nasional Terakreditasi	Nasional ***)	
		<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	2			1,55
b.	Ruang lingkup dan kedalaman pembahasan (30%)	6			9,7
c.	Kecukupan dan kemitakhiran data/informasi dan metodologi (30%)	6			5,65
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	6			5,9
	Total	100%			18,3

Catatan Penilaian ARTIKEL oleh Reviewer :

a. Kelengkapan dan kesesuaian unsur.....
 Paper ditulis dengan baik, hasil sesuai judul penelitian, mengikuti kaidah penelitian.

Jurnal ini sudah cukup lengkap dan juga terdapat kesesuaian antara unsur dan isinya

b. Ruang lingkup & kedalaman pembahasan ... Penelitian tentang potensi ekstrak tempah (engkoh, kayumanis, tumbar, kapulaga) sebagai antioksidan secara in vivo

Ruang lingkup bahasan sudah memadai dan ada kedalaman dalam setiap
tejanya

c. Kecukupan & kemutakhiran data serta metodologi

Data-data dan sumber referensi yang digunakan cukup mutakhir serta kesimpulan cukup sah

d. Kelengkapan unsur dan kualitas penerbit

Ide dasar penelitian baik membahas membandingkan dengan senyawa murni yang
terdapat pada berbagai ekstrak rempah

Kualitas penerbit sudah memiliki reputasi yang baik

e. Indikasi plagiasi tidak terdapat indikasi plagiarisme atau self-plagiarisme

Secara umum belum terlihat adanya unsur plagiasi

f. Kesesuaian bidang ilmu

Paper bidang botanika, farmasi, sesuai dengan bidang Ilmu Penulis

Jurnal ini sudah sesuai dengan bidang ilmu yang diteliti penulis

Medan,
Reviewer 2

(Prof. Dr. Ermi Girsang, M.Kes)
NIK : 0117057501
UNIVERSITAS PRIMA INDONESIA

Medan,
Reviewer 1

(Prof. Dr. Chrismis Novalinda Ginting, M.Kes)
NIK : 0115127801
UNIVERSITAS PRIMA INDONESIA