

- **Manuscript ID** : NPS-15-0062
- **Manuscript Type** : Regular Article
- **Manuscript Category** : None (or N/A)
- **Manuscript Title** : Tea Flavonoids Protect Endothelial Progenitor Cells (EPCs) from Oxidative Damage by Reducing Reactive Oxygen Species (ROS) Level

Dear Dr. Wahyu Widowati:

This automatic message acknowledges the receipt of the above manuscript that you submitted via the *Natural Product Sciences* online submission system. Your manuscript has been successfully checked-in and editors will be assigned for evaluation of your manuscript. Please refer to the assigned manuscript ID number in any correspondence with the NPS Editorial Office or with the editor.

Your paper will be reviewed by two reviewers assigned by editors in the NPS editorial board and final decision made by the editor will be informed by email in due course. Reviewers' suggestions and editor's comments will be then available at the online system. You can monitor the review process for your paper by visiting the "Status of my manuscript" at the NPS online submission system.

If your manuscript is accepted for publication, NPS editorial office will contact you for the production of your manuscript.

Thank you very much for submitting your manuscript to the *Natural Product Sciences*.

Sincerely,

**Korean Society of Pharmacognosy**

Dr. Editor-in-Chief (MD, PhD)

Editor-in-Chief

Tel: +82-51-510-2803

E-mail: [jhjung@pusan.ac.kr](mailto:jhjung@pusan.ac.kr)

Homepage: <http://www.e-nps.or.kr>

- **Manuscript ID** : NPS-15-0062
- **Manuscript Type** : Regular Article
- **Manuscript Title** : Tea Flavonoids Induced Differentiation of Peripheral Blood-derived Mononuclear Cells into Peripheral Blood-derived Endothelial Progenitor Cells and Suppressed Intracellular Reactive Oxygen Species Level of Peripheral Blood-derived Endothelial Progenitor Cells

Dear Dr. Wahyu Widowati:

Your revised manuscript titled `Tea Flavonoids Induced Differentiation of Peripheral Blood-derived Mononuclear Cells into Peripheral Blood-derived Endothelial Progenitor Cells and Suppressed Intracellular Reactive Oxygen Species Level of Peripheral Blood-derived Endothelial Progenitor Cells` has been submitted successfully.

NPS Editorial Office

*Natural Product Sciences*

Tel: +82-51-510-2803

E-mail: [journal@e-nps.or.kr](mailto:journal@e-nps.or.kr)  
Homepage: <http://www.e-nps.or.kr>

- **Manuscript ID** : NPS-15-0062
- **Manuscript Type** : Regular Article
- **Manuscript Title** : Tea Flavonoids Protect Endothelial Progenitor Cells (EPCs) from Oxidative Damage by Reducing Reactive Oxygen Species (ROS) Level

Dear Dr. Wahyu Widowati:

We would like to remind you that the deadline for revised manuscript resubmission is Jan 3, 2016. Please note that revised manuscript must be submitted by this deadline. Please submit the revised manuscript to `Natural Product Sciences` as soon as possible. Thanks for your cooperation.

NPS Editorial Office  
*Natural Product Sciences*  
Tel: +82-51-510-2803  
E-mail: [journal@e-nps.or.kr](mailto:journal@e-nps.or.kr)  
Homepage: <http://www.e-nps.or.kr>

- **Manuscript ID** : NPS-15-0062
- **Manuscript Type** : Regular Article
- **Manuscript Title** : Tea Flavonoids Induced Differentiation of Peripheral Blood-derived Mononuclear Cells into Peripheral Blood-derived Endothelial Progenitor Cells and Suppressed Intracellular Reactive Oxygen Species Level of Peripheral Blood-derived Endothelial Progenitor Cells

Dear Dr. Wahyu Widowati:

Your revised manuscript titled `Tea Flavonoids Induced Differentiation of Peripheral Blood-derived Mononuclear Cells into Peripheral Blood-derived Endothelial Progenitor Cells and Suppressed Intracellular Reactive Oxygen Species Level of Peripheral Blood-derived Endothelial Progenitor Cells` has been submitted successfully.

NPS Editorial Office  
*Natural Product Sciences*  
Tel: +82-51-510-2803  
E-mail: [journal@e-nps.or.kr](mailto:journal@e-nps.or.kr)  
Homepage: <http://www.e-nps.or.kr>

- **Manuscript ID** : NPS-15-0062
- **Manuscript Type** : Regular Article
- **Manuscript Title** : Tea Flavonoids Induced Differentiation of Peripheral Blood-derived Mononuclear Cells into Peripheral Blood-derived Endothelial Progenitor Cells and Suppressed Intracellular Reactive Oxygen Species Level of Peripheral Blood-derived Endothelial Progenitor Cells

Dear Dr. Wahyu Widowati:

We are pleased to inform you that the above-mentioned paper has been accepted for publication at *Natural Product Sciences*. The manuscript has been forwarded to the Manuscript and English editors for further possible revision. You may be requested to reply some more revisions from them.

You will receive a galley proof for your correction from the publisher about a few months in advance of publication. Please return corrections within 5 working days.

If you have not submit assignment of copyright transfer and disclosure of conflict of interest' to editorial office yet, please send it to the office either by fax or e-mail. Without that, we cannot process the editorial work for the publication of your article.

We appreciate you and your colleagues for your contribution to the NPS and anticipate that you will submit your future research articles to the journal.

Sincerely, #EDITOR\_COPYRIGHT#

### **Evaluations (2nd Review) :**

Editorial Comment

**Recommendation** Accept

Ms. Ref. No.: NPS-15-0062

Title:**Tea Flavonoids Protect Endothelial Progenitor Cells (EPCs) from Oxidative Damage by Reducing Reactive Oxygen Species (ROS) Level**

Dear Professor Wahyu Widowati,

**Comments to the Author**

A final disposition of "Accept" has been registered for the above-mentioned manuscript.

I will be contacted in the near future as the associate editor regarding the page proofs for your manuscript. After you approve your page proofs, your manuscript will be published on the next issue of NPS journal. In view of this rapid

publication time, it is important to review your page proofs carefully. Once your page proofs return to NPS it is considered published. Any change to the manuscript will then need to be submitted to the Journal office as an addition or correction.

Thank you again for choosing Natural Product Sciences for the publication of your work.

With kind regards,

**Files**

None (or N/A)

## Tea Flavonoids Induced Differentiation of Peripheral Blood-derived Mononuclear Cells into Peripheral Blood-derived Endothelial Progenitor Cells and Suppressed Intracellular Reactive Oxygen Species Level of Peripheral Blood-derived Endothelial Progenitor Cells

Wahyu Widowati<sup>1,\*</sup>, Laura Wijaya<sup>2</sup>, Dian Ratih Laksmiawati<sup>3</sup>, Rahma Micho Widyanto<sup>4</sup>, Pande Putu Erawijantari<sup>5</sup>, Nurul Fauziah<sup>5</sup>, Indra Bachtiar<sup>2</sup>, and Ferry Sandra<sup>6,7,\*</sup>

<sup>1</sup>Medical Research Center, Faculty of Medicine, Maranatha Christian University, Bandung 40164, Indonesia

<sup>2</sup>Stem Cell and Cancer Institute, Jakarta 13210, Indonesia

<sup>3</sup>Faculty of Pharmacy, Pancasila University, Jakarta 12640, Indonesia

<sup>4</sup>Faculty of Agricultural Technology, Brawijaya University, Malang 65145, Indonesia

<sup>5</sup>Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung 40163, Indonesia

<sup>6</sup>Faculty of Dentistry, Trisakti University, Jakarta 11440, Indonesia

<sup>7</sup>Prodia Clinical Laboratory, Jakarta 10430, Indonesia

**Abstract** – Endothelial dysfunction in atherosclerosis is associated with increasing oxidative stress that could be reversed by antioxidant. Therefore epigallocatechin gallate (EGCG), epicatechin gallate (ECG), epigallocatechin (EGC) and catechin (C) of tea flavonoids were investigated for their roles in regenerating endothelial cell. Peripheral blood mononuclear cells (PB-MNCs) were isolated, plated and cultured in medium with/without treatment of EGCG, ECG, EGC and C. Results showed that among all EGCG, ECG, EGC and C concentrations tested, 12.5  $\mu\text{mol/L}$  was not cytotoxic for peripheral blood-derived endothelial progenitor cells (PB-EPCs). Treatment of EGCG, ECG, EGC or C increased the percentages of CD34, CD133, VEGFR-2 expressions and suppressed hydrogen peroxide-induced percentages of reactive oxygen species (ROS) level in PB-EPCs. Taken together, our current results showed that EGCG, ECG, EGC or C of tea flavonoids could induce differentiation of PB-MNCs into PB-EPCs as well as protect PB-EPCs from oxidative damage by suppressing the intracellular ROS levels.

**Keywords** – Tea flavonoids, Antioxidant, Endothelial progenitor cell, Differentiation, ROS, Apoptosis

### Introduction

Endothelial dysfunctions play a role in the pathogenesis of atherosclerosis and its complications.<sup>1</sup> Increasing oxidative stress and disorder antioxidant mechanism, especially inactivation of nitric oxide (NO) by reactive oxygen species (ROS) such as **superoxide anion radical**, hydrogen peroxide ( $\text{H}_2\text{O}_2$ ), **hydroxyl radical**, and lipid peroxides, may contribute to the endothelial dysfunction, intact vessel wall, up to the rupture of a lipid-rich atherosclerotic plaque.<sup>2</sup> Endothelial dysfunction in atherosclerosis is

associated with increasing oxidative stress that could be reversed by antioxidant therapy.<sup>3</sup> Antioxidants counterbalance the ROS production in cells and modify the regulatory pathways of cell growth.<sup>4</sup>

In several numbers of previous studies, tea consumption as a source of flavonoids was associated with low cardiovascular disease (CVD) and stroke-related mortality.<sup>5</sup> The mortality was significantly decreased along with the increase of tea flavonoids intake in a dose-dependent manner.<sup>6</sup> Some *in vitro* studies showed that flavonoids have strong ROS scavenging activity and prevent low density lipoprotein (LDL) to oxidize.<sup>1</sup> Atherosclerotic animal models study indicated that consuming dietary flavonoids can reduce atherosclerotic plaque development.<sup>8</sup> Flavonoids were reported to have anti-platelet, anti-inflammatory, and antioxidant activities.<sup>9,10</sup> In addition, an epidemiological study reported that consuming flavonoid-containing foods and beverages could reduce CVD risk,


\*Author for correspondence

Wahyu Widowati, Medical Research Center, Faculty of Medicine Maranatha Christian University, Jl. Prof. drg. Surya Sumantri No. 1, Bandung 40164, Indonesia

Tel: +62-81910040010; E-mail: wahyu\_w60@yahoo.com

Ferry Sandra, Faculty of Dentistry, Trisakti University, Jl. Kyai Tapa No.260, Grogol, Jakarta 11440, Indonesia

Tel: +62-81280777803; E-mail: ferrysandra@yahoo.com



**Fig. 1.** Chemical Structures of catechins. Structures of (-)-epigallocatechin gallate (EGCG) (a), (-)-epicatechin gallate (ECG) (b), (-)-epigallocatechin (EGC) (c), and (+)-catechin (C) (d).

as well as increase endothelial function.<sup>7</sup>

The effects of green tea mainly associated with polyphenols content, especially flavonoids with approximately 30% of dry leaf weight.<sup>11</sup> Predominantly flavonoids consist of epigallocatechin gallate (EGCG), epicatechin gallate (ECG), epigallocatechin (EGC) and catechin (C).<sup>12,13</sup> In our previous study, we found that EGCG enhanced proliferation of adipose tissue-derived mesenchymal stem cells (AD-MSCs) and differentiation of AD-MSCs into endothelial progenitor cells (EPCs). Tea flavonoids including EGCG, ECG, EGC and C could also have effects on peripheral blood-derived EPCs (PB-EPCs). Therefore we conducted a study in order to investigate proliferation and differentiation inducing properties of EGCG, ECG, EGC, and C (Fig. 1) on PB-EPCs.

## Experimental

**General Experimental Procedure** – (-)-epigallocatechin gallate (EGCG), (-)-epicatechin gallate (ECG), (-)-epigallocatechin (EGC), and (+)-catechin (C) with purity 95-99% (Fig. 1.) were purchased from Biopurify Phytochemical Ltd. (Chengdu, China). Human fibronectin was purchased from Roche (Basel, Switzerland). VascGrow™ medium was obtained from Stem Cell and Cancer Institute (Jakarta, Indonesia). Ficoll-Paque reagent was purchased from GE Healthcare (Uppsala, Sweden). Fluorescein isothiocyanate (FITC)-conjugated ulex europaeus agglutinin I (UEA-I), and 1,1-dioctadecyl-3,3,3,3-tetramethylindole carbocyanine-labeled acetylated LDL (Dil-acLDL) were purchased from Sigma-Aldrich (St. Louis, MO, USA). The 4',6-diamidino-2-phenylindole (DAPI)

and 2'-7'-dichlorofluorescein diacetate (DCF-DA) were purchased from Invitrogen (Carlsbad, CA, USA). FcR Blocking and phycoerythrin (PE)-conjugated anti-CD133 antibodies were purchased from Miltenyi Biotec (Bergisch Gladbach, Germany). FITC-conjugated anti-CD45/PE-conjugated anti-CD34 antibody was purchased from Becton Dickinson (Franklin Lakes, NJ, USA). PE-conjugated anti-vascular endothelial growth factor (VEGFR)-2/Kinase Domain Receptor (KDR) antibody was purchased from R&D System (Minneapolis, MN, USA). The 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium (MTS) was purchased from Promega (Madison, WI, USA). Fluorescence-activated cell sorting (FACS) was performed using a FACS Calibur Flow Cytometer Becton Dickinson (Franklin Lakes, NJ, USA). Cells were observed using inverted fluorescent microscope (Axiovert 40 CFL, Zeiss).

**Isolation and Culture** – Briefly, total PB-MNCs were isolated from peripheral blood of healthy young human volunteers with Ficoll density gradient centrifugation. PB-MNCs were plated in human fibronectin-coated culture dishes and cultured using VascGrow™ at 37 °C in a humidified, 5% CO<sub>2</sub> incubator. New media was applied after 4 days of culture and the culture was maintained for 7 days. Prior to blood collection, all volunteers signed informed consent. All procedures were approved by the ethics committee of Stem Cell and Cancer Institute, Jakarta and the joint ethics committee of Faculty of Medicine, Maranatha Christian University, Bandung, Indonesia and Immanuel Hospital, Bandung, Indonesia.

**Fluorescent Staining** – Fluorescent staining was performed to detect FITC-UEA-I and Dil-acLDL. To identify the uptake Dil-acLDL, adherent cells were incubated with 1 mg/mL Dil-acLDL at 37 °C for 4 hours. Cells were fixed with 3% paraformaldehyde for 10 minutes. Soon after cleansed, cells were incubated with 1 mg/mL FITC-UEA-I at 37 °C for 1 hour, followed by nucleus staining using DAPI. After staining, cells were observed using an Axiovert 40 CFL inverted fluorescent microscope from Carl Zeiss (Jena, Germany). Cells with double positive for both Dil-acLDL and FITC-UEA-I were defined as PB-EPCs.

**MTS Assay** – This assay was done to quantify viable PB-EPCs under treatment of EGCG, ECG, EGC and C. Briefly, 5 × 10<sup>3</sup> PB-EPCs were seeded in 96-well plates using serum-free medium for 24 hours. After EGCG, ECG, EGC or C treatment, PB-EPCs were added with 20 μL MTS and incubated at 37 °C in a humidified, 5% CO<sub>2</sub> incubator for 4 hours. Optical Density value was measured at 490 nm using a microplate reader from Biorad (Hercules, CA, USA).

**Apoptosis Assay** – PB-EPCs were cultured in 12-well plate and treated with EGCG, ECG, EGC or C for 24 hours. After that, PB-EPCs were collected, washed and stained with propidium iodide solution in phosphate buffer saline (PBS) for 15 minutes. The apoptotic cells were determined as SubG1 area using a fluorescence-activated cell sorting (FACS) Calibur flow cytometer from Becton Dickinson and presented as a percentage of total cells.

**Immunophenotyping** – PB-EPCs were detached using 2 mM ethylenediaminetetraacetate (EDTA) and pre-incubated for 15 minutes at room temperature with FcR Blocking. Then PB-EPCs were incubated at 4 °C with FITC-conjugated anti-CD45/PE-conjugated anti-CD34 and PE-conjugated anti-CD133 antibodies for 15 minutes, followed by PE-conjugated VEGFR-2/KDR antibody for another 40 minutes. Isotype-identical antibodies were also applied and used as controls to negative bindings. Each expression was analyzed using a FACSCalibur.


**Intracellular ROS Assay** – Treated PB-EPCs were incubated in 37 °C with 10 μmol/L DCF-DA for 30 minutes and washed with PBS containing KCl solution. Level of intracellular ROS were measured using a FACSCalibur.

**Statistical Analysis** – Data are presented as mean and standard deviation. Statistical analysis was performed using IBM SPSS Statistics for Windows software version


20.0. (Armonk, NY, USA). One-way analysis of variance test was performed, followed by Duncan's post-hoc tests.

## Result and Discussion

**Lectin binding and Dil-acLDL uptake of the CD34<sup>+</sup>, CD133<sup>+</sup> and VEGFR-2<sup>+</sup> cells** – Cultured cells were adherent with spindle-shaped morphology. Fig. 2B showed lectin binding property, while Fig. 2C showed Dil-acLDL uptake of the cells, marked by fluorescent cells in both Fig. 2B and 2C. These 2 characteristics



**Fig. 2.** PB-EPCs lectin binding and LDL uptake. PB-EPCs were seeded in 96-well plate and subjected to Fluorescent Staining as described in Experimental. A: DAPI, B: FITC-UEA-I, C: Dil-acLDL, D: Merge of A, B and C. White bar: 50 mm.



**Fig. 3.** PB-EPCs immunophenotypes. PB-EPCs were detached and subjected to Immunophenotyping as described in Experimental. A: isotype, B: CD34/45, C: CD133 and its isotype, D: VEGF-R2 and its isotype.

**Table 1.** Effect of EGCG, ECG, EGC and C on PB-EPCs viability

Treatment	Percentage of Viable PB-EPCs (Mean±SD)			
	EGCG	ECG	EGC	C
Untreated	100.00 ± 8.27 <sup>cA</sup>	100.00 ± 8.27 <sup>cA</sup>	100.00 ± 8.27 <sup>cA</sup>	100.00 ± 8.27 <sup>cA</sup>
0.5% DMSO	98.68 ± 5.04 <sup>cA</sup>	98.68 ± 5.04 <sup>cA</sup>	98.68 ± 5.04 <sup>cA</sup>	98.68 ± 5.04 <sup>cA</sup>
12.5 µmol/L	93.20 ± 5.29 <sup>cA</sup>	94.00 ± 2.27 <sup>cA</sup>	93.73 ± 1.04 <sup>cA</sup>	93.39 ± 0.33 <sup>bcA</sup>
25 µmol/L	78.28 ± 4.02 <sup>bA</sup>	68.45 ± 4.96 <sup>bA</sup>	69.23 ± 4.32 <sup>cA</sup>	93.96 ± 4.84 <sup>bcB</sup>
50 µmol/L	71.29 ± 4.18 <sup>bA</sup>	62.18 ± 2.27 <sup>bA</sup>	63.47 ± 3.55 <sup>bA</sup>	86.21 ± 5.81 <sup>bB</sup>
100 µmol/L	30.59 ± 4.96 <sup>aA</sup>	32.43 ± 4.96 <sup>aA</sup>	54.67 ± 3.06 <sup>aB</sup>	61.76 ± 5.94 <sup>aB</sup>

PB-EPCs were treated with 12.5, 25, 50 or 100 µmol/L of EGCG, ECG, EGC or C for 24 hours. MTS Assay was carried out as described in Experimental. Each treatment was done in triplicate. <sup>a,b,c</sup>Means in the same column containing the same superscript are not significant ( $p \geq 0.05$ ), while means in the same column containing different superscript in small letter indicate significant differences ( $p < 0.05$ ). <sup>A,B</sup>Means in the same row containing the same superscript are not significant ( $p \geq 0.05$ ), while means in the same row containing different superscript in capital letter indicate significant differences ( $p < 0.05$ ). Statistical analysis was performed based on Duncan's post-hoc test. SD: standard deviation.

**Table 2.** Effect of EGCG, ECG, EGC and C on apoptosis of PB-EPCs

Treatment	Percentage of SubG1
Untreated	11.68 ± 1.20 <sup>a</sup>
0.5% DMSO	14.93 ± 1.60 <sup>a</sup>
EGCG	14.48 ± 6.83 <sup>a</sup>
ECG	14.57 ± 7.67 <sup>a</sup>
EGC	12.31 ± 3.00 <sup>a</sup>
C	15.97 ± 3.62 <sup>a</sup>

Ten thousands PB-EPCs were seeded in each 12-well plate and treated with 12.5 µmol/L EGCG, ECG, EGC or C for 24 hours. After 24 hours, Apoptosis Assay was carried out as described in Experimental. Each treatment was done in triplicate. The apoptotic cells were determined on the basis of the SubG1 area. The data are presented as mean ± standard deviation. <sup>a</sup>Means in the same column containing the same superscript are not significant ( $p \geq 0.05$ ). Statistical analysis was performed based on Duncan's post-hoc test.

implied the cultured cells as PB-EPCs. The cells were further immunophenotyped for CD34, CD133 and VEGFR-2. Flow cytometric results showed that the cells expressed CD34, CD133 and VEGFR-2 (Fig. 3). Hence, our cultured cells were confirmed as EPCs.

**EGCG, ECG, EGC and C reduced percentages of viable PB-EPCs** – Based on the results of MTS assay (Table 1), PB-EPCs treated with higher concentration of EGCG, ECG, EGC, and C showed significant lower percentages of viable PB-EPCs. Among all concentrations EGCG, ECG, EGC and C tested, 12.5 µmol/L had the lowest potential in reducing viable PB-EPCs percentages. In accordance, under treatment of 12.5 µmol/L EGCG, ECG, EGC or C, percentages of apoptosis resulted were almost similar to the untreated (Table 2). These results were similar to our previous research showing that >25

**Table 3.** Effect of EGCG, ECG, EGC and C on PB-EPCs immunophenotypes

Treatment	CD34 (%)	CD133 (%)	VEGFR-2 (%)
Untreated	52.86 ± 8.44 <sup>b</sup>	0.63 ± 0.02 <sup>ab</sup>	0.22 ± 0.02 <sup>a</sup>
0.5% DMSO	28.96 ± 9.68 <sup>a</sup>	0.36 ± 0.14 <sup>a</sup>	1.78 ± 0.15 <sup>c</sup>
EGCG	51.09 ± 2.77 <sup>b</sup>	0.83 ± 0.41 <sup>ab</sup>	1.22 ± 0.19 <sup>b</sup>
ECG	60.48 ± 2.60 <sup>b</sup>	0.61 ± 0.08 <sup>ab</sup>	2.42 ± 0.13 <sup>c</sup>
EGC	54.70 ± 3.74 <sup>b</sup>	1.13 ± 0.71 <sup>b</sup>	1.21 ± 0.22 <sup>b</sup>
C	62.24 ± 9.22 <sup>b</sup>	1.01 ± 0.13 <sup>ab</sup>	1.47 ± 0.33 <sup>bc</sup>

Isolated-PB-MNCs were cultured with addition of 12.5 µmol/L EGCG, ECG, EGC or C. PB-EPCs Culture was then carried out as described in Experimental. PB-EPCs were then detached for Immunophenotyping as described in Experimental. Each treatment was done in triplicate. The data are presented as mean ± standard deviation. <sup>a,b,c</sup>Means in the same column containing the same superscript are not significant ( $p \geq 0.05$ ), while means in the same column containing different superscript in small letter indicate significant differences ( $p < 0.05$ ). Statistical analysis was performed based on Duncan's post-hoc test.

µg/mL green tea extract induced cytotoxic in PB-EPCs<sup>14</sup>, while 12.5 µmol/L was not cytotoxic for EPCs.

**EGCG, ECG, EGC and C increased expressions of CD34, CD133 and VEGFR-2** – EPCs are defined by their cell surface expressions, including CD34, CD133 and VEGFR-2<sup>15,16</sup>. CD34 and CD133 indicate the plasticity of the cells bearing the stem cell characteristic, whereas VEGFR-2 indicates endothelial characteristic. Based on our current flow cytometric results, EGCG, ECG, EGC or C increased the percentages of CD34, CD133 and VEGFR-2 expressions (Table 3). The highest percentage of CD34 expression was induced by ECG, the highest percentage of CD133 expression was induced by EGC, meanwhile the highest percentage of VEGFR-2 expression was induced by ECG. These results suggested that EGCG,



**Table 4.** Effect of EGCG, ECG, EGC and C on ROS of H<sub>2</sub>O<sub>2</sub>-induced PB-EPCs

Treatment	ROS Level (%)	Ratio of All to Negative Control (%)	Ratio of All to Positive Control (%)
Untreated (negative control)	7.20 ± 1.65 <sup>a</sup>	99.95 ± 22.97 <sup>a</sup>	22.81 ± 5.24 <sup>a</sup>
H <sub>2</sub> O <sub>2</sub> (positive control)	31.55 ± 1.10 <sup>c</sup>	438.15 ± 15.22 <sup>e</sup>	99.99 ± 3.47 <sup>e</sup>
EGCG + H <sub>2</sub> O <sub>2</sub>	12.92 ± 0.70 <sup>c</sup>	179.49 ± 9.67 <sup>c</sup>	40.96 ± 2.21 <sup>c</sup>
ECG + H <sub>2</sub> O <sub>2</sub>	10.83 ± 2.35 <sup>bc</sup>	150.46 ± 32.62 <sup>bc</sup>	34.34 ± 7.44 <sup>bc</sup>
EGC + H <sub>2</sub> O <sub>2</sub>	18.66 ± 2.81 <sup>d</sup>	259.21 ± 39.09 <sup>d</sup>	59.15 ± 8.92 <sup>d</sup>
C + H <sub>2</sub> O <sub>2</sub>	8.85 ± 1.13 <sup>ab</sup>	122.96 ± 15.73 <sup>ab</sup>	28.06 ± 3.59 <sup>ab</sup>

PB-EPCs were treated with 12.5 µmol/L of EGCG, ECG, EGC or C for 30 minutes. PB-EPCs were then treated with H<sub>2</sub>O<sub>2</sub> with final concentration of 200 µmol/L for 1 hour. Treated-PB-EPCs were subjected to Intracellular ROS Assay as described in Experimental. Each treatment was done in triplicate. The data are presented as mean±standard deviation. <sup>a,b,c,d</sup>Means in the same column containing the same superscript are not significant ( $p \geq 0.05$ ), while means in the same column containing different superscript in small letter indicate significant differences ( $p < 0.05$ ). Statistical analysis was performed based on Duncan's post-hoc test.

ECG, EGC and C induced differentiation of peripheral blood mononuclear cells (PB-MNCs) into PB-EPCs. Our previous results also showed that EGCG induced differentiation of AD-MSCs into EPCs<sup>17</sup>. In addition, resulted PB-EPCs showed that VEGFR-2 was the most elevated marker, indicating high endothelial characteristic.

**EGCG, ECG, EGC and C suppressed H<sub>2</sub>O<sub>2</sub>-induced ROS level in PB-EPCs** – DCF-DA has been used for several studies dealing with the effect of ROS in cell culture<sup>18,19</sup>. DCF trapped within the cells was measured to represent the intracellular ROS level<sup>20,21</sup>. Compared with the untreated, induction of H<sub>2</sub>O<sub>2</sub> increased percentages of ROS level in PB-EPCs significantly (Table 4). By pretreatment of EGCG, ECG, EGC or C, H<sub>2</sub>O<sub>2</sub>-induced percentages of ROS level was suppressed. This results was consistent with our previous results. Green tea extract that was supposed to contain flavonoids including EGCG, ECG, EGC and C, suppressed H<sub>2</sub>O<sub>2</sub>-induced ROS level in PB-EPCs<sup>14</sup>. Among all investigated flavonoids, C had the highest ROS suppressing capacity in PB-EPCs, since C was reported to have high 1,1-diphenyl-2-picryl-hydrazyl (DPPH) free radical scavenging and high superoxide dismutase activities<sup>22</sup>.

Taken together, our current results showed that EGCG, ECG, EGC or C of tea flavonoids, could induce differentiation of PB-MNCs into PB-EPCs as well as protect PB-EPCs from oxidative damage by suppressing the intracellular ROS levels. In addition, concentration of 12.5 µmol/L was not cytotoxic for EPCs. This study suggests that tea flavonoids might be related to PB-EPCs production and protection.

### Acknowledgments

This research was funded by the Ministry of Research, Technology and Higher Education, Republic of Indonesia

(No:1.41.02/SEK/IR/PPK./I/2011). Stem Cell and Cancer Institute (SCI), Jakarta for supporting the research methodology and facilities. We thank Maesaroh, I Dewa Gde Sathya Deva and Balqis Balqis at Biomolecular and Biomedical Research Center, Aretha Medika Utama for their technical support.

### References

- (1) Bonetti, P. O.; Lerman, L. O.; Lerman, A. *Arterioscler. Thromb. Vasc. Biol.* **2003**, *23*, 168-175.
- (2) Higashi, Y.; Noma, K.; Yoshizumi, M.; Kihara, Y. *Circ. J.* **2009**, *73*, 411-418.
- (3) Diaz, M. N.; Frei, B.; Vita, J. A.; Keaney J. F. Jr. *N. Engl. J. Med.* **1997**, *337*, 408-416.
- (4) Lin, J.; Cook, N. R.; Albert, C.; Zaharris, E.; Gaziano, J. M.; Van Denburgh, M.; Buring, J. E.; Manson, J. E. *J. Natl. Cancer Inst.* **2009**, *101*, 14-23.
- (5) Yang, C. S.; Landau, J. M. *J. Nutr.* **2000**, *130*, 2409-2412.
- (6) Trevisanato, S. I.; Kim, Y. I. *Nutr. Rev.* **2000**, *58*, 1-10.
- (7) Shenouda, S. M.; Vita, J. A. *J. Am. Coll. Nutr.* **2007**, *26*, 366S-372S.
- (8) Grassi, D.; Mulder, T. P.; Draijer, R.; Desideri, G.; Mollhuizen, H. O.; Ferri, C. *J. Hypertens.* **2009**, *27*, 774-781.
- (9) Hertog, M. G.; Feskens, E. J.; Hollman, P. C.; Katan, M. B.; Kromhout, D. *Lancet* **1993**, *342*, 1007-1011.
- (10) Mukamal, K. J.; Maclure, M.; Muller, J. E.; Sherwood, J. B.; Mittleman, M. A. *Circulation* **2002**, *105*, 2476-2481.
- (11) Balentine, D. A.; Wiseman, S. A.; Bouwens, L. C. *Crit. Rev. Food Sci. Nutr.* **1997**, *37*, 693-704.
- (12) Wolfram, S. *J. Am. Coll. Nutr.* **2007**, *26*, 373S-388S.
- (13) Sano, M.; Tabata, M.; Suzuki, M.; Degawa, M.; Miyase, T.; Maeda-Yamamoto, M. *Analyst* **2001**, *126*, 816-820.
- (14) Widowati, W.; Widyanto, R. M.; Husin, W.; Ratnawati, H.; Laksmiawati, D. R.; Setiawan, B.; Nugrahenny, D.; Bachtiar, I. *Iran J. Basic Med. Sci.* **2014**, *17*, 702-709.
- (15) Wollert, K. C.; Drexler, H. *Circ. Res.* **2005**, *96*, 151-163.
- (16) Sagara, Y.; Vanhnasy, J.; Maher, P. *J. Neurochem.* **2004**, *90*, 1144-1155.
- (17) Widowati, W.; Sardjono, C. T.; Wijaya, L.; Laksmiawati, D. R.; Sandra, F. *J. US-China Med. Sci.* **2012**, *9*, 22-29.
- (18) Murphy, T. H.; Miyamoto, M.; Sastre, A.; Schnaar, R. L.; Coyle, J. T. *Neuron* **1989**, *2*, 1547-1558.
- (19) LeBel, C. P.; Ischiropoulos, H.; Bondy, S. C. *Chem. Res. Toxicol.*

1992, 5, 227-231.

(20) Jie, G.; Lin, Z.; Zhang, L.; Lv, H.; He, P.; Zhao, B. *J. Agric. Food Chem.* **2006**, *54*, 8058-8064.

(21) Valko, M.; Leibfritz, D.; Moncol, J.; Cronin, M. T.; Mazur, M.; Telser, J. *Int. J. Biochem. Cell Biol.* **2007**, *39*, 44-84.

(22) Evacuasiyany, E.; Ratnawati, H.; Liana, L. K.; Widowati, W.; Maesaroh, M.; Mozef, T.; Risdian, C. *Oxid. Antioxid. Med. Sci.* **2014**, *3*, 141-146.

Received April 0, 2016

Revised May 0, 2016

Accepted June 0, 2016