

Complimentary and personal copy for

Nguyen Phuong Thao, Bui Thi Thuy Luyen, Wahyu Widowati,
Nurul Fauziah, Maesaro Maesaroh, Tati Herlina, Zahid
Manzoor, Irshad Ali, Young Sang Koh, Young Ho Kim

www.thieme.com

**Anti-inflammatory Flavonoid
C-Glycosides from *Piper aduncum*
Leaves**

DOI 10.1055/s-0042-108737
Planta Med 2016; 82: 1475–1481

This electronic reprint is provided for non-commercial and personal use only: this reprint may be forwarded to individual colleagues or may be used on the author's homepage. This reprint is not provided for distribution in repositories, including social and scientific networks and platforms.

Publishing House and Copyright:

© 2016 by
Georg Thieme Verlag KG
Rüdigerstraße 14
70469 Stuttgart
ISSN 0032-0943

Any further use
only by permission
of the Publishing House

 **Thieme**

Editorial Board

Planta Medica

An International Journal of Natural Products and Medicinal Plant Research

Editors-in-Chief

Oliver Kayser, Dortmund, Germany
Robert Fürst, Frankfurt, Germany

Editors

Alessandra Braca, Pisa, Italy
Olivier Potterat, Basel, Switzerland
Jose-Luis Rios, Valencia, Spain
Judith Rollinger, Vienna, Austria

Associate Editor for Planta Medica International Open

Luc Pieters, Antwerp, Belgium

Editorial Office

René Prétôt, Muttenz, Switzerland

Honorary Advisory Board

Yoshinori Asakawa, Tokushima, Japan
Kurt Hostettmann, Geneva, Switzerland
Arnold Vlietinck, Antwerp, Belgium

Advisory Board

John Arnason, Ottawa, Canada
Anna-Rita Bilia, Florence, Italy
Mark Butler, S. Lucia, Australia
Veronika Butterweck, Romanshorn, Switzerland
Angela Calderon, Auburn, AL, USA
Claus Cornett, Copenhagen, Denmark
Hartmut Derendorf, Gainesville, USA
Nunziatina De Tommasi, Salerno, Italy
Alberto Dias, Braga, Portugal
Alfonso Garcia-Pineros, Frederick, MD, USA
Jürgen Gertsch, Bern, Switzerland
De-an Guo, Shanghai, China
Matthias Hamburger, Basel, Switzerland
Andreas Hensel, Münster, Germany
Judit Hohmann, Szeged, Hungary
Ikhlas Khan, Oxford MS, USA
Jinwoong Kim, Seoul, Korea
Anna Kiss, Warsaw, Poland
Wolfgang Kreis, Erlangen, Germany
Noberto Pepporine Lopez, Ribeirão Preto, Brasil
Matthias Melzig, Berlin, Germany
Eduardo Munoz, Cordoba, Spain
Nicholas H. Oberlies, Greensboro NC, USA
Peter Proksch, Düsseldorf, Germany
Ping-Jyun Sung, Pingtung, Taiwan
Deniz Tasdemir, Kiel, Germany
Michał Tomczyk, Białystok, Poland
Heikki Vuorela, Helsinki, Finland
Jean-Luc Wolfender, Geneva, Switzerland
Yang Ye, Shanghai, China

Georg Thieme Verlag KG

Stuttgart · New York

Rüdigerstraße 14
D-70469 Stuttgart
Postfach 30 11 20
D-70451 Stuttgart

Thieme Publishers

333 Seventh Avenue
New York, NY 10001, USA
www.thieme.com



Year
2016

Issue

- 18: Special Issue Saponin
- 17: 1447-1492
- 16: 1387-1445
- 15: 1309-1386
- 14: 1225-1307
- 13: 1143-1223 | E6
- 11/12: Special Issue Prof.
- 09/10: Special Issue: Mar

> Current Issue
> Sample Issue (01/2019)

DOI: 10.1055/s-00000058

Planta Medica



Issue 17 · Volume 82 · November 2016

DOI: 10.1055/s-006-32941

eFirst Issue Most Read

Reviews

- 1447 Herman, Anna; Herman, Andrzej P.:
Topically Used Herbal Products for the Treatment of Psoriasis – Mechanism of Action, Drug Delivery, Clinical Studies

[Full Text](#) [HTML](#) [PDF \(146 kb\)](#)

Biological and Pharmacological Activity

Original Papers

- 1456 Vattanarongkup, Jaturavit; Piyachaturawat, Pawinee; Tuchinda, Patoomratana; Sanvarinda, Pimtip; Sanvarinda, Yupin; Jantaratnotai, Nattinee:

Protective Effects of a Diarylheptanoid from *Curcuma comosa* Against Hydrogen Peroxide-Induced Astroglial Cell Death

[Full Text](#) [HTML](#) [PDF \(310 kb\)](#)

- 1463 Marques, Lucia C. F.; Pinheiro, Aruanã J. M. C. R.; Araújo, João G. G.; de Oliveira, Raimundo A. G.; Silva, Selma N.; Abreu, Iracelle C.; de Sousa, Eduardo M.; Fernandes, Elizabeth S.; Luchessi, André D.; Silbiger, Vivian N.; Nicolete, Roberto; Lima-Neto, Lidio G.:

Anti-Inflammatory Effects of a Pomegranate Leaf Extract in LPS-Induced Peritonitis

[Full Text](#) [HTML](#) [PDF \(139 kb\)](#)

- 1468 Dlodla, Phiyayinkosi Vusi; Muller, Christo John Frederick; Joubert, Elizabeth; Louw, Johan; Gabuza, Kwazi Bethuel; Huisamen, Barbara; Essop, M. Faadiel; Johnson, Rabia:

Phenylpyruvic Acid-2-O-β-D-Glucoside Attenuates High Glucose-Induced Apoptosis in H9c2 Cardiomyocytes

[Full Text](#) [HTML](#) [PDF \(275 kb\)](#)

> Supporting Information

- 1475 Thao, Nguyen Phuong; Luyen, Bui Thi Thuy; Widowati, Wahyu; Fauziah, Nurui; Maesaroh, Maesaroh; Herlina, Tati; Manzoor, Zahid; Ali, Irshad; Koh, Young Sang; Kim, Young Ho:

Anti-inflammatory Flavonoid C-Glycosides from *Piper aduncum* Leaves

[Full Text](#) [HTML](#) [PDF \(199 kb\)](#)

> Supporting Information

Natural Product Chemistry and Analytical Studies

Original Papers

- 1482 Masoumi-Ardakani, Yaser; Mandegary, Ali; Esmailpour, Khadijeh; Najafipour, Hamid; Shariffar, Fariba; Pakravanan, Mahboobeh; Ghazvini, Hamed:

Chemical Composition, Anticonvulsant Activity, and Toxicity of Essential Oil and Methanolic Extract of *Elettaria cardamomum*

[Full Text](#) [HTML](#) [PDF \(160 kb\)](#)

- 1487 Mudge, Elizabeth M.; Liu, Ying; Lund, Jensen A.; Brown, Paula N.:

Single-Laboratory Validation for the Determination of Flavonoids in Hawthorn Leaves and Finished Products by LC-UV

[Full Text](#) [HTML](#) [PDF \(122 kb\)](#)

Top of Page

Anti-inflammatory Flavonoid C-Glycosides from *Piper aduncum* Leaves

Authors

Nguyen Phuong Thao^{1,2*}, Bui Thi Thuy Luyen^{1*}, Wahyu Widowati³, Nurul Fauziah⁴, Maesaro Maesaroh⁴, Tati Herlina⁵, Zahid Manzoor⁶, Irshad Ali⁶, Young Sang Koh⁶, Young Ho Kim¹

Affiliations

The affiliations are listed at the end of the article

Key words

- *Piper aduncum*
- Piperaceae
- Flavonoid C-glycoside
- IL-12 p40
- IL-6
- TNF- α
- LPS

Abstract

Four new compounds, acacetin 8-C-[β -D-apiofuranosyl-(1 \rightarrow 2)- β -D-glucopyranoside] (**1**), 7-methoxyacacetin 8-C-[β -D-apiofuranosyl-(1 \rightarrow 3)- β -D-glucopyranoside] (**2**), 7-methoxyacacetin 8-C-[β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside] (**3**), and 4''-O-acetylacacetin 8-C-[α -L-rhamnopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside] (**4**), along with ten known compounds (**5–14**), were isolated from *Piper aduncum* leaves. The effects of these compounds on lipopolysaccharide-induced expression of the proinflammatory cytokines IL-12 p40, IL-6, and TNF- α in bone marrow-

derived dendritic cells were evaluated. Compounds **2**, **3**, **6**, **8**, **9**, and **11–13** inhibited the production of both IL-12 p40 and IL-6, with IC₅₀ values ranging from 0.35 \pm 0.01 to 1.40 \pm 0.04 μ M and 1.22 \pm 0.02 to 3.79 \pm 0.10 μ M, respectively. Compounds **5** and **10** only showed strong inhibition effects on the production of IL-12 p40, with IC₅₀ values of 2.76 \pm 0.08 and 0.39 \pm 0.05 μ M, respectively. However, all compounds showed weak activity or no activity on TNF- α production at the tested concentrations.

Supporting information available online at <http://www.thieme-connect.de/products>

Introduction

The usual outcome of acute inflammation is successful resolution and repair of tissue damage. However, it has also been recognized as the central cause of many prevalent diseases, including arthritis, periodontal disease, cardiovascular diseases, cancer, and Alzheimer's disease [1]. Cytokines are primarily involved in host responses to infection, immune responses, inflammation, and trauma. Several cytokines, called proinflammatory cytokines, tend to increase disease symptoms and produce fever, inflammation, tissue destruction, shock, and even death when they are administered to humans. Therefore, compounds that could block or inhibit the production of proinflammatory cytokines might be beneficial anti-inflammatory agents [2].

The genus *Piper* (Piperaceae) is well known for producing a large number of physiologically active compounds and is widely used in folklore medicine in the West Indies and Latin America. The Piperaceae family, in which the genus *Piper*

belongs, comprises approximately 2000 species distributed in the tropical regions of the world [3,4]. *Piper aduncum* L. is widely used in folk medicine to treat stomachaches (Jamaica) and as an anti-inflammatory, diarrhea (Peru), and antiseptic to heal wounds (Brazil and Papua New Guinea) [3–5]. Previous phytochemical studies of *P. aduncum* reported the isolation of dihydrochalcones, chalcones, flavanones, chromene, phenylpropanoids, and benzoic acid derivatives [6–8]. Several pharmacological effects of *P. aduncum* extracts have been demonstrated, including antileishmanial, antibacterial, cytotoxic, and antifungal activities [6,9]. Furthermore, some of the isolated compounds have been shown to be active against promastigote and intracellular amastigotes, causing damaging effects on DNA, and antileishmanial, antimicrobial, molluscicidal, antitumor, and antifungal activities [8,10,11]. However, anti-inflammatory studies on *P. aduncum* are limited. To date, only an essential oil (dillapiol) of this plant was investigated for anti-inflammatory activity in the carrageenan-induced rat paw edema model [12].

As a part of our ongoing investigations on the chemical constituents of medicinal plants with

received February 25, 2016
revised May 1, 2016
accepted May 5, 2016

Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-108737>
Published online June 10, 2016
Planta Med 2016; 82: 1475–1481 © Georg Thieme Verlag KG Stuttgart · New York · ISSN 0032-0943

Correspondence

Prof. Dr. Young Ho Kim
College of Pharmacy
Chungnam National University
99 Yuseong-gu
Daejeon 305-764
Korea
Phone: + 82 4 2821 5933
Fax: + 82 4 28 23 65 66
yhk@cnu.ac.kr

* These authors contributed equally to this work.




Fig. 1 Chemical structures of compounds 1–14 isolated from *P. aduncum* leaves. (Color figure available online only.)

anti-inflammatory activities, we found that the methanolic extract of *P. aduncum* showed significant anti-inflammatory effects *in vitro* with an inhibition value of 65.9% (20.0 µg/mL). In the present study, we report the isolation and structural elucidation of four new flavonoids (1–4) and ten known compounds (5–14) from *P. aduncum* leaves (Fig. 1). Also, the anti-inflammatory effects of these compounds on lipopolysaccharide (LPS)-induced expression of the proinflammatory cytokines IL-12 p40, IL-6, and TNF-α in bone marrow-derived dendritic cells (BMDCs) were evaluated.

Results and Discussion

A MeOH extract of the leaves of *P. aduncum* was suspended in H₂O and successively partitioned with CH₂Cl₂ and EtOAc. The EtOAc fraction was separated by various chromatographic experiments to afford 14 C-glycosyl flavones. Ten known compounds including isospinosin (5) [13], 2-O-β-D-glucosyl-8-C-β-D-glucosyl-4'-O-methylapigenin (6) [14], apigenin-8-C-neohesperidoside (7) [15], 2''-O-α-rhamnosyl-4'-O-methyl vitexin (8) [16], ficuflavoside (8-C-(2''-O-β-D-apiofuranosyl)-β-D-glucopyranosyl apigenin) (9) [17], apigenin 8-C-β-glucopyranoside (10) [18], cytisoside (11) [19], isoembigenin (12) [20], orientin (13) [21], and 4'-O-methylorientin (14) [22] were identified by NMR analysis as well as comparison with previous reports (Fig. 1). The C-glycosyl flavones were isolated for the first time from the leaves of this plant, indicating that these compounds could be considered chemotaxonomic markers for the identification of *P. aduncum*.

Compound 1 was obtained as a yellow amorphous powder. The HR-ESI-MS of 1 was consistent with the molecular formula of C₂₇H₃₀O₁₄. The ¹³C NMR and DEPT spectra (Table 1) revealed 27 carbon signals, including 15 signals that were assigned to a flavone aglycon, as well as two anomeric and nine other oxygenated carbon signals, of which three methylenes were ascribable to C-4''' of an apiofuranosyl unit, and C-6''' of a glucopyranosyl unit, respectively. Assignment of the protons and carbons in the aglycone moiety was done based on 2D spectra. All of the sugar proton resonances were assigned by a COSY experiment, and the corresponding ¹³C resonances were then identified by the HMQC.

The COSY spectra of compound 1 indicated two individual correlation systems of sugar units [H-1'' (δ_H 4.99)/H-2'' (δ_H 4.24)/H-3'' (δ_H 3.65)/H-4'' (δ_H 3.70)/H-5'' (δ_H 3.46)/H-6'' (δ_H 3.84 and 4.01)] and [H-1''' (δ_H 5.16)/H-2''' (δ_H 3.78)]. The connectivity of these fragments was determined through HMBC. The downfield shift was observed for C-2'' (δ_C 77.4), as well as HMBC correlations of H-1''' (δ_H 5.16) with C-2'' (δ_C 77.4), and H-2'' (δ_H 4.24) with C-1''' (δ_C 111.1), C-1'' (δ_C 73.7), and C-8 (δ_C 105.4), demonstrating the (1 → 2) linkage between the apiosyl and glucosyl units in 1. Furthermore, acid hydrolysis of 1, followed by TLC, GC analysis, and a comparison with authentic D-apiose, confirmed the presence of a D-apiose moiety in that compound (see Materials and Methods). A C-8 substituted acetin structure was suggested by the ¹H (δ_H 6.25, s, H-6) and ¹³C NMR (δ_C 105.4 for C-8 and 99.3 for C-6) signals characteristic of C-8-glycosylated flavones [23]. In addition, the HMBC correlations of H-1'' (δ_H 4.99) with C-7 (δ_C 164.1), C-8 (δ_C 105.4), and C-9 (δ_C 158.4) confirmed that the disaccharide chain was bonded by a C-glycosidic linkage at the 8-position. Accordingly, compound 1 was identified as acetin 8-C-[β-D-apiofuranosyl-(1 → 2)-β-D-glucopyranoside].

Compound 2 was obtained as a yellow amorphous powder. The HR-ESI-MS was consistent with the molecular formula of C₂₈H₃₂O₁₄. The ¹H and ¹³C NMR spectra (Table 1) of 2 were similar to those of 1. The differences between the spectra of compounds 1 and 2 were with regard to the presence of two methoxy groups and the linkage between two sugar units, glucose and apiose. The downfield shift of C-3'' (δ_C 84.8) and the upfield shift of C-2'' (δ_C 70.9) of the glucopyranosyl unit, in comparison with those observed in 1, were indicative of a glycosidation at C-3''. In addition, the interglycosidic linkage of the apiofuranosyl moiety in 2 was established unambiguously by the HMBC spectrum to be at the C-3'' position of the glucopyranosyl unit based on the cross-peak between H-1''' (δ_H 5.20) with C-3'' (δ_C 84.8) and H-3'' (δ_H 3.45) with C-1''' (δ_C 109.8). The position of the disaccharide chain was determined as C-8 based on the HMBC correlations of the anomeric proton H-1'' (δ_H 4.76) of glucosyl with C-7 (δ_C 164.0), C-8 (δ_C 105.2), and C-9 (δ_C 155.8) of an aglycon. Two methoxy groups were located at C-7 (δ_C 164.0) and C-4' (δ_C 163.0), as indicated by the HMBC correlations of a methoxy proton at δ_C 3.89 with C-7 (δ_C 164.0), and at δ_H 3.87 with C-4' (δ_C 163.0), respectively. In addition, acid hydrolysis of 2, followed by TLC, GC analysis, and a comparison with authentic D-apiose, confirmed the presence of a D-apiose moiety in 2 (see Materials and Methods). Therefore, the structure of 2 was identified as 7-methoxyacetin 8-C-[β-D-apiofuranosyl-(1 → 3)-β-D-glucopyranoside]. Compound 3 was obtained as a yellow powder. The HR-ESI-MS spectra of 3 indicated the molecular formula of C₂₉H₃₄O₁₅. The ¹H and ¹³C NMR spectra of 3 showed the similar aglycon of 2 as 7-methoxyacetin (Table 1). The presence of a glucopyranosyl unit, instead of a terminal apiofuranosyl unit, was the main difference in the structure of 3. Two individual correlations for the spin systems of H-1''/H-2''/H-3''/H-4''/H-5''/H-6'' and H-1'''/H-2'''/H-3'''/H-4'''/H-5'''/H-6''' were observed in the COSY spectra of 3, which supports the backbone proton assignments of the two sugar units. HMBC correlations of the anomeric proton H-1''' (δ_H 3.87)/C-2'' (δ_C 81.3) and proton H-2'' (δ_H 4.05)/C-1''' (δ_C 105.3), as well as the downfield shift of C-2'' (δ_C 81.3), suggested the structure of a β-D-glucopyranosyl (1 → 2) β-D-glucopyranosyl disaccharide. The position of the sugar moiety in 3 through a C-C linkage was identified as C-8 based on the HMBC correlations of the anomeric proton H-1'' (δ_H 4.83) with C-8 (δ_C 105.3), C-7 (δ_C 164.0), and C-9 (δ_C 155.9). Acid hydrolysis experiments con-

Table 1 ^1H and ^{13}C NMR spectroscopic data for compounds **1–3**.

Pos.	1		2		3	
	$\delta_{\text{H}}^{\text{a,c}}$ (mult., J in Hz)	$\delta_{\text{C}}^{\text{a,d}}$	$\delta_{\text{H}}^{\text{b,c}}$ (mult., J in Hz)	$\delta_{\text{C}}^{\text{b,d}}$	$\delta_{\text{H}}^{\text{b,c}}$ (mult., J in Hz)	$\delta_{\text{C}}^{\text{b,d}}$
2	–	166.1	–	164.5	–	164.3
3	6.55 (s)	104.0	6.92 (s)	103.8	6.88 (s)	103.9
4	–	184.0	–	182.9	–	182.8
5	–	162.5	–	162.0	–	162.0
6	6.25 (s)	99.3	6.55 (s)	95.6	6.50 (s)	95.7
7	–	164.1	–	164.0	–	164.0
8	–	105.4	–	105.2	–	105.3
9	–	158.4	–	155.8	–	155.9
10	–	105.7	–	105.6	–	104.8
1'	–	124.7	–	123.7	–	123.8
2'	8.01 (d, 8.4)	129.8	8.16 (d, 9.0)	129.5	8.15 (d, 8.4)	129.5
3'	7.01 (d, 9.0)	115.5	7.09 (d, 9.0)	115.0	7.09 (d, 9.0)	115.0
4'	–	164.3	–	163.0	–	162.9
5'	7.01 (d, 9.0)	115.5	7.09 (d, 9.0)	115.0	7.09 (d, 9.0)	115.0
6'	8.01 (d, 8.4)	129.8	8.16 (d, 9.0)	129.5	8.15 (d, 8.4)	129.5
7-OCH ₃	–	–	3.89 (s)	57.1	3.84 (s)	57.1
4'-OCH ₃	3.85 (s)	56.0	3.87 (s)	56.1	3.83 (s)	56.1
1''	4.99 (d, 10.2)	73.7	4.76 (d, 10.2)	73.8	4.83 (d, 9.6)	71.9
2''	4.24 (t, 9.0)	77.4	3.99*	70.9	4.05 (t, 9.0)	81.3
3''	3.65 (t, 9.0)	80.9	3.45 (t, 8.4)	84.8	3.48*	78.8
4''	3.70 (t, 9.0)	72.2	3.49 (m)	69.3	3.45*	70.6
5''	3.46 (m)	82.8	3.29 (m)	82.3	3.23 (m)	82.4
6''	3.84*	63.0	3.59*	61.4	3.54 (m)	61.4
	4.01 (dd, 1.2, 12.0)		3.77*		3.73 (m)	
1'''	5.16 (br s)	111.1	5.20 (d, 1.8)	109.8	3.87*	105.3
2'''	3.78 (br s)	77.8	3.71*	76.4	2.71 (m)	75.0
3'''	–	80.6	–	79.5	2.40 (m)	76.6
4'''	2.55 (d, 9.6)	74.7	3.55*	74.0	2.87*	69.8
	3.15 (d, 9.6)		4.01*			
5'''	3.28 (d, 12.0)	65.9	3.36 (m)	64.0	2.87*	76.7
	3.40 (d, 12.0)					
6'''	–	–	–	–	3.06 (m)	60.7
					3.12 (m)	

^a CD₃OD; ^b DMSO-*d*₆; ^c 600 MHz; ^d 150 MHz. *Overlapped signals. Assignments were done by DEPT, HMQC, HMBC, and COSY

confirmed the presence of D-glucose in **3**. This conclusion was further supported by TLC, GC analysis, and comparisons with an authentic D-glucose sample. Thus, compound **3** was identified as 7-methoxyacacetin 8-C-[β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside].

Compound **4** was obtained as yellow powder. The HR-ESI-MS spectra of **4** indicated the molecular formula of C₃₀H₃₄O₁₅. The ^1H NMR spectra of **4** showed signals of six aromatic protons, including two singlet signals, four aromatic protons, one methoxy group, two methyl groups, and two anomeric protons, together with carbinol protons, suggesting the presence of two sugar units. The ^{13}C NMR spectra displayed 30 carbon resonance signals, including 15 carbons belonging to an aglycon moiety, 12 carbons of sugar units, 1 methoxy carbon, and 2 carbons of an acetyl group.

The ^1H and ^{13}C NMR data showed doubling of the signals due to restricted rotation (Table 2). The rotameric signals show almost the same ^1H NMR signal intensity of 3:2. Hence, the rotameric partners of **4** could also be observed in the ^{13}C NMR spectrum. Because of the similar ^1H and ^{13}C resonance signals between **4** and **1**, the aglycon of **4** was identified as acacetin. The presence of glucose and rhamnose units was observed in the ^1H , ^{13}C , and COSY spectra of **4**. Furthermore, acid hydrolysis of **4**, followed by TLC, GC analysis, and a comparison with authentic L-rhamnose,

confirmed the presence of an L-rhamnose moiety in **4** (see Materials and Methods). The HMBC correlations of the anomeric proton of rhamnose H-1''' (δ_{H} 5.42/5.30) and C-2'' (δ_{C} 76.2/76.1) of glucose, proton H-2'' (δ_{H} 4.20) of glucose and C-1''' (δ_{C} 101.3) of rhamnose, and C-8 (δ_{C} 105.9/105.7) of the A ring suggested the structure of an α -L-rhamnopyranosyl (1 \rightarrow 2)- β -D-glucopyranosyl disaccharide. The position of an acetyl group at C-4''' was confirmed by the cross-peak observed between H-4''' (δ_{H} 4.53/4.61) and C-1''' (δ_{C} 172.5), as well as the downfield shift of C-4''' (δ_{C} 75.2). The C-8 substituted structure was determined by the HMBC correlation of H-1'' (δ_{H} 5.05) of the glucopyranosyl unit with C-8 (105.9/105.7), C-7 (δ_{C} 164.4/164.3), and C-9 (δ_{C} 157.7/156.2) of the aglycon. Therefore, compound **4** was assigned as 4'''-O-acetylacacetin 8-C-[α -L-rhamnopyranosyl-(1 \rightarrow 2)- β -D-glucopyranoside].

The isolated compounds were assessed for anti-inflammatory activity on LPS-induced expression of the proinflammatory cytokines IL-12 p40, IL-6, and TNF- α by BMDCs (see Materials and Methods). Many natural products have been shown to exhibit anti-inflammatory activities by inhibiting the production of proinflammatory cytokines, including IL-12 p40, IL-6, and TNF- α , which play a crucial role in host defenses and inflammation [24]. Thus, agents that block the excessive production of these cytokines might be candidates for use in the treatment of inflam-

Table 2 ¹H and ¹³C NMR spectroscopic data for compound **4** in CD₃OD.

Pos.	Aglycon		No.	Sugar moiety	
	$\delta_{\text{H}}^{\text{a}}$ (mult., J in Hz)	$\delta_{\text{C}}^{\text{b}}$		$\delta_{\text{H}}^{\text{a}}$ (mult., J in Hz)	$\delta_{\text{C}}^{\text{b}}$
2	–	166.0 165.3	1''	5.05 (d, 9.6)	73.7 75.1
3	6.60 (s)	104.0	2''	4.20 (dd, 7.8, 9.6)	76.2 76.1
4	–	184.0	3''	3.67 (m)	81.7 81.4
5	–	162.9 162.7	4''	3.68*	72.3 72.0
6	6.28 (s) 6.23 (s)	99.8 101.1	5''	3.49 (m)	83.0 82.9
7	–	164.4 164.3	6''	4.01* 3.84*	63.0 62.5
8	–	105.9 105.7	1'''	5.42 (br s) 5.30 (br s)	101.3
9	–	157.7 156.2	2'''	3.57 (t, 9.6) 3.67*	71.7 71.5
10	–	105.5	3'''	3.48 (dd, 3.6, 9.6) 2.97 (dd, 3.6, 9.6)	69.9
1'	–	124.4	4'''	4.53 (t, 9.6) 4.61 (t, 9.6)	75.2
2'	8.03 (d, 9.0) 7.85 (d, 8.4)	129.9 129.3	5'''	2.31 (m) 2.05 (m)	67.2 66.7
3'	7.03 (d, 9.0) 6.98 (d, 8.4)	115.6 115.5	6'''	0.67 (d, 6.0) 0.50 (d, 6.0)	17.9 17.8
4'	–	164.2 164.1	1''''	–	172.5
5'	7.03 (d, 9.0) 6.98 (d, 8.4)	115.6 115.5	2''''	2.02 (s) 1.91 (s)	20.9 20.9
6'	8.03 (d, 9.0) 7.85 (d, 8.4)	129.9 129.3			
4'-OCH ₃	3.86 (s) 3.84 (s)	56.0			

^a 600 MHz; ^b 150 MHz. *Overlapped signals. Assignments were done by DEPT, HMQC, HMBC, and COSY

matory diseases. BMDs were pretreated with the isolated compounds for 18 h, and then viability was measured using the MTT assay. At a concentration of 5.0 μM , compound **4** showed toxicity. Other compounds did not exhibit significant cytotoxic effects (data not shown). Thus, the effects of compounds **1–3** and **5–14** on LPS-stimulated production of proinflammatory cytokines, including IL-12 p40, IL-6, and TNF- α , in BMDs were evaluated at concentrations of 0.2, 0.5, 1.0, 2.0, and 5.0 μM . SB203580, a bicyclic imidazole compound and an inhibitor of p38 mitogen-activated protein (MAP) kinase [25], was used as a positive control, with IC₅₀ values of 5.00 \pm 0.16, 3.50 \pm 0.12, and 7.20 \pm 0.13 μM for IL-12 p40, IL-6, and TNF- α , respectively. Compounds **2, 3, 5, 6, and 8–13** potently inhibited IL-12 p40 production in LPS-stimulated BMDs with IC₅₀ values ranging from 0.35 \pm 0.01 to

2.76 \pm 0.08 μM (Table 3 and Fig. 2A). Among them, compound **9** showed the greatest inhibitory activity, up to 83.74% at a concentration of 5.0 μM and 42.66% after dilution to 0.2 μM . Other compounds were weak or inactive (Fig. 2A).

To investigate the effects of the tested compounds on other pro-inflammatory cytokines, their effects on IL-6 production were measured. Compounds **2, 3, 6, 8, 9, and 11–13** showed potent inhibition of IL-6 production, with IC₅₀ values ranging from 1.22 \pm 0.02 to 3.79 \pm 0.10 μM (Table 3). Compound **3** showed the greatest inhibition of IL-6 production, with 61.63% at a concentration of 5.0 μM and 37.47% when diluted to 0.2 μM (Fig. 2B). Compounds **5** and **10** showed moderate activity, with inhibition percentages ranging from 4.11% to 39.11% at the tested concentrations. Other compounds were inactive. In addition, the isolated compounds were also evaluated for TNF- α production. However, all of them showed weak activity or were inactive at the tested concentrations from 0.2 to 5.0 μM (Fig. 2C). As shown in Table 3, compounds **2, 3, 6, 8, 9, and 11–13** inhibited the production of both IL-12 p40 and IL-6. Compounds **5** and **10** showed potent inhibitory effects only on the production of IL-12 p40. Compounds **1, 7, and 14** were inactive in terms of production of both IL-12 p40 and IL-6.

Flavonoids, one of the most widespread groups of natural products, possess a wide variety of pharmacological and biochemical properties, including antioxidant, antimicrobial, antiallergenic, and anti-inflammatory effects. Among them, flavonoid C-glycosides such as vitexin, isoorientin, orientin, and isovitexin were shown to have significant anti-inflammatory activity [26,27]. Therefore, our results were in agreement with those from previous studies. This study has provided evidence of the anti-inflammatory properties of flavonoid C-glycosides that could be promising in the development of anti-inflammatory agents.

Materials and Methods

General experimental procedures

Optical rotation was recorded on a JASCO DIP-370 automatic digital polarimeter. The NMR spectra were measured using a JEOL ECA 600 spectrometer with TMS as the internal standard. The ESI mass spectra were performed on an AGILENT 1100 LC-MSD trap spectrometer. The HR-ESI-MS were obtained from an Agilent 6530 Accurate-Mass Q-TOF LC/MS system. GC used an instrument (SHIMADZU GC-14B). Silica gel (70–230, 230–400 mesh, Merck) and YMC RP-18 resins (75 μm , Fuji Silysia Chemical Ltd.) were used as absorbents in the CC. TLC plates (silica gel 60 F₂₅₄ and RP-18 F₂₅₄, 0.25 μm , Merck) were purchased from Merck KGaA. Spots were detected under UV radiation (254 and 365 nm) and by spraying the plates with 10% H₂SO₄, followed by heating with a heat gun. Other chemical reagents and standard compounds were purchased from Sigma-Aldrich.

Plant material

The leaves of *P. aduncum* were collected from Coblong-Bandung, West Java, Indonesia in September 2014 and taxonomically identified by the staff at the Herbarium Laboratory, Department of Biology, School of Life Sciences and Technology, Bandung Institute of Technology, Bandung, West Java, Indonesia. A voucher specimen (BIT-1481) was deposited at the Herbarium of the Department of Biology, School of Life Sciences and Technology, Bandung Institute of Technology.

Table 3 Anti-inflammatory effects of selected compounds on LPS-stimulated BMDCs.

Compounds	IC ₅₀ values (μM)	
	IL-12 p40	IL-6
2	0.87 ± 0.06	1.54 ± 0.09
3	0.56 ± 0.01	1.22 ± 0.02
5	2.76 ± 0.08	> 5.0
6	0.41 ± 0.02	3.17 ± 0.23
8	1.40 ± 0.04	2.95 ± 0.17
9	0.35 ± 0.01	3.79 ± 0.10
10	0.39 ± 0.02	> 5.0
11	1.26 ± 0.02	2.07 ± 0.21
12	0.56 ± 0.03	2.27 ± 0.05
13	0.39 ± 0.05	3.71 ± 0.04
SB203580 ^a	5.00 ± 0.16	3.50 ± 0.12

^a SB203580 was used as a positive control

Extraction and isolation

The dried leaves of *P. aduncum* (3.2 kg) were extracted with 70% ethanol (10 L × 3 times) under reflux condition. Evaporation of the solvent under reduced pressure gave the crude extract (379 g), which was suspended in H₂O and successively separated with CH₂Cl₂ and EtOAc to yield CH₂Cl₂ (93 g) and EtOAc extracts (27 g).

The EtOAc extract (27 g) was fractionated on a silica gel CC eluting with gradient solvent systems of CH₂Cl₂-MeOH (0–100% MeOH, stepwise) to obtain six fractions (C.1 through C.6). Compound **12** (20 mg) was isolated from fraction C.3 by YMC reverse-phase (RP)-18 CC using MeOH-H₂O (1/1, v/v) as the eluent, and further purified by YMC RP-18 CC eluting with acetone-H₂O (1/2, v/v). Fraction C.4 was separated on silica gel CC eluting with CH₂Cl₂-MeOH-H₂O (4/1/0.1, v/v/v) to give six subfractions (C.4.1 through C.4.6). Compounds **2** (10 mg), **3** (70 mg), and **11** (38 mg) were separated from fraction C.4.2 by silica gel CC using CH₂Cl₂-MeOH (6/1, v/v) and further purified by Sephadex[®] LH-20 CC using MeOH-H₂O (1/1, v/v) as the eluent. Fraction C.4.4 was fractionated on YMC RP-18 CC eluting with MeOH-H₂O (1/2, v/v) to obtain three fractions (C.4.4.1 through C.4.4.3). Compounds **1** (12 mg) and **14** (10 mg) were obtained from fraction C.4.4.2 by Sephadex[®] LH-20 CC eluting with MeOH-H₂O (1/1, v/v). Fraction C.5 was isolated on YMC RP-18 CC eluting with MeOH-H₂O (1/2, v/v) to obtain four fractions (C.5.1 through C.5.4). Similarly, compound **13** (40 mg) was afforded from the fraction C.5.1 by Sephadex[®] LH-20 CC using MeOH-H₂O (1/1, v/v) as the eluent. Compounds **5** (8 mg), **9** (9 mg), and **10** (7 mg) were isolated from fraction C.5.2 by Sephadex[®] LH-20 CC eluting with MeOH-H₂O (1/1, v/v). And finally, fraction C.5.4 were separated by Sephadex[®] LH-20 CC using MeOH-H₂O (1/1, v/v) and further purified by silica gel CC using CH₂Cl₂-MeOH-H₂O (4/1/0.1, v/v/v) as eluents to afford compounds **4** (20 mg), **6** (10 mg), **7** (15 mg), and **8** (100 mg).

Acacetin 8-C-[β-D-apiofuranosyl-(1 → 2)-β-D-glucopyranoside] (1): Yellow amorphous powder; [α]_D²⁵: -65.3° (c 0.15, MeOH); IR (KBr) ν_{max}: 3367, 2925, 1650, 1602, 1561, 1496, 1070, and 1000 cm⁻¹; UV (MeOH) λ_{max} (log ε) 329 (4.39), 270 (4.31) nm; HR-ESI-MS m/z: 579.1726 [M + H]⁺ (Calcd. C₂₇H₃₁O₁₄ for 579.1714) and 601.1544 [M + Na]⁺ (Calcd. C₂₇H₃₀NaO₁₄ for 601.1533); ¹H NMR (600 MHz, CD₃OD) and ¹³C NMR (150 MHz, CD₃OD) are given in **Table 1**.




Fig. 2 A Effect of compounds **2**, **3**, **5**, **6**, and **8–13** on IL-12 p40 production by LPS-stimulated BMDCs. B Effect of compounds **2**, **3**, **5**, **6**, and **8–13** on IL-6 production by LPS-stimulated BMDCs. C Effect of compounds **2**, **3**, **5**, **6**, and **8–13** on TNF-α production by LPS-stimulated BMDCs. The data are presented as inhibition rate (%). Data represent the mean ± SD of at least three independent experiments performed in triplicate. Statistical significance is indicated as determined by one-way ANOVA followed by Dunnett's multiple comparison test, $p < 0.05$, using GraphPad Prism 6.

7-Methoxyacacetin 8-C-[β-D-apiofuranosyl-(1 → 3)-β-D-glucopyranoside] (2): Yellow amorphous powder; [α]_D²⁵: -75.8° (c 0.15, MeOH); IR (KBr) ν_{max}: 3375, 2931, 1650, 1602, 1570, 1523, 1459, 1350, and 1055 cm⁻¹; UV (MeOH) λ_{max} (log ε) 331 (4.42), 272 (4.32) nm. HR-ESI-MS m/z: 615.1685 [M + Na]⁺ (Calcd. C₂₈H₃₂NaO₁₄ for 615.1684); ¹H NMR (600 MHz, DMSO-*d*₆) and ¹³C NMR (150 MHz, DMSO-*d*₆) are given in **Table 1**.

7-Methoxyacacetin 8-C-[β-D-glucopyranosyl-(1 → 2)-β-D-glucopyranoside] (3): Yellow powder; [α]_D²⁵: -49.7° (c 0.1, MeOH); IR (KBr) ν_{max}: 3369, 2925, 1651, 1610, 1570, 1480, 1100, and 1040 cm⁻¹; UV (MeOH) λ_{max} (log ε) 332 (4.43), 268 (4.27) nm. HR-ESI-MS m/z: 623.1979 [M + H]⁺ (Calcd. C₂₉H₃₅O₁₅ for 623.1970), 645.1800 [M + Na]⁺ (Calcd. C₂₉H₃₄NaO₁₅ for 645.1790); ¹H NMR (600 MHz, DMSO-*d*₆) and ¹³C NMR (150 MHz, DMSO-*d*₆) are given in **Table 1**.

4''-O-Acetylacacetin 8-C-[α-L-rhamnopyranosyl-(1 → 2)-β-D-glucopyranoside] (4): Yellow powder; [α]_D²⁵: -56.0° (c 0.15, MeOH); IR (KBr) ν_{max}: 3414, 2936, 1704, 1650, 1570, 1490, 1446, 1350, and 1045 cm⁻¹; UV (MeOH) λ_{max} (log ε) 334 (4.44), 271 (4.28)

nm. HR-ESI-MS m/z : 635.1975 $[M + H]^+$ (Calcd. $C_{30}H_{35}O_{15}$ for 635.1970) and 657.1794 $[M + Na]^+$ (Calcd. $C_{30}H_{34}NaO_{15}$ for 657.1790); 1H NMR (600 MHz, CD_3OD) and ^{13}C NMR (150 MHz, CD_3OD) are given in **Table 2**.

Acid hydrolysis and sugar identification

Each compound (2.0 mg) was dissolved in 1.0 N HCl (dioxane/ H_2O , 1:1, v/v, 1.0 mL) and then heated to 80 °C in a water bath for 3.5 h. The acidic solution was neutralized with silver carbonate and the solvent thoroughly driven out under N_2 gas overnight. After extraction with ethyl acetate, the aqueous layer was first TLC analyzed [individual and co-analysis with standard sample: glucose (R_f 0.29), rhamnose (R_f 0.57), and apiose (R_f 0.59); $CHCl_3/MeOH/H_2O$, 3:2:0.3] and then concentrated to dryness using N_2 gas. The residue was dissolved in 0.1 mL of dry pyridine, and then L-cysteine methyl ester hydrochloride in pyridine (0.06 M, 0.1 mL) was added to the solution. The reaction mixture was heated at 60 °C for 2 h, and 0.1 mL of trimethylsilylimidazole solution was added, followed by heating at 60 °C for 1.5 h. The dried product was partitioned with *n*-hexane and H_2O (0.1 mL, each), and the organic layer was analyzed by GC; Column: SPB-1 (0.25 mm \times 30 m); detector FID, column temp 210 °C, injector temperature 270 °C, detector temperature 300 °C, carrier gas He. The absolute configuration of the monosaccharide was confirmed to be D-glucose, L-rhamnose, and D-apsiose by comparison of the retention time of the monosaccharide derivative (t_R 14.11, 4.50, and 4.67 min, respectively) with that of authentic sugar derivative samples prepared in the same manner (the retention times of standards D-glucose, L-rhamnose, and D-apsiose were 14.12, 4.50, and 4.68 min, respectively).

Cell culture and measurement of cytokine production

BMDCs were grown from wild-type C57BL/6 mice (Orient Bio Inc.) as previously described [28]. All animal procedures were approved and performed according to the guidelines of the Institutional Animal Care and Use Committee of Jeju National University (#2010-0028). Briefly, the mouse tibia and femur were obtained by flushing with Dulbecco's modified Eagle's medium to yield bone marrow cells. The cells were cultured in RPMI 1640 medium containing 10% heat-inactivated FBS (Gibco), 50.0 μ M β -mercaptoethanol, and 2 mM glutamine supplemented with 3% J558 L hybridoma cell culture supernatant containing granulocyte-macrophage colony stimulating factor. The culture medium was replaced with fresh medium every second day. On day 6 of the culture, non-adherent cells and loosely adherent DC aggregates were harvested, washed, and resuspended in RPMI 1640 supplemented with 5% FBS.

The BMDCs were incubated in 48-well plates in 0.5 mL containing 1×10^5 cells per well, and then treated with the isolated compounds at different concentrations for 1 h before stimulation with 10.0 ng/mL LPS from Salmonella Minnesota (Alexis). Supernatants were harvested 18 h after stimulation. Concentrations of compounds IL-12 p40, IL-6, and TNF- α in the culture supernatant were determined by ELISA (BD PharMingen) according to the manufacturer's instructions. SB203580 was the product of Calbiochem.

The inhibitory activity (*I*) was expressed as the inhibition rate (%), which was calculated from the following formula:

$$I = \frac{C_{dcv} - C_{dcc}}{C_{dcv}} \times 100$$

Where: *C*_{dcv}: Cytokine level (ng/mL) in vehicle-treated DC; *C*_{dcc}: Cytokine level (ng/mL) in compound-treated DC.

Statistical analysis

All data represent the mean \pm SD of at least three independent experiments performed in triplicate. Statistical significance is indicated as determined by one-way ANOVA followed by Dunnett's multiple comparison test, $p < 0.05$, using the GraphPad Prism 6.01 program (GraphPad Software Inc.).

Supporting information

Spectral data of compounds 1–4 are available as Supporting Information.

Acknowledgements

This study was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2009-0093815).

Conflict of Interest

The authors declare no conflict of interest.

Affiliations

- College of Pharmacy, Chungnam National University, Daejeon, Republic of Korea
- Institute of Marine Biochemistry (IMBC), Vietnam Academy of Science and Technology (VAST), Cau Giay, Hanoi, Vietnam
- Faculty of Medicine, Maranatha Christian University, Bandung, West Java, Indonesia
- Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung, West Java, Indonesia
- Department of Chemistry, Mathematic and Natural Sciences, University of Padjadjaran, Jatinangor Sumedang, West Java, Indonesia
- School of Medicine, Brain Korea 21 PLUS Program, and Institute of Medical Science, Jeju National University, Jeju, Republic of Korea

References

- Schwab JM, Serhan CN. Lipoxins and new lipid mediators in the resolution of inflammation. *Curr Opin Pharmacol* 2006; 6: 414–420
- Dinarelo CA. Proinflammatory cytokines. *Chest* 2000; 118: 503–508
- Okunade AL, Hufford CD, Clark AM, Lentz D. Antimicrobial properties of the constituents of *Piper aduncum*. *Phytother Res* 1997; 11: 142–144
- Morandim Ade A, Bergamo DCB, Kato MJ, Cavalheiro AJ, Bolzani Vda S, Furlan M. Circadian rhythm of anti-fungal prenylated chromene in leaves of *Piper aduncum*. *Phytochem Anal* 2005; 16: 282–286
- Morandim ADA, Kato MJ, Cavalheiro AJ, Furlan M. Intraspecific variability of dihydrochalcone, chromenes and benzoic acid derivatives in leaves of *Piper aduncum* L. (Piperaceae). *Afr J Biotechnol* 2009; 8: 2157–2162
- Orjala J, Wright AD, Behrends H, Folkers G, Sticher O, Rüggeger H, Rali T. Cytotoxic and antibacterial dihydrochalcones from *Piper aduncum*. *J Nat Prod* 1994; 57: 18–26
- Morandim AA, Bergamo DC, Kato MJ, Cavalheiro AJ, Bolzani VS, Furlan M. Circadian rhythm of anti-fungal prenylated chromene in leaves of *Piper aduncum*. *Phytochem Anal* 2005; 16: 282–286
- Baldoqui DC, Kato MJ, Cavalheiro AJ, Bolzani Vda S, Young MCM, Furlan M. A chromene and prenylated benzoic acid from *Piper aduncum*. *Phytochemistry* 1999; 51: 899–902
- Okunade AL, Hufford CD, Clark AM, Lentz D. Antimicrobial properties of the constituents of *Piper aduncum*. *Phytother Res* 1997; 11: 142–144
- Piccolo CRD, Bezerra MP, Gomes KS, Passero LFD, Laurenti MD, Martins EGA, Sartorellia P, Lago JHG. Antileishmanial activity evaluation of adunchalcone, a new prenylated dihydrochalcone from *Piper aduncum* L. *Fitoterapia* 2014; 97: 28–33
- Orjala J, Erdelmeier CA, Wright AD, Rali T, Sticher O. Five new prenylated *p*-hydroxybenzoic acid derivatives with antimicrobial and mollusci-

- cidal activity from *Piper aduncum* leaves. *Planta Med* 1993; 59: 546–551
- 12 Parise-Filho R, Pastrello M, Camerlingo CEP, Silva GJ, Agostinho LA, de Souza T, Magri FMM, Ribeiro RR, Brandt CA, Polli MC. The anti-inflammatory activity of dillapiole and some semisynthetic analogues. *Pharm Biol* 2011; 49: 1173–1179
- 13 Cheng G, Bai Y, Zhao Y, Tao J, Liua Y, Tu G, Ma L, Liao N, Xu X. Flavonoids from *Ziziphus jujuba* Mill var. *spinosa*. *Tetrahedron* 2000; 56: 8915–8920
- 14 Khallouki F, Hmamouchi M, Younos C, Soulimani R, Essassic EM. A new flavonoid from the aerial parts of *Chrysanthemum viscidhirtum*. *Fitoterapia* 2000; 71: 413–416
- 15 BJORØY Ø, RAYYAN S, FOSSEN T, KALBERG K, ANDERSEN ØM. C-glycosylanthocyanidins synthesized from C-glycosylflavones. *Phytochemistry* 2009; 70: 278–287
- 16 Larionova M, Spengler I, Nogueira C, Quijano L, Ramírez-Gualito K, Cortés-Guzmán F, Cuevas G, Calderón JS. A C-glycosylflavone from *Piper ossanum*, a compound conformationally controlled by CH/ π and other weak intramolecular interactions. *J Nat Prod* 2010; 73: 1623–1627
- 17 Kiem PV, Cuong NX, Nhiem NX, Thu VK, Ban NK, Minh CV, Tai BH, Hai TN, Lee SH, Jang HD, Kim YH. Antioxidant activity of a new C-glycosylflavone from the leaves of *Ficus microcarpa*. *Bioorg Med Chem Lett* 2011; 21: 633–637
- 18 Rayyan S, Fossen T, Nateland HS, Andersen ØM. Isolation and identification of flavonoids, including flavone rotamers, from the herbal drug 'Crataegi Folium Cum Flore' (Hawthorn). *Phytochem Anal* 2005; 16: 334–341
- 19 Sharaf M, El-Ansari MA, Matlin SA, Saleh NAM. Four flavonoid glycosides from *Peganum harmala*. *Phytochemistry* 1997; 44: 533–536
- 20 Hilsenbeck RA, Mabry TJ. C-glycosylflavones from *Siphonoglossa sessilis*. *Phytochemistry* 1983; 22: 2215–2217
- 21 Kato T, Morita Y. C-glycosylflavones with acetyl substitution from *Rumex acetosa* L. *Chem Pharm Bull* 1990; 38: 2277–2280
- 22 Nicholls KW, Bruce AB. Flavonoids of *Lupinus arboreus*. *Phytochemistry* 1979; 18: 1078
- 23 Markham KR, Whitehouse LA. Unique flavonoid glycosides from the New Zealand white pine, *Dacrycarpus dacrydioides*. *Phytochemistry* 1984; 23: 1931–1936
- 24 Wei WC, Sung PJ, Duh CY, Chen BW, Sheu JH, Yang NS. Anti-inflammatory activities of natural products isolated from soft corals of Taiwan between 2008 and 2012. *Mar Drugs* 2013; 11: 4083–4126
- 25 Lee JC, Laydon JT, McDonnell PC, Gallagher TF, Kumar S, Green D, McNulty D, Blumenthal MJ, Heys JR, Landvatter SW, Strickler JE, Mclaughlin MM, Siemens IR, Fisher SM, Livi GL, White JR, Adams JL, Young PR. A protein kinase involved in the regulation of inflammatory cytokine biosynthesis. *Nature* 1994; 372: 739–746
- 26 Xiaoa J, Capanoglu E, Jassbi AR, Miron A. Advance on the flavonoid C-glycosides and health benefits. *Cri Rev Food Sci Nut*, advance online publication 13 October 2015; DOI: 10.1080/10408398.2015.1067595
- 27 Borghi SM, Carvalho TT, Staurengo-Ferrari L, Hohmann MSN, Pinge-Filho P, Casagrande R, Verri jr. WA. Vitexin inhibits inflammatory pain in mice by targeting TRPV1, oxidative stress, and cytokines. *J Nat Prod* 2013; 76: 1141–1149
- 28 Koo JE, Hong HJ, Dearth A, Kobayashi KS, Koh YS. Intracellular invasion of orientia tsutsugamushi activates inflammasome in ASC-dependent manner. *PLoS One* 2012; 7: e39042



LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau *PEER REVIEW*

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Anti-inflammatory Flavonoid C-Glycosides from Piper aduncum Leaves
 Jumlah Penulis : 10 Orang
 Nama-nama Penulis : Nguyen Phuong Thao, Bui Thi Thuy Luyen, **Wahyu Widowati**, Nurul Fauziah, Maesaroh Maesaroh, Tati Herlina, Zahid Manzoor, Irshad Ali, Young Sang Koh, Young Ho Kim.
 Status Penulis : **Penulis Pertama** / Penulis ke 3 / **Penulis Korespondensi ****
 Identitas Jurnal Ilmiah : a. Nama jurnal : Planta Medica
 b. Nomor ISSN : 0032-0943
 c. Vol., No., Bulan, Tahun : Vol 82; No 17; 2016;
 d. Penerbit : George thieme Verlag KG
 e. DOI Artikel (jika ada) : 10.1055/s-0042-108737
 f. Alamat Web Jurnal : www.thieme-connect.com/products
 g. Terindeks di : Scopus Q1, SJR 0.422

Kategori Publikasi Jurnal Ilmiah: **Jurnal Ilmiah Internasional / Internasional Bereputasi ****
 (beri tanda \checkmark yang dipilih)
 Jurnal Ilmiah Nasional Terakreditasi
 Jurnal Ilmiah Nasional / Nasional terindeks ***

HASIL PENILAIAN (*Peer Review*) :


No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi <input checked="" type="checkbox"/>	Nasional Terakreditasi <input type="checkbox"/>	Nasional *** <input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,5
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			4,8
c.	Kecukupan dan kemitakhiran data/informasi dan metodologi (30%)	12			4,5
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			4,4
	Total 100%	40			38,2

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur... *Penulisan memenuhi standard karya ilmiah akademis, metoda yang digunakan sesuai dan tepat digunakan pada penelitian ini*
- b. Ruang lingkup & kedalaman pembahasan... *Penelitian tentang isolasi, senyawa baru dari Piper aduncum sebagai bahan antiinflamasi*
- c. Kecukupan & kemitakhiran data serta metodologi... *Ida dasar penelitian baik membahas mekanisme senyawa baru menghambat IL-12, IL-16, TNF- α pada sel BMDC yang diinduksi LPS*
- d. Kelengkapan unsur dan kualitas penerbit... *Planta Medica terindeks Scopus Q1 SJR 0,422. Penerbit George Thieme Verlag KG*
- e. Indikasi plagiasi... *Tidak terdapat indikasi plagiarisme atau self plagiarisme*

f. Kesesuaian bidang ilmu Paper bidang kimia organik, biokimia kurang sesuai dengan bidang ilmu penerap.

REVIEWER 1



(Prof. Dr. Chrismis Novalinda Ginting, M.Kes)

NIK : 0115127801

UNIVERSITAS PRIMA INDONESIA

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau PEER REVIEW

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Anti-inflammatory Flavonoid C-Glycosides from Piper aduncum Leaves
 Jumlah Penulis : 10 Orang
 Nama-nama Penulis : Nguyen Phuong Thao, Bui Thi Thuy Luyen, **Wahyu Widowati**, Nurul Fauziah, Maesaroh Maesaroh, Tati Herlina, Zahid Manzoor, Irshad Ali, Young Sang Koh, Young Ho Kim.
 Status Penulis : ~~Penulis Pertama~~ / Penulis ke 3 / ~~Penulis Korespondensi **~~
 Identitas Jurnal Ilmiah : a. Nama jurnal : Planta Medica
 b. Nomor ISSN : 0032-0943
 c. Vol., No., Bulan, Tahun : Vol 82; No 17; 2016;
 d. Penerbit : George thieme Verlag KG
 e. DOI Artikel (jika ada) : 10.1055/s-0042-108737
 f. Alamat Web Jurnal : www.thieme-connect.com/products
 g. Terindeks di : Scopus Q1, SJR 0.422

Kategori Publikasi Jurnal Ilmiah: ~~Jurnal Ilmiah Internasional~~ / Internasional Bereputasi **)
 (beri tanda \surd yang dipilih)
 Jurnal Ilmiah Nasional Terakreditasi
 Jurnal Ilmiah Nasional / Nasional terindeks ***)

HASIL PENILAIAN (Peer Review) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi <input checked="" type="checkbox"/>	Nasional Terakreditasi <input type="checkbox"/>	Nasional ***) <input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,5
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			4,7
c.	Kecukupan dan kemutakhiran data/ informasi dan metodologi (30%)	12			11,7
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			11,4
	Total	100%	40		36,3

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur
Artikel ini ditulis sudah sesuai dengan kaidah penulisan dan penyusunan serta unsur dan isinya sudah memiliki kesesuaian.
- b. Ruang lingkup & kedalaman pembahasan
Ruang lingkup bahasan sudah memailui dan ada kedalaman dalam pembahasannya dan urutannya.
- c. Kecukupan & kemutakhiran data serta metodologi
Secara umum metodologi yang dipakai sudah tercukupi dan sudah mutakhir
- d. Kelengkapan unsur dan kualitas penerbit
Unsur-unsur dalam artikel ini sudah lengkap dan kualitas penerbit sudah masuk kedalam kategori bereputasi yang baik.
- e. Indikasi plagiasi

Tidak ada ditemukannya indikasi plagiasi

f. Kesesuaian bidang ilmu

Artikel ini sesuai dengan bidang ilmu yang diteliti penulis.

REVIEWER 2



(Prof. Dr. Ermi Girsang, M. Kes)

NIK : 0117057501

UNIVERSITAS PRIMA INDONESIA

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau *PEER REVIEW*

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Anti-inflammatory Flavonoid C-Glycosides from Piper aduncum Leaves
 Jumlah Penulis : 10 Orang
 Nama-nama Penulis : Nguyen Phuong Thao, Bui Thi Thuy Luyen, **Wahyu Widowati**, Nurul Fauziah, Maesaroh Maesaroh, Tati Herlina, Zahid Manzoor, Irshad Ali, Young Sang Koh, Young Ho Kim.
 Status Penulis : ~~Penulis Pertama~~ / Penulis ke 3 / ~~Penulis Korespondensi~~ **)
 Identitas Jurnal Ilmiah : a. Nama jurnal : Planta Medica
 b. Nomor ISSN : 0032-0943
 c. Vol., No., Bulan, Tahun : Vol 82; No 17; 2016;
 d. Penerbit : George thieme Verlag KG
 e. DOI Artikel (jika ada) : 10.1055/s-0042-108737
 f. Alamat Web Jurnal : www.thieme-connect.com/products
 g. Terindeks di : Scopus Q1, SJR 0.422

Kategori Publikasi Jurnal Ilmiah: ~~Jurnal Ilmiah Internasional~~ / Internasional Bereputasi **)
 (beri tanda \checkmark yang dipilih)
 Jurnal Ilmiah Nasional Terakreditasi
 Jurnal Ilmiah Nasional / Nasional terindeks ***)

HASIL PENILAIAN (*Peer Review*) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi <input checked="" type="checkbox"/>	Nasional Terakreditasi <input type="checkbox"/>	Nasional ***) <input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,5
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			11,75
c.	Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12			11,6
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			11,4
Total		40			38,25

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur. *Penulisan memenuhi standard karya ilmiah akademis, metode yang digunakan sesuai dan tepat digunakan pada penelitian ini. Artikel ini ditulis sudah sesuai dengan kaidah penulisan dan penyusunan serta unsur dan lainnya sudah memiliki kesesuaian*
- b. Ruang lingkup & kedalaman pembahasan. *Penelitian tentang isolasi, senyawa baru dari Piper aduncum sebagai bahan antiinflamasi*

Ruang lingkup bahasan sudah memadai dan ada kedalaman dalam pembahasan dan analisisnya

c. Kecukupan & kemutakhiran data serta metodologi. Ide dasar penelitian baik membahas mekanisme senyawa baru menghambat IL-12, IL-16, TNF- α pada sel BMDC yang dirangsang LPS

Caranya umum metodologi yang dipakai sudah terakurasi dan sudah mutakhir

d. Kelengkapan unsur dan kualitas penerbit

Planta Medica terindeks Scopus Q1 SJR 0,422. Penerbit George Thieme Verlag KG

Unsur-unsur dalam artikel ini sudah lengkap dan kualitas penerbit sudah masuk kedalam kategori bereputasi yang baik

e. Indikasi plagiaris

Tidak terdapat adanya indikasi plagiaris atau self plagiaris

Tidak ada ditemukannya indikasi plagiaris

f. Kesesuaian bidang ilmu

Paper bidang kimia organik, biokimia kurang sesuai dengan bidang ilmu penulis

Artikel ini sudah sesuai dengan bidang ilmu yang dilakoni penulis

Medan,
Reviewer 2



(Prof. Dr. Ermi Girsang, M.Kes)

NIK : 0117057501

UNIVERSITAS PRIMA INDONESIA

Medan,
Reviewer 1



(Prof. Dr. Chrismis Novalinda Ginting, M.Kes)

NIK : 0115127801

UNIVERSITAS PRIMA INDONESIA