




JOURNAL OF NATURAL REMEDIES

Dedicated to Medicinal Plant Research

ISSN : 0972 - 5547





Editorial Board

Editor-In-Chief

Dr. M. K. Mohan Maruga Raja

Senior Scientist, TanBio R&D Solution, Penyar Technology Business Incubator, Periyar Maniammal University, Thanjavur, Tamil Nadu

Co-Editors

Dr. D. Prashanth

Natural Remedies, Bangalore

Balu Ranganathan

Palms Connect LLC, Sandy, Utah, USA

Dr. M. K. Kathiravan

Department of Pharmaceutical Chemistry, SRM University, Kattankulathur, Tamil Nadu

Dr. Jollus Gimbut

Deputy Director of Centre of Excellent for Advanced Research in Fluid Flow (CARIFF), Assoc. Professor, Faculty of Chemical and Natural Resources Engineering, The Gimbut Group, Universiti Malaysia Pahang, Gambang, Pahang, Malaysia

Editorial Board Members

Dr. Acharya Sanjeev R.

Institute of Pharmacy, Nirma University, Ahmedabad

Dr. Akula Chinapolaiah

ICAR Indian Council of Agricultural Research, Anand Gujarat

Dr. Anant Bhandarkar

Goa College of Pharmacy, Govt. of Goa

Dr. Anil Kumar Uttamchand Tatiya

R.C.Patel College of Pharmacy Shirgur

Dr. Arun B. Joshi

Goa College of Pharmacy, Panaji, Goa

Dr. Chandrashekar K. S.

Manipal College of Pharmaceutical Sciences, Manipal University, Manipal

Dr. Dayanand M. Kannur

SCCES'S Indira College of Pharmacy, Pune

Dr. Deepak Kumar Semwal

Uttarakhand Ayurved University, Dehradun

Dr. Kuntal Das

Krupanidhi College of Pharmacy, Bangalore

Dr. Laxminarayana Baiyy

Kasturba Medical College, Manipal

Dr. Madhavan V.

M.S.Ramaiah College of Pharmacy, Bangalore

Dr. Manik Ghosh

Birla Institute of Technology, Mesra, Ranchi

Dr. Manjunath Shetty

Manipal College of Pharmaceutical Sciences, Manipal University, Manipal

Dr. Mohd. Yaseen

Central Institute of Medicinal and Aromatic Plants(CIMAP), Lucknow

Dr. Mruthunjaya K.

JSS College of Pharmacy, Mysore

Dr. Nagarajan A.

PSG College of Pharmacy, Coimbatore

Dr. Peter Achunike Akah

University of Nigeria, Nsukka, Nigeria

Dr. Rajendra Gyawali

Department of Pharmacy, Kathmandu

Dr. Sathianarayanan S.

Amrita School of Pharmacy Amrita Vishwavidyapeetram University, Kochi

Dr. Shailendra S. Gurav

Goa College of Pharmacy, Panaji, Goa

Dr. Sinha B. N.

Birla Institute of Technology, Mesra, Ranchi

Dr. Subhash C Mandal

Jadavpur University, Kolkata

Dr. Vasantha Kumar

ICAR, Indian Institute of Horticultural Research, Bengaluru

Dr. Vidhu Aeri

Jamia Hamdani Deemed University, New Delhi

Dr. Vikram Chauhan

Planet Ayurveda Clinic and Krishna Herbal Company, Mohali, India

Dr. Vimal Kumar

Institute of Pharmacy, Nirma University, Ahmedabad

Dr. Padmaa M Paarakh

Pharmacognostical, Phytochemical and Pharmacological studies, Banaras Hindu University, Varanasi

Prof.(Dr.) R. S. Tomar

Director, Amity Institute of Biotechnology, Madhya Pradesh

Manish K. R. Singh

Department of Poultry Science, Duvasu, Mathura, Uttar Pradesh

Dr. N. Kasthuri Bai

Interdisciplinary Institute of Indian System of Medicine, Tamil Nadu, India

Dr. Mani Ganesh

Hanseon University, Chungcheongnam-do, South Korea

Editorial Manager

Kavitha Kumar

Informatics Publishing Ltd.

Journal Coordinator

Priyadarshini C

Informatics Publishing Ltd.

Quick Links

- [Abstracting and Indexing](#)
- [Instructions to Authors](#)
- [Pre-submission checklist](#)
- [Publication Fees](#)
- [Journal Policies and Ethics](#)
- [Submission Process](#)
- [Information for Reviewers](#)
- [Email Alerts](#)
- [CrossMark Policy](#)

Archives

[Expand All]

[2019](#)

[2018](#)

[2017](#)

[2016](#)

[2015](#)

[2014](#)

[2013](#)

[2012](#)

[More..](#)


 All Search

- Home
- Current
- Archives
- Authors
- Institutions
- Editorial Board
- Submit Article
- Online First

Volume 16, Issue 2, April 2016

Table of Contents

Articles

Evaluation of Antipyretic and Anti-inflammatory Activity of Aqueous Extract of *Leptadenia Reticulata* in Animal Models

Sneha Bherji, M. Ganga Raju, Namile Divya

DOI: 10.18311/jnr/2016/468, Pagination: 40-44

[Full-Text PDF](#)

Abstract Views: 447 | Pdf Views: 392

An Innovation for Retarded Healing Process of a Chronic Ulcer by Aloe Vera Gel Treatment

Majid Avijgan, Mahtab Avijgan, Atousa Hakamifard, Niloufar Razavi

DOI: 10.18311/jnr/2016/479, Pagination: 45-51

[Full-Text PDF](#)

Abstract Views: 527 | Pdf Views: 558

Cytotoxic Activity of Mangosteen (*Garcinia mangostana* L.) Peel Extract and α -mangostin toward Leukemia Cell Lines (HL-60 and K-562)

Arina Novilla, Dedi S. Djamhuri, Nurul Fauziah, Maesaroh Maesaroh, Balqis Balqis, Wahyu Widowati

DOI: 10.18311/jnr/2016/842, Pagination: 52-59

[Full-Text PDF](#)

Abstract Views: 628 | Pdf Views: 673

Tea Polyphenols as Natural Products for Potential Future Management of HIV Infection - an overview

Hamid Reza Sodagari, Roodabeh Bahramsoltani, Mohammad Hosein Farzaei, Amir Hossein Abdolghaffari, Nima Rezaei, Andrew W. Taylor-Robinson

DOI: 10.18311/jnr/2016/4782, Pagination: 60-72

[Full-Text PDF](#)

Abstract Views: 681 | Pdf Views: 560

Anti-Inflammatory Potential of Gandarusa (*Gendarussa vulgaris* Nees) and Soursoup (*Annona muricata* L) Extracts in LPS Stimulated-Macrophage Cell (RAW264.7)

Dian Ratih Laksmiawati, Ajeng Prima Prasanti, Nadia Larasinta, Gloria Agitha Syauta, Rivanny Hilda, Hesty Utami Ramadaniati, Anisa Widyastuti, Nadia Karami, Merry Afni, Dwi Davidson Rihibiha, Hanna Sari W. Kusuma, Wahyu Widowati

DOI: 10.18311/jnr/2016/5367, Pagination: 73-81

[Full-Text PDF](#)

Abstract Views: 816 | Pdf Views: 391

Quick Links

- [Abstracting and Indexing](#)
- [Instructions to Authors](#)
- [Pre-submission checklist](#)
- [Publication Fees](#)
- [Journal Policies and Ethics](#)
- [Submission Process](#)
- [Information for Reviewers](#)
- [Email Alerts](#)
- [CrossMark Policy](#)

Archives

[\[Expand All\]](#)

[2019](#)

[2018](#)

[2017](#)

[2016](#)

[2015](#)

[2014](#)

[2013](#)

[2012](#)

[More...](#)

Current Issue

ISI	1.0
ISI	2.0
ISI	1.0



Cytotoxic Activity of Mangosteen (*Garcinia mangostana* L.) Peel Extract and α -Mangostin toward Leukemia Cell Lines (HL-60 and K-562)

Arina Novilla¹, Dedi S. Djamhuri¹, Nurul Fauziah², Maesaroh Maesaroh², Balqis Balqis² and Wahyu Widowati^{3*}

¹School of Health Sciences Jenderal Achmad Yani, Cimahi, Indonesia

²Biomedical and Biomolecular Research Centre, Aretha Medika Utama, Bandung, Indonesia

³Faculty of Medicine, Maranatha Christian University, Bandung, Indonesia; wahyu_w60@yahoo.com

Abstract

Fruit of Mangosteen (*Garcinia mangostana* L.) is well-known in Indonesia and other Southeast Asian countries. Studies have shown that extract of the pericarp of mangosteen contained mostly of xanthenes exhibit many biological activities, especially as an antitumor. This study aimed to investigate the cytotoxic activity and selectivity of Mangosteen Peel Extract (MPE) and α -mangostin against two leukemia cell lines (HL-60 and K-562) and the normal lymphocyte cells from two different donors. The cytotoxic activity was performed using 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium (MTS) assay. Imatinib and Isotretinoin were used as a positive control to the K-562 and HL-60 cells, respectively. The MPE and α -mangostin revealed higher mortality toward leukemia cell lines rather than toward lymphocyte cells, with more than 80% of HL-60 and K-562 cells died at 6.25 and 25 μ g/ml, respectively. MPE was more toxic and selective against K-562 with IC50 of 2.79 μ g/ml and SI of 8.27, while α -mangostin was more toxic and selective against HL-60 with IC50 of 1.12 μ g/ml and SI of 22.34. MPE and α -mangostin showed potent sensitivity and selectivity to leukemia cells, hence these are considered as promising sources for future leukemia treatment.

Keywords: Mangosteen Peel Extract, α -mangostin, acute myeloid leukemia (AML), chronic myeloid leukemia (CML), selectivity index

1. Introduction

Cancer has been the second cause of death after cardiovascular diseases in the world. There were around 8.2 million of people died because of cancer and about 14.1 million of new cancer cases were detected in 2012¹. Particularly in Indonesia, there was about 0.1% of total people suffering from cancer with 74.6% of mortality rate

in male and 63.3% in female². Lung or bronchus cancer, prostate cancer in men, breast cancer in the female, and colon or rectum cancer reveal as the most deadly cancer worldwide. However, rare cancer such as leukemia is also a cause of death because of its destructive nature³. According to the GLOBACAN, the percentage of leukemia was 2.5% of total new cases, contributing to 3.2% of total deaths by cancer in 2012¹ which was mostly suffered by children⁴.

*Author for correspondence

Email: wahyu_w60@yahoo.com

Characterized by high numbers of abnormal white blood cells, leukemia has been grouped into lymphoblastic and myeloid leukemia, namely Chronic Lymphocytic Leukemia (CLL) or Acute Lymphocytic Leukemia (ALL) and Chronic Myelogenous Leukemia (CML) or acute Myelogenous Leukemia (AML). Leukemia can be treated using interferon- α (IFN- α), Stem Cell Transplantation (SCT), radiotherapy and chemotherapy. Imatinib, a chemotherapy agent of tyrosine-kinase inhibitor, has been used as the first-line treatment for CML. It induces apoptosis and inhibition of proliferation by inhibiting the phosphorylation and binding process of Bcr-Abl oncoprotein⁵. Another chemotherapy agent for leukemia is retinoic acid, vitamin A derivative such as All-Trans Retinoic Acid (ATRA) and isotretinoin. They are used for several cancer treatment and remission including for Acute Promyelocytic Leukemia (APL), a subtype of AML by inducing cell apoptosis⁶. However, the usage of imatinib and retinoic acid as well as the other chemotherapy agents can generate side effects by damaging the normal cells and lead to the development of the multidrug resistance cancer⁷. Therefore, the finding of the safe, selective, and sensitive chemicals that have anticancer property is still needed.

Mangosteen (*Garcinia mangostana* L.) has been used as a traditional medicine to treat diseases related to immune and gastrointestinal for decades. Its extract contains several active compounds, including terpenes, anthocyanins, tannins, phenols, and xanthenes⁸. Xanthenes have been shown to have several pharmacological properties, including as an antitumor⁹. The most abundant and the most studied xanthenes are α -mangostin, β -mangostin, γ -mangostin, garcinone E, and gartanin, and those xanthenes possess great biological activities such as anti-allergy, anti-inflammatory, antioxidant, anti-tumor, antibacterial, antifungal and antiviral activities, neuroprotective, cardioprotective, and immunomodulation¹⁰⁻¹². Prior studies revealed that α -mangostin were the most abundance xanthone derivative from the aqueous, acetone and methanol extract of mangosteen^{9,13}. Therefore, this study aimed to investigate the cytotoxic activity and selectivity of Mangosteen Peel Extract (MPE) and α -mangostin against two leukemia cell lines (HL-60 and K-562) and the normal lymphocyte cell from two different donors. The 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium MTS) assay was used to investigate the cytotoxic effect of the MPE, α -mangostin, imatinib, and isotretinoin. Those data are used to reveal the IC₅₀ and Selectivity Index

(SI) of the MPE and three other compounds to the HL-60 and K-562 cell lines. The *in vitro* result demonstrates that MPE and α -mangostin have a potential to use against leukemia by showing the higher cytotoxic effect and selective index to leukemia rather than the imatinib and isotretinoin.

2. Materials and Methods

2.1 Mangosteen Peel Extract (MPE) Preparation

The peels of *Garcinia mangostana* L. from Cisalak-Subang, West Java, Indonesia were determined by the herbarium staff of Biology Department, School of Life Sciences and Technology, Bandung Institute of Technology, Bandung, West Java, Indonesia. Around 350 g of chopped and dried-mangosteen peels were extracted using steady-state extraction (maceration) method. The chopped and dried peels were immersed in ethanol 70% for 24 hours and filtrated. The filtration was repeated until the filtrate became colorless. The filtrate was evaporated with a rotary evaporator at 40°C and the ethanolic extract of mangosteen peel was collected and stored in a freezer with the temperature of -20°C^{12,13}.

2.2 Leukimia Cell Lines Cultivation

Two different leukemia cell lines, human acute promyelocytic (HL-60) and human chronic myelogenous (K-562) were obtained from Stem Cells and Cancer Institute (SCI) Jakarta, Indonesia. Cell lines were cultured in supplemented Iscove's Modified Dulbecco's Medium (IMDM) (Biowest, France), 2% Penicillin-Streptomycin (Biowest, France), 10% Fetal Bovine Serum (FBS) (Biowest, France) and were incubated at 37°C in a humidified incubator with 5% CO₂. After 24 hours of incubation, the number of viable cells was counted using a haemocytometer with trypan blue staining to obtain a sufficient cell number for the cytotoxic assay.

2.3 Lymphocyte Isolation and Cultivation

Lymphocyte isolation from two healthy donors was approved by Ethic Commission Number 232/KEP/I/2016 of Maranatha Christian University-Immanuel Hospital Bandung, Indonesia. The healthy donors have signed the informed consent. After taken from donors, to separate

the lymphocyte from other leukocytes histopaque (Sigma-Aldrich, U.S.A.) was added into the whole blood. Whole blood then was centrifuged at 1,800 rpm for 5 minutes to obtain the blood buffy coat. The blood buffy coat was added with 1x Phosphate Buffer Saline (PBS) (Biowest, France) and was centrifuged at 1,800 rpm for 10 minutes in 4°C. Then, the pellet was added with 1x PBS and was centrifuged at 1,500 rpm for 10 minutes in 4°C. The final pellet contained leukocytes was maintained in Roswell Park Memorial Institute Medium (RPMI) (Biowest, France) and supplemented with 1% non-essential amino acids, 1% l-glutamine, 100 U/ml penicillin, 10mg/mL streptomycin and 10% heat-inactivated fetal calf serum (Biowest, France) then incubated at 37°C in humidified air containing 5% CO₂¹⁴.

2.4 Cytotoxic Assay

Approximately 100 µL of lymphocyte, HL-60 and K-562 cell suspension contained 10⁵ cells/ml were seeded in each well of 96-well plate. Following that, 100 µL of Isotretinoin (Roche, Switzerland) in different levels (100, 85, 70, 55, 40, and 25 µM) were added into the wells of lymphocyte and HL-60. Imatinib (Glivec, Novartis) in different levels (100, 50, 25, 12.5, 6.25, and 3.125 µM) were added into the wells of lymphocyte and K-562. α-mangostin (Cengdu Biopurify Phytochemical, China) (100, 50, 25, 12.5, 6.25, and 3.125 µM) and Mangosteen Peel Extract (MPE) (100, 50, 25, 12.5, 6.25, and 3.125 µg/mL) were added into the wells of lymphocytes, HL-60, and K-562. Cells then incubated at 37°C with 5% of CO₂ for 24 hours. Then, 20 µL of 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxyme-thoxyphenyl)-2-(4-sulphophenyl)-2H-tetrazolium (MTS) (Promega, U.S.A.) was added into each well and plate was incubated at 5% CO₂, 37°C for 3 hours. Finally, the absorbance used to calculate the percentage of cell mortality (Equation 1) was measured at 490 nm with Multiskan GO (Thermo Scientific, U.S.A.). Wells of each cell that not supplemented with the compounds were presented as the negative control.

$$\text{Cell Mortality (\%)} = \frac{(A_{\text{untreated cell}} - A_{\text{treated cell}})}{A_{\text{untreated cell}}} \times 100\% \quad (1)$$

$A_{\text{treated cell}}$: Absorbance of cells treated with either MPE, α-mangostin, imatinib, or isotretinoin;

$A_{\text{untreated cell}}$: Absorbance of untreated cells

2.5 Selectivity Analysis

The Selectivity Index (SI) describes the selectivity of the MPE, α-mangostin, imatinib, and isotretinoin toward the leukemic cell lines. The index was calculated from IC₅₀ ratio of the lymphocytes and the cancerous cell line (HL-60 or K-562) (Equation 2)¹⁵. The compound with SI higher than 3 revealed a high selective compound against the HL-60 or K-562, and the opposite was a less selective compound.

$$\text{Selectivity Index (SI)} = \frac{\text{IC}_{50} \text{ of normal cell}}{\text{IC}_{50} \text{ of cancer cell}} \quad (2)$$

IC₅₀ of normal cell: The value of median inhibitory concentration (IC₅₀) of lymphocyte cells;

IC₅₀ of cancer cell: The value of median inhibitory concentration (IC₅₀) of cancer cells

2.6 Statistical Analysis

All treatment in this study was done independently and conducted in triplicate. SPSS Statistic 17.0 for Windows was used to calculate the IC₅₀ of the compounds and the significant differences using probit analysis and one-way ANOVA, respectively. The mortality and viability data were presented as mean ± Standard Deviation (SD).

3. Results

3.1 Cytotoxic Effect of Each Compound toward HL-60, K-562, and Leukocyte

Table 1 to 4 show the cytotoxic effect of imatinib, isotretinoin, α-mangosteen and MPE presented as the percentage mortality of HL-60, K-562, and leukocyte. Isotretinoin and α-mangostin started at the concentration of 25 µM had more than 50% mortality to HL-60 and K-562, respectively. Imatinib and α-mangostin started at 12.5 µM showed more than 50% mortality to K-562 and HL-60, respectively (Table 1-3). Those of MPE started at 3.125 µg/ml and at 6.25 µg/ml revealed more than 50% mortality to HL-60 and K-562, respectively (Table 4). Furthermore, Figure 1 shows HL-60 and K-562 cell lines have lower density in a higher concentration of MPE and α-mangostin treatments.

Table 1: The percentage mortality (%) of Lymphocyte and K-562 treated with Imatinib in six different concentrations

Cells	Concentration (μ M)					
	100	50	25	12.5	6.25	3.125
Lymphocyte	74.67 \pm 11.09 ^b	10.71 \pm 5.82 ^a	6.09 \pm 6.48 ^a	5.67 \pm 6.57 ^a	-5.81 \pm 1.92 ^a	-22.88 \pm 10.86 ^a
K-562	94.64 \pm 0.11 ^f	76.89 \pm 1.01 ^e	63.43 \pm 0.74 ^d	55.95 \pm 0.99 ^c	51.95 \pm 0.9 ^b	47.5 \pm 0.8 ^a

Data is presented as mean \pm standard deviation, different superscript letters in each row (Lymphocyte, K-562) showed a significant difference at $p < 0.05$ (Tukey HSD post hoc test) among the concentrations.

Table 2: The percentage mortality (%) of Lymphocyte and HL-60 treated with Isotretinoin in six different concentrations

Cells	Concentration (μ M)					
	100	85	70	55	40	25
Lymphocyte	91.74 \pm 12.07 ^b	60.39 \pm 17.70 ^b	54.93 \pm 34.16 ^b	39.4 \pm 14.56 ^{ab}	15.05 \pm 4.31 ^{ab}	-38.14 \pm 11.43 ^a
HL-60	99.05 \pm 1.11 ^e	95.41 \pm 2.19 ^{de}	83.48 \pm 1.02 ^{cd}	80.53 \pm 2.48 ^{bc}	69.52 \pm 4.66 ^b	51.73 \pm 2.86 ^a

Data is presented as mean \pm standard deviation, different superscript letters in each row (Lymphocyte, HL-60) showed a significant difference at $p < 0.05$ (Tukey HSD post hoc test) among the concentrations.

Table 3: The percentage mortality (%) of Lymphocyte, HL-60, and K-562 treated with α -Mangosteen in six different concentrations

Cells	Concentration (μ M)					
	100	50	25	12.5	6.25	3.125
Lymphocyte	67.53 \pm 25.83 ^{aA}	49.9 \pm 1.88 ^{aA}	33.8 \pm 26.82 ^{aA}	29.32 \pm 7.07 ^{aA}	22.6 \pm 37.28 ^{aA}	19.38 \pm 1.11 ^{aB}
HL-60	95.72 \pm 0.43 ^{dA}	94.55 \pm 0.07 ^{dB}	94.51 \pm 0.65 ^{dA}	75.1 \pm 0.27 ^{cB}	-29.44 \pm 0.5 ^{bA}	-45.36 \pm 7.4 ^{aA}
K-562	96.2 \pm 0.11 ^{dA}	94.65 \pm 0.3 ^{dB}	91.29 \pm 0.27 ^{dA}	16.31 \pm 0.65 ^{cA}	-21.72 \pm 3.08 ^{bA}	-68.47 \pm 8.35 ^{aA}

Data is presented as mean \pm standard deviation, different lowercase of superscript letters in each row (Lymphocyte, K-562, HL-60) showed a significant difference at $p < 0.05$ (Tukey HSD post hoc test) among the concentrations, different uppercase of superscript letters in each column (100, 50, 25, 12.5, 6.25, 3.125) showed a significant difference at $p < 0.05$ (Tukey HSD post hoc test) among the cells.

Table 4: The percentage mortality (%) of Lymphocyte, HL-60, and K-562 treated with MPE in six different concentrations

Cells	Concentration (μ g/ml)					
	100	50	25	12.5	6.25	3.125
Lymphocyte	94.96 \pm 12.46 ^{cA}	55.91 \pm 8.82 ^{bA}	44.86 \pm 1.15 ^{abA}	42.2 \pm 1.46 ^{abA}	37.3 \pm 5.4 ^{abA}	16.17 \pm 3.71 ^{aA}
HL-60	106.7 \pm 1.85 ^{bcA}	100.97 \pm 0.67 ^{bb}	106.8 \pm 0.52 ^{bc}	109.1 \pm 1.7 ^{cc}	101.65 \pm 1.36 ^{bc}	89.7 \pm 1.11 ^{ac}
K-562	98.08 \pm 0.83 ^{dA}	95.06 \pm 2.95 ^{cdB}	93.14 \pm 4.21 ^{cdB}	86.24 \pm 0.76 ^{cB}	73.16 \pm 0.23 ^{bb}	46.23 \pm 0.65 ^{ab}

Data is presented as mean \pm standard deviation, different lowercase of superscript letters in each row (Lymphocyte, K-562, HL-60) showed a significant difference at $p < 0.05$ (Tukey HSD post hoc test) among the concentrations, different uppercase of superscript letters in each column (100, 50, 25, 12.5, 6.25, 3.125) showed a significant difference at $p < 0.05$ (Tukey HSD post hoc test) among the cells.




Fig. 1. Morphological appearance of HL-60 and K-562 cells with no treatment (1,2), treated with Imatinib (3-8), Isotretinoin (9-14), MPE (15-26), and α -mangostin (27-38). Scale bar: 200 μ m. Control (untreated) cells showed higher density than treated cells. The density of HL-60 and K-562 were low in higher MPE and α -mangostin concentrations.

3.2 Sensitivity and Selectivity of Each Compound toward HL-60, K-562, and Leukocyte

Further analysis of percentage mortality of the compounds using probit regression revealed that the IC_{50} of imatinib, isotretinoin, MPE, and α -mangostin were high to normal lymphocyte and the IC_{50} MPE and α -mangostin were comparably low to the leukemia cell lines (Figure 2). MPE


possessed the lowest IC_{50} (2.79 μ g/ml) toward K-562 whereas α -mangostin possessed the lowest IC_{50} (1.12 μ g/ml) to HL-60 (Table 5). The result of selectivity index showed that isotretinoin, a recent chemotherapy agent used to treat leukemia was less selective than MPE and α -mangostin toward HL-60, whilst imatinib was more selective toward K-562 than MPE and α -mangostin. Nevertheless, MPE and α -mangostin showed high selectivity toward both of HL-60 and K-562, with selectivity index was more than 3 (Table 5).

Table 5: The cytotoxicity and selectivity of MPE, α -mangostin, imatinib and isotretinoin against HL-60, K-562, and leukocyte

Compounds	Cell Line				
	HL-60		K-562		Lymphocyte
	IC ₅₀ (μ g/ml)	SI ^a	IC ₅₀ (μ g/ml)	SI ^a	IC ₅₀ (μ g/ml)
MPE ^b	1.16	19.83	2.79	8.27	23.08
α -mangostin	1.12	22.34	7.21	3.47	25.02
Isotretinoin	7.66	2.86	-	-	21.93
Imatinib	-	-	2.84	12.07	34.35

a. Selectivity Index, calculated based on the ratio of IC₅₀ lymphocyte and the respective leukemic cell line.

b. Mangosteen peel extract.

**Fig. 2.** IC₅₀ values of isotretinoin, imatinib, Mangosteen Peel Extract (MPE) and α -mangostin toward leukemia cell lines (HL-60 and K-562) and normal lymphocyte.

4. Discussion

The activities of xanthenes and its derivatives to inhibit certain molecular target in cancer progression are related to the tricyclic scaffold and its position⁸. As an anti-cancer, xanthenes arrest cell cycle, induce apoptosis, and differentiation but inhibit the tumor cell proliferation, adhesion, invasion, and metastasis¹⁶⁻¹⁸. Xanthenes were also reported to prevent initiation stage of cancer by inducing Quinine Reductase (QR) and inhibiting cytochrome P450 (CYP) activity^{9,19}.

In this study, imatinib, isotretinoin, MPE, and α -mangostin showed a cytotoxic effect on the HL-60, K-562, and lymphocyte cells in a concentration-dependent manner. Microscopic images showed the cell density gradually

decreased in the higher concentration of each treatment (Figure 1). This result was supported by the percentage of mortality data that showed the cell mortality increased in higher concentration of each treatment. Based on the Table 5, the IC₅₀ value of MPE and α -mangostin toward HL-60 and K-562 cell lines was lower than the IC₅₀ value toward lymphocyte, with the selectivity index higher than 3. These findings suggested that MPE and α -mangosteen were safe against the normal lymphocyte, and possessed a high selectivity and sensitivity toward HL-60 and K-562^{8,15}. These results were consistent with Matsumoto *et al.*²² study that showed the antiproliferative activity of xanthenes in mangosteen pericarp against human leukemia²⁰. The α -mangostin mediates mitochondrial apoptotic pathway in human promyelocytic leukemia (HL-60) by activating

the caspase-3 and caspase-9²¹. Together with β -mangostin, γ -mangostin, and methoxy- β -mangostin, these compounds arrest the cell cycle via expression of cyclin proteins in the human colon cancer cells (DLD-1)¹¹. Furthermore, other anti-tumor activities of mangostins in several cancers have also been reported, including the inhibition of TCF/ β -catenin transcription in colon cancer cells and inhibition of cell growth signaling pathways in chondrosarcoma²². These findings suggest that xanthenes work by various pathways to cancer cells.

In contrast, isotretinoin was either less selective or less sensitive toward the HL-60 cells compared with MPE and α -mangostin. Cancer has been known to be able to develop resistance towards chemotherapy. The resistance of leukemic cell lines to retinoic acid derivate might occur in molecular level by affecting several proteins functions and the mutations of the RAR α receptor in APL can block the initiation of differentiation by retinoic acid²³. On the contrary, Imatinib shows good sensitivity and selectivity toward K-562 human Chronic Myeloid Leukemia (CML) cells. CML is characterized by the presence of a Bcr-Abl fusion gene, which is caused by a reciprocal translocation of chromosomes 9 and 22²⁴. The cytotoxic activity of imatinib was supported by other study, demonstrated that imatinib was able to inhibit Bcr-Abl kinase activity led to inactivate of survival pathways and induce long-term activation of caspases that responsible for the degradation and inactivation of Bcr-Abl tyrosine kinase as well as apoptosis of the K562 cells²⁵.

5. Conclusion

Take together, our data suggest that MPE and α -mangostin possessed potent sensitivity and selectivity against leukemia. Both of them revealed higher selectivity and sensitivity than isotretinoin toward HL-60 cell line, while MPE also show high sensitivity and selectivity toward K-562 cell line, showing its great potential for pharmaceutical application. We suggest that MPE can be produced as a safe, efficient and low cost of an alternative remedy to fight leukemia. Therefore, the further study of mangosteen peel extract in molecular and *in vivo* study must be conducted.

6. Acknowledgements

We gratefully acknowledge the financial support from Research Grant of Hibah Bersaing 2016 by Ministry of Research, Technology and Higher Education of the

Republic of Indonesia and Biomedical and Biomolecular Research Centre Aretha Medika Utama, Bandung, Indonesia for research grant, research method and facilities support. We are also thankful to I Dewa Gde Sathya Deva from Biomedical and Biomolecular Research Centre Aretha Medika Utama, Bandung, Indonesia for his valuable assistance.

7. References

1. Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, et al. Cancer incident and mortality worldwide: Source, methods and major patterns in GLOBOCAN 2012. *Int J Cancer*. 2014; 136:349–86.
2. McDonald M, Hertz R, Lowenthal SW. *Pfizer facts: The burden of cancer in Asia USA*: Pfizer Medical Division; 2008.
3. Modak H, Kulkarni SS, Kadakol GS, Hiremath SV, Patil BR, Hallikeri U, et al. Prevalence and risk of leukemia in the multi-ethnic population of North Karnataka. *Asian Pac J Cancer Prev*. 2011; 12:671–5.
4. Belson M, Kingsley B, Holmes A. Risk factors for acute leukemia in children: A review. *Environ Health Persp*. 2007; 115(1):138–45.
5. Henkes M, van der Kuip H, Aulitzky WE. Therapeutic options for chronic myeloid leukemia: focus on imatinib (Glivec®, Gleevec™). *Ther Clin Risk Manag*. 2008; 4(1):163–87.
6. Niles RM. Recent advances in the use of Vitamin A (Retinoids) in the prevention and treatment of cancer. *Nutr*. 2000; 16:1084–90.
7. Gottesman MM, Fojo T, Bates ES. Multidrug resistance in cancer. *Nat Rev Cancer*. 2002; 2:48–58.
8. Shan T, Ma Q, Guo K, Liu J, Li W, Wang F, et al. Xanthenes from mangosteen extracts as natural chemopreventive agents: potential anticancer drugs. *Curr Mol Med*. 2011; 11(8):666–77.
9. Foti RS, Pearson JT, Rock DA, Wahlstrom JL, Wienkers LC. In vitro inhibition of multiple cytochrome P40 isoforms by xanthone derivatives from mangosteen extract. *Drug Metab Dispos*. 2009; 37(9):1848–55.
10. Yang J, Liu RH, Halim L. Antioxidant and antiproliferative activities of common edible nut seeds. *Lwt-Food Sci Technol*. 2009; 42(1):1–8.
11. Aisha AFA, Abu-Salah KM, Ismail Z, Abdul-Majid AMS. In vitro and in vivo anti-colon cancer effects of Garcina

- mangostana xanthenes extract. BMC Compl Alternative Med. 2012; 12(104):1–10.
12. Darsono L, Hidayat M, Maesaroh M, Fauziah N, Widowati W. Ex vivo study of *Garcinia mangostana* L. (Mangosteen) peel extract and xanthenes as anti-adipogenesis in HepG2 cell model. Int J Med Res Health Sci. 2015; 4(3):566–71.
 13. Widowati W, Darsono L, Suherman J, Yelliantty Y, Maesaroh M. High Performance Liquid Chromatography (HPLC) analysis, antioxidant, antiaggregation of mangosteen peel extract (*Garcinia mangostana* L.). Int J Biosci Biochem Bioinfo. 2014; 4(6):458–66.
 14. Boyum A. Isolation of mononuclear cells and granulocytes from human blood. Isolation of monuclear cells by one centrifugation, and of granulocytes by combining centrifugation and sedimentation at 1 g. Scand J Clin Lab Invest Suppl 1968; 97:77–89.
 15. Mahavorasirikul W, Viyanant V, Chaijaroenkul W, Itharat A, Na-Bangchang K. Cytotoxic activity of Thai medicinal plants against human cholangiocarcinoma, laryngeal and hepatocarcinoma cells in vitro. BMC Complement Altern Med. 2010; 10(55):1–8.
 16. Akao Y, Nakagawa Y, Iinuma M, Nozawa Y. Anti-cancer effects of xanthenes from pericarps of mangosteen. Int J Mol Sci. 2008; 9:355–70.
 17. Pedraza-Chaverri J, Cardenas-Rodriguez N, Orozco-Ibarra M, Perez-Rojas JM. Medicinal properties of mangosteen (*Garcinia mangostana*). Food Chem Toxicol. 2008; 46:3227–39.
 18. Hung SH, Shen KH, Wu CH, Liu CL, Shih YW. Alpha-mangostin suppresses PC-3 human prostate carcinoma cell metastasis by inhibiting matrix metalloproteinase-2/9 and urokinase-plasminogen expression through the JNK signaling pathway. J Agric Food Chem. 2009; 57(4):1291–8.
 19. Chin YW, Jung HA, Chai H, Keller WJ, Kinghorn AD. Xanthenes with quinone reductase-inducing activity from the fruits of *Garcinia mangostana* (Mangosteen). Phytochemistry. 2008; 69(3):754–8.
 20. Matsumoto K, Akao Y, Kobayashi E, Ohguchi K, Ito T, Tanaka T, et al. Induction of apoptosis by xanthenes from mangosteen in human leukemia cell lines. J Nat Prod. 2003; 66:1124–7.
 21. Matsumoto K, Akao Y, Yi H, Ohguchi K, Ito T, Tanaka T, et al. Preferential target is mitochondria in alpha-mangostin-induced apoptosis in human leukemia HL60 cells. Bioorg Med Chem. 2004; 12(22):5799–806.
 22. Krajarng A, Nakamura Y, Suksamram S, Watanapokasin R. Preferential target is mitochondria in alpha-mangostin-induced apoptosis in human leukemia HL60 cells. J Agric Food Chem. 2011; 59:5746–54.
 23. Niles RM. Recent advances in the use of vitamin A (Retinoids) in the prevention and treatment of cancer. Nutr. 2000; 16:1084–90.
 24. Deininger M, Buchdunger E, Druker BJ. The development of imatinib as a therapeutic agent for chronic myeloid leukemia. Blood. 2005; 105(7):2640–53.
 25. Jacquelin A, Herrant M, Legros L, Belhacene N, Luciano F, Pages G, et al. Imatinib induces mitochondria-dependent apoptosis of the Bcr-Abl-positive K562 cell line and its differentiation toward the erythroid lineage. FASEB J. 2003; 17(14):2160–2.

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau PEER REVIEW

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Cytotoxic Activity of Mangosteen (*Garcinia mangostana* L.) Peel Extract and α -mangostin toward Leukemia Cell Lines (HL-60 and K-562)

Jumlah Penulis : 6 Orang

Nama-nama Penulis : Arina Novilla, Dedi S. Djambhuri, Nurul Fauziah, Maesaroh Maesaroh, Balqis Balqis, **Wahyu Widowati**

Status Penulis : ~~Penulis Pertama~~ / Penulis ke 6 / Penulis Korespondensi **)

Identitas Jurnal Ilmiah : a. Nama jurnal : Journal of Natural Remedies
b. Nomor ISSN : 2320-3358
c. Vol., No., Bulan, Tahun : Vol 16; No 2, April 2016
d. Penerbit : Informatics Publishing Limited
e. DOI Artikel (jika ada) : 10.18311/jnr/2016/842
f. Alamat Web Jurnal : www.informaticsjournals.com/index.php
g. Terindeks di : Scopus Q4, SJR 0.154

Kategori Publikasi Jurnal Ilmiah: Jurnal Ilmiah Internasional / Internasional Bereputasi **)
(beri tanda \surd yang dipilih)

Jurnal Ilmiah Nasional Terakreditasi

Jurnal Ilmiah Nasional / Nasional terindeks ***)

HASIL PENILAIAN (Peer Review) :


No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi <input checked="" type="checkbox"/>	Nasional Terakreditasi <input type="checkbox"/>	Nasional ***) <input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,6
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			11,5
c.	Kecukupan dan kemitakhiran data/informasi dan metodologi (30%)	12			11,6
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			11,4
	Total	40			38,1

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur. Penulisan memenuhi standar karya ilmiah akademis, metoda yang digunakan sesuai dan tepat digunakan pada penelitian ini
- b. Ruang lingkup & kedalaman pembahasan. Penelitian tentang potensi ekstrak kulit manggis dan senyawa A-mangosteen menghambat proliferasi sel kanker leukemia
- c. Kecukupan & kemitakhiran data serta metodologi. lde dasar baik membahas tingkat selektivitas ekstrak kulit manggis, senyawa A-mangosteen sebagai antikanker leukemia
- d. Kelengkapan unsur dan kualitas penerbit. Jurnal JNR terindeks Scopus Q4 SJR 0,154 Penerbit Informatics Publishing Limited
- e. Indikasi plagiasi. tidak terdapat indikasi plagiarism atau self plagiarism

f. Kesesuaian bidang ilmu *Paper bidang biokimia, biomedik sesuai dengan bidang Ilmu sejenis*

REVIEWER 1



(Prof. Dr. Chrismis Novalinda Ginting, M.Kes)
NIK: 0115127801
UNIVERSITAS PRIMA INDONESIA

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau *PEER REVIEW*

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Cytotoxic Activity of Mangosteen (*Garcinia mangostana* L.) Peel Extract and α -mangostin toward Leukemia Cell Lines (HL-60 and K-562)

Jumlah Penulis : 6 Orang

Nama-nama Penulis : Arina Novilla, Dedi S. Djamhuri, Nurul Fauziah, Maesaroh Maesaroh, Balqis Balqis, **Wahyu Widowati**

Status Penulis : ~~Penulis Pertama~~ / Penulis ke 6 / Penulis Korespondensi **)

Identitas Jurnal Ilmiah : a. Nama jurnal : Journal of Natural Remedies
b. Nomor ISSN : 2320-3358
c. Vol., No., Bulan, Tahun : Vol 16; No 2, April 2016
d. Penerbit : Informatics Publishing Limited
e. DOI Artikel (jika ada) : 10.18311/jnr/2016/842
f. Alamat Web Jurnal : www.informaticsjournals.com/index.php
g. Terindeks di : Scopus Q4, SJR 0.154

Kategori Publikasi Jurnal Ilmiah : Jurnal Ilmiah Internasional / Internasional Bereputasi **)
(beri tanda \surd yang dipilih)

Jurnal Ilmiah Nasional Terakreditasi

Jurnal Ilmiah Nasional / Nasional terindeks ***)

HASIL PENILAIAN (*Peer Review*) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi	Nasional Terakreditasi	Nasional ***)	
		<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,5
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			4,5
c.	Kecukupan dan kemutakhiran data/ informasi dan metodologi (30%)	12			11,7
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			11,6
	Total	100%			38,3

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur
Jurnal ini sudah cukup lengkap dan adanya kesesuaian antara unsur dan isinya
- b. Ruang lingkup & kedalaman pembahasan
Setiap ruang lingkup dan bahasan sudah meneliti keululaman pembahasan
- c. Kecukupan & kemutakhiran data serta metodologi
Secara umum metodologi sudah memadai dan lengkap kemutakhiran data juga sudah terpenuhi
- d. Kelengkapan unsur dan kualitas penerbit
Kualitas penerbit sudah memadai dan memenuhi kaidah dan kualitas jurnal, Artikel ini termasuk ke dalam kategori bereputasi yang baik.
- e. Indikasi plagiasi

Belum terlihat adanya indikasi plagiat.

f. Kesesuaian bidang ilmu

Artikel ini sudah sesuai dengan bidang ilmu yang diteliti oleh penulis.

REVIEWER 2



(Prof. Dr. Ermi Girsang, M. Kes)

NIK : 0117057501

UNIVERSITAS PRIMA INDONESIA

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau *PEER REVIEW*

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Cytotoxic Activity of Mangosteen (*Garcinia mangostana* L.) Peel Extract and α -mangostin toward Leukemia Cell Lines (HL-60 and K-562)

Jumlah Penulis : 6 Orang

Nama-nama Penulis : Arina Novilla, Dedi S. Djamhuri, Nurul Fauziah, Maesaroh Maesaroh, Balqis Balqis, **Wahyu Widowati**

Status Penulis : ~~Penulis Pertama~~ / Penulis ke 6 / Penulis Korespondensi **)

Identitas Jurnal Ilmiah : a. Nama jurnal : Journal of Natural Remedies
 b. Nomor ISSN : 2320-3358
 c. Vol., No., Bulan, Tahun : Vol 16; No 2, April 2016
 d. Penerbit : Informatics Publishing Limited
 e. DOI Artikel (jika ada) : 10.18311/jnr/2016/842
 f. Alamat Web Jurnal : www.informaticsjournals.com/index.php
 g. Terindeks di : Scopus Q4, SJR 0.154

Kategori Publikasi Jurnal Ilmiah: Jurnal Ilmiah Internasional / Internasional Bereputasi **)
 (beri tanda \checkmark yang dipilih)

Jurnal Ilmiah Nasional Terakreditasi

Jurnal Ilmiah Nasional / Nasional terindeks ***)

HASIL PENILAIAN (*Peer Review*) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi	Nasional Terakreditasi	Nasional ***)	
		<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,55
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			11,5
c.	Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12			11,65
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			11,5
	Total	100%	40		38,2

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur. *Penulisan memenuhi standard karya ilmiah akademis, metoda yang digunakan sesuai dan tepat digunakan pada penelitian ini*
Jurnal ini sudah cukup lengkap dan adanya kesesuaian antara unsur dan isinya
- b. Ruang lingkup & kedalaman pembahasan *Penelitian tentang potensi ekstrak kulit manggis dan senyawa A-mangosteen menghambat proliferasi sel kanker leukemia*

Seluruh ruang lingkup dan bahasan sudah memiliki kedalaman pembahasan.

c. Kecukupan & kemutakhiran data serta metodologi. Ide dasar baik membahas tingkat selektivitas ekstrak kulit manggis, senyawa A- mangosteen sebagai antikanker leukemia.

Secara umum metodologi sudah memadai dan lengkap kemutakhiran data juga sudah terpenuhi.

d. Kelengkapan unsur dan kualitas penerbit

Jurnal JNA terindeks Scopus Q1, STAR 0,1514. Penerbit Informatics Publishing Limited.

Kualitas penerbit sudah memadai dan memenuhi kaidah dan kualitas jurnal. Artikel ini termasuk ke dalam kategori bereputasi yang baik.

e. Indikasi plagiasi tidak terdapat indikasi plagiarism atau self plagiarism.

Belum terlihat adanya indikasi plagiasi.

f. Kesesuaian bidang ilmu


Paper bidang biokimia, biomedik sesuai dengan bidang Ilmu sjenis.

Artikel ini sudah sesuai dengan bidang Ilmu yang diteliti oleh penulis.

Medan,
Reviewer 2


(Prof. Dr. Ermi Girsang, M.Kes)
NIK : 0117057501
UNIVERSITAS PRIMA INDONESIA

Medan,
Reviewer 1


(Prof. Dr. Chrismis Novalinda Ginting, M.Kes)
NIK : 0115127801
UNIVERSITAS PRIMA INDONESIA