

DAFTAR PUSTAKA

1. Indriyani S, Retayasa WI, Surjono A, Suryantoro P. 2009. Percentage birth weight loss and hyperbilirubinemia during the first week of life in term newborns. *Paediatrica Indonesiana*, 3 (49): 149-154.
2. Maisels MJ, Gifford K, Antle CE, Leib GR. 1988. Jaundice in the healthy newborn infant: A new approach to an old problem. *PEDIATRICS*, 4 (81): 505-511.
3. Macdonald PD, Ross SRM, Grant L, Young D. 2003. Neonatal weight loss in breast and formula fed infants. *Arch Dis Child Fetal Neonatal*, (88): 472-476.
4. Bertini G, Dani C, Tronchin M, Rubaltelli FF. 2001. Is breastfeeding really favoring early neonatal jaundice? *PEDIATRICS*, 3 (107): 1-5.
5. Mulder PJ, Johnson TS, Baker LC. 2010. Excessive weight loss in breastfed infants during the postpartum hospitalization. *Journal of Obstetric Gynecologic and Neonatal Nursing*, 1 (39): 15-26.
6. Davanzo R, Cannioto Z, Ronfani L, Monasta L, Demarini S. 2012. Breastfeeding and neonatal weight loss in healthy term infants. *Journal of Human Lactation*, 20 (10): 1-9.
7. Chang RJ, Chou HC, Chang YH, et al. 2012. Weight loss percentage prediction of subsequent neonatal hyperbilirubinemia in exclusively breastfed neonatus. *Pediatrics and Neonatology*, (53): 41-44.

8. Yang WC, Zhao LL, Li YC, *et al.* 2013. Bodyweight loss in predicting neonatal hyperbilirubinemia 72 hours after birth in term newborn infants. *BMC Pediatrics*, (13): 1-7.
9. Sarici SU, Serdar MA, Korkmaz A, *et al.* 2004. Incidence, course, and prediction of hyperbilirubinemia in near-term and term newborns. *PEDIATRICS*, 4 (113): 775-780.
10. Alatas H et al. 1985. *Ilmu kesehatan anak 3. 4th ed* : INFOMEDIKA. p 1050.
11. Weiss JN, Woodend AK, Peterson WE, Gibb W, Groll DL. 2011. An observational study of associations among maternal fluids during parturition, neonatal output, and breastfed newborn weight loss. *International Breastfeeding Journal*, (6): 1-10.
12. Fajrina A. 2012. Hubungan pertambahan berat badan selama hamil dan faktor lain dengan berat badan lahir di Rumah Bersalin Lestari Ciampea Bogor tahun 2010-2011 : 1-77.
13. Korones SB. 1976. *High-risk newborn infants*. 2nd ed : The C.V Mosby Company. p 65, 193-203.
14. Polin RA. 2011. *Fetal and Neonatal Physiology*. 4th ed : Elsevier Saunders. p 648-661.
15. Conita I. 2013. Hubungan penurunan berat badan dengan kadar bilirubin neonatus pada hari ketiga pasca lahir : 1-36.

16. Guyton AC, Hall JE. 2006. *Textbook of medical physiology*. 11th ed : Elsevier Saunders. p 1038-1041.
17. Kosim MS, Yunanto A, Dewi R, Sarosa GI, Usman A. 2014. *Buku Ajar Neonatologi*. 1st ed : Badan Penerbit IDAI. p 147-157, 161, 167.
18. Murray RK, Granner DK, Mayes PA, Rodwell VW. 2003. *Harper's Illustrated Biochemistry*. 26th ed : McGraw-Hill Companies. p 270-277.
19. Alatas H et al. 1985. *Ilmu kesehatan anak 2*. 4th ed : INFOMEDIKA. p 519-522.
20. R. Arvind. 2006. *Applied neonatology*. 1st ed : Jaypee Brothers. p 190-196, 202-205.
21. Arant BS. 1981. Adaptation of the infant to an external milieu. Gruslan AB (Ed). *Pediatric nephrology: Proceedings of the fifth international pediatric nephrology symposium*. 3rd ed. The Hague: Martinus Hijhoff Publishers. p 265.
22. Stevenson DK, Maisels MJ, Watchko JF. 2012. *Care of the Jaundiced Neonate* : The McGraw-Hill Company. p 66, 71.