

## DAFTAR PUSTAKA

1. Singer M, Deutschman CS, Seymour C, Shankar-Hari M, Annane D, Bauer M, et al. The third international consensus definitions for sepsis and septic shock (sepsis-3). *JAMA - J Am Med Association*;2016;315(8):801–10.
2. Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock. *Intensive Care Medicine*;2017;43(3):304–77.
3. World Health Organization. Sepsis. Media centre Seventieth World Health Assembly update;2017.p.3–5.
4. World Health Organization. C-reactive protein concentrations as a marker of inflammation or infection for interpreting biomarkers of micronutrient status. *Vitamin as Mineral Nutrition Information System*;2014.p.1–4.
5. Nierhaus A, Klatte S, Linssen J, Eismann NM, Wichmann D. Revisiting the white blood cell count: immature granulocytes count as a diagnostic marker to discriminate between SIRS and sepsis - a prospective, observational study. *BMC Immunology*;2013;13(1):1–20.
6. Afriyanti E. Peranan C-Reaktif Protein (CRP) Sebagai Parameter Diagnosis Sepsis Neonatorum. Fakultas Kedokteran Unand; 2010.
7. Kristiorini H. Perbandingan antara C-Reaktif Protein dan Prokalsitonin pada Anak dengan Disfungsi Multi Organ Akibat Sepsis. *digital library*.Surakarta;Universitas Sebelas Maret; 2014.
8. Antonelli M, DeBacker D, Dorman T, Kleinpell R, Levy MM, Rhodes A. Surviving Sepsis Campaign Responds to Sepsis-3;2016.p. 4–5. Available from: [www.sccm.org/sepsisredefined](http://www.sccm.org/sepsisredefined)
9. World Health Organisation. Improving the prevention, diagnosis and clinical management of sepsis. WHO;2018.p.1–2. Available from: <http://www.who.int/sepsis/en/>
10. Ramachandran G. Gram-positive and gram-negative bacterial toxins in sepsis: A brief review. *Virulence*; 2014;5(1):196–201.
11. Starr ME, Saito H. Sepsis in Old Age: Review of Human and Animal Studies;*Aging Disease*;2014;5(2):126–37.

12. Keputusan Menteri Kesehatan Republik Indonesia Nomor Hk.01.07/Menkes/342/2017 Tentang Pedoman Nasional Pelayanan Kedokteran Tata Laksana Sepsis. Jakarta:Kemenkes RI;2017.p.1–82.
13. Dinas Kesehatan Jawa Barat. Profil Kesehatan Indonesia. Bandung;Dinkes JABAR;2012.p.1–138.
14. Juniatiningsih A, Aminullah A, Firmansyah A. Profil Mikroorganisme Penyebab Sepsis Neonatorum di Departemen Ilmu Kesehatan Anak Rumah Sakit Cipto Mangunkusumo. *Sari Pediatri*. Jakarta; 2016;10(1):60.
15. Centre for Disease Control and Prevention (CDC); Basic Information of Sepsis. Available from: <https://www.cdc.gov/sepsis/basic/index.html>
16. Cunha ABA, Editor C, Stuart M. Bacterial Sepsis. *Medscape Drugs Disease*; 2015.p.1–5.
17. Angus DC, van der Poll T. Severe Sepsis and Septic Shock. *New England Journal of Medicine*;2013;369(9):840–51.
18. Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, Cook D, et al. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Intensive Care Med*; 2003;29(4):530–8.
19. Marshall JC. Predisposition to Sepsis. *Anaesthesia, Pain, Intensive Care Emergency; APICE*;2008.p.241–54.
20. Wiersinga WJ, Seymour CW. Handbook of Sepsis. *Springer International Publishing AG*;2018.p.31–43.
21. Minasyan H. Sepsis: Pathophysiology and Treatment. *Journal Intensive Critical Care*;2017;03(02):10–2.
22. Kalil A. Septic Shock Clinical Presentation. *Medscape Drugs Disease*;2015.p.1–5.
23. Salazar J, Martínez MS, Chávez-castillo M, Núñez V, Añez R, Torres Y, et al. C-Reactive Protein : An In-Depth Look into Structure , Function , and Regulation. *Hindawi Publishing Corporation*;2014.p.2–3
24. Ansar W, Ghosh S. *Biology of C Reactive Protein in Health and Disease*;2016.p.160.
25. Andareswari N, Maulana S. Penetapan Kadar CRP secara Kualitatif. Kementerian Riset Teknologi Dan Pendidikan Tinggi Universitas Jenderal Soedirman Fakultas Biologi. Purwokerto ; 2015.p.7–8

26. Fiscbach F, Dunning M, *a manual of laboratory and diagnostic tests ninth edition*. Wolters Kluwer;2015.p.72–5
27. Zhang H, Wang X, Zhang Q, Xia Y, Liu D. Comparison of procalcitonin and high-sensitivity C-reactive protein for the diagnosis of sepsis and septic shock in the oldest old patients. *BMC Geriatrics*;2017;17(1):1–6.
28. Hangandi F. Sensitivity of serum C-reactive protein ( CRP ) levels compared to white blood cell count ( WBC ) as a predictor of surgical site infection in patients undergoing major surgery. *J Med Sci*;2013;45:159–66.
29. Mare TA, Treacher DF, Shankar-Hari M, Beale R, Lewis SM, Chambers DJ, et al. The diagnostic and prognostic significance of monitoring blood levels of immature neutrophils in patients with systemic inflammation. *Critical Care*;2015;19(1):1–11.
30. Ruotsalo K, Tant M. Complete Blood Count. *VCA Hospital*;2008; Available from: <https://vcahospitals.com/know-your-pet/complete-blood-count>
31. Lipiński M, Rydzewska G. Immature granulocytes predict severe acute pancreatitis independently of systemic inflammatory response syndrome. *Przegląd Gastroenterologiczny*;2017;12(2):140–4.
32. Senthilnayagam B, Kumar T, Sukumaran J, Jeya M, Rao K. R. Automated measurement of immature granulocytes: Performance characteristics and utility in routine clinical practice. *Pathology Research International*;2012.p.1-5
33. Sysmex Europe GmbH. The Immature Granulocyte Count – The first to know about inflammation;2011. p. 1–4. Available from: [www.sysmex-europe.com](http://www.sysmex-europe.com)
34. Sysmex Educational Enhancement and Development. *SEED Haematology*; 2012. p. 1–7. Available from: [www.sysmex-europe.com](http://www.sysmex-europe.com)
35. Mathews EK, Griffin RL, Mortellaro V, Beierle EA, Harmon CM, Chen MK, et al. Utility of immature granulocyte percentage in pediatric appendicitis. *Journal of Surgical Research*;2014;190(1):230–4.
36. Kementrian Kesehatan Republik Indonesia. Pedoman Interpretasi Data Klinik.Jakarta;2011.

37. Honda T, Uehara T, Matsumoto G, Arai S, Sugano M. Neutrophil left shift and white blood cell count as markers of bacterial infection. *Clinica Chimica Acta*;2016;457:46–53.
38. Girard TD, Opal SM, Ely EW. Insights into Severe Sepsis in Older Patients: From Epidemiology to Evidence-Based Management. *Clinical Infection Disease*; 2005;40(5):719–27.
39. Wester AL, Dunlop O, Melby KK, Dahle UR, Wyller TB. Age-related differences in symptoms, diagnosis and prognosis of bacteremia. *BMC Infection Disease*;2013;13(1):346.
40. Melamed A, Sorvillo FJ. The burden of sepsis-associated mortality in the United States from 1999 to 2005: An analysis of multiple-cause-of-death data. *Critical Care*;2009;13(1):1–8.
41. Angele MK, Frantz MC, Chaudry IH. Gender and sex hormones influence the response to trauma and sepsis: potential therapeutic approaches. *Clinics* ;2006;61(5):479–88.
42. Buoro S, Mecca T, Azzarà G, Apassiti Esposito S, Seghezzi M, Vavassori M, et al. Extended leukocyte differential count and C-reactive protein in septic patients with liver impairment: diagnostic approach to evaluate sepsis in intensive care unit. *Annals of Translational Medicine* ;2015;3(17):244.
43. Pavare J, Grope I, Gardovska D. Measurement of immature granulocytes (IG) percentage to assess severe bacterial infection in Latvian children: a secondary data analysis. *Department of Pediatrics, Riga Stradins University, Dzirciema street 16, Riga, Latvia, Riga*;2018.