

## DAFTAR PUSTAKA

1. World Health Organization (WHO). Global Strategy on Diet [Internet]. 2018. Available from: <http://www.who.int/dietphysicalactivity/pa/en/>
2. Mills O. Exercise\_Guide. Maryland: Medifast Inc; 2007.
3. Lira VA, Okutsu M, Zhang M, Greene NP, Laker RC, Breen DS, et al. Autophagy is required for exercise training-induced skeletal muscle adaptation and improvement of physical performance. *FASEB J.* 2013;27(10):4184–93.
4. Boya P, Codogno P. Autophagy : Mechanisms and Function. 2013;28040.
5. Mejías-Peña Y, Rodriguez-Miguel P, Fernandez-Gonzalo R, Martínez-Flórez S, Almar M, de Paz JA, et al. Effects of aerobic training on markers of autophagy in the elderly. *Age (Omaha)*. 2016;38(2).
6. Martin-Rincon M, Morales-Alamo D, Calbet JAL. Exercise-mediated modulation of autophagy in skeletal muscle. *Scand J Med Sci Sport*. 2018;28(3):772–81.
7. Vainshtein A, Hood DA. The regulation of autophagy during exercise in skeletal muscle. *J Appl Physiol* [Internet]. 2016;120(6):664–73. Available from: <http://jap.physiology.org/lookup/doi/10.1152/japplphysiol.00550.2015>
8. Rodgers AB. Exercise & Physical Activity. Your Everyday Guid from Natl Inst Aging NIH Exerc Phys Act. 2013;
9. Soya H MA, Deocaris CC, Ohiwa N Chang H NT, Al E. Threshold-like pattern of neuronal activation in the hypothalamus during treadmill running : Establishment of a minimum running stress ( MRS ) rat model. 2017;
10. Lubis L S. Response of Long-Term Memory to Molecular Changes of BDNF in Hippocampus in Various Intensities of Physical Activity. 2016;
11. Lesmana R, Iwasaki T, Iizuka Y, Amano I SN. The change in thyroid hormone signaling by altered training intensity in male rat skeletal muscle. 2016;
12. Bernardo A, Petriz, Clarissa P.C. Gomes, Jeeser A, Almeida, Getulio P. De Oliveira Jr, Filipe M. Ribeiro, Rinaldo W. Pereira OLF. The Effect of Acute and Chronic Exercise on Skeletal Muscle. 2016;
13. Richard L D, Vogl AW, Mitchell AWM. Gray Dasar-Dasar Anatomi. Kalanjati VP, editor. Singapura: Elsevier Inc.; 2012.
14. Eroschenko VP. Atlas Histologi diFiore dengan Korelasi Fungsional. edisi 11. Dharmawan, Didiek. Yesdelita N, editor. Amerika Serikat: Lippincott

- Williams & Wilkins; 2008. 117 p.
15. Hall JE, Arthur C G. Guyton Dan Hall Buku Ajar Fisiologi Kedokteran. edisi kedu. Singapura: Elsevier Inc.; 2014.
  16. Bhutia SK, Praharaj PP, Bhol CS, Panigrahi DP, Mahapatra KK, Patra S, et al. Monitoring and measuring mammalian autophagy. *Methods Mol Biol*. 2019;1854:209–22.
  17. Mizushima N. Physiological functions of autophagy. *Curr Top Microbiol Immunol*. 2009;335(1):71–84.
  18. Mizushima N, Yoshimori T, Levine B. Methods in Mammalian Autophagy Research. *Cell*. 2010;140(3):313–26.
  19. Huang CC, Wang T, Tung YT, Lin WT. Effect of exercise training on skeletal muscle SIRT1 and PGC-1 $\alpha$  expression levels in rats of different age. *Int J Med Sci*. 2016;13(4):260–70.
  20. Britannica TE of E. Polymerase Chain Reaction [Internet]. 2018. Available from: <https://www.britannica.com/science/polymerase-chain-reaction>
  21. Bachman J. Reverse-transcription PCR (RT-PCR) [Internet]. 1st ed. Vol. 530, Methods in Enzymology. Elsevier Inc.; 2013. 67-74 p. Available from: <http://dx.doi.org/10.1016/B978-0-12-420037-1.00002-6>
  22. Academy K. Polymerase Chain Reaction (PCR). 2018.
  23. American Psychological Association. APS Resource Book for the Design of Animal Exercise Protocols. 2006.
  24. Wang Y, Zhang Y, Qian C, Cai M, Li Y, Li Z, et al. GSK3 $\beta$ /β-catenin signaling is correlated with the differentiation of glioma cells induced by Wogonin. *Toxicol Lett* [Internet]. 2013;222(2):212–23. Available from: <http://dx.doi.org/10.1016/j.toxlet.2013.07.013>
  25. Qi L. An Indian Journal. Biotechnology. 2014;4(3):3–6.
  26. Schwalm C, Jamart C, Benoit N, Naslain D, Prémont C, Prevet J, et al. Activation of autophagy in human skeletal muscle is dependent on exercise intensity and AMPK activation. *FASEB J*. 2015;29(8):3515–26.
  27. Li FH, Li T, Ai JY, Sun L, Min Z, Duan R, et al. Beneficial autophagic activities, mitochondrial function, and metabolic phenotype adaptations promoted by high-intensity interval training in a rat model. *Front Physiol*. 2018;9(MAY).
  28. Pauly M, Assense A, Rondon A, Thomas A, Dubouchaud H, Freyssenet D, et

al. High intensity aerobic exercise training improves chronic intermittent hypoxia-induced insulin resistance without basal autophagy modulation. *Sci Rep* [Internet]. 2017;7(January):1–12. Available from: <http://dx.doi.org/10.1038/srep43663>

