




JOURNAL OF NATURAL REMEDIES

Dedicated to Medicinal Plant Research

ISSN : 0972 - 5547





Editorial Board

Editor-In-Chief

Dr. M. K. Mohan Maruga Raja

Senior Scientist, TanBio R&D Solution, Penyar Technology Business Incubator, Periyar Maniammal University, Thanjavur, Tamil Nadu

Co-Editors

Dr. D. Prashanth

Natural Remedies, Bangalore

Balu Ranganathan

Palms Connect LLC, Sandy, Utah, USA

Dr. M. K. Kathiravan

Department of Pharmaceutical Chemistry, SRM University, Kattankulathur, Tamil Nadu

Dr. Jollus Gimbut

Deputy Director of Centre of Excellence for Advanced Research in Fluid Flow (CARIFF), Assoc. Professor, Faculty of Chemical and Natural Resources Engineering, The Gimbut Group, Universiti Malaysia Pahang, Gambang, Pahang, Malaysia

Editorial Board Members

Dr. Acharya Sanjeev R.

Institute of Pharmacy, Nirme University, Ahmedabad

Dr. Akula Chinapolaiah

ICAR Indian Council of Agricultural Research, Anand Gujarat

Dr. Anant Bhandarkar

Goa College of Pharmacy, Govt. of Goa

Dr. Anil Kumar Uttamchand Tatiya

R.C.Patel College of Pharmacy Shirgur

Dr. Arun B. Joshi

Goa College of Pharmacy, Panaji, Goa

Dr. Chandrashekar K. S.

Manipal College of Pharmaceutical Sciences, Manipal University, Manipal

Dr. Dayanand M. Kamur

SCCES S.Indira College of Pharmacy, Pune

Dr. Deepak Kumar Semwal

Uttarakhand Ayurved University, Dehradun

Dr. Kuntal Das

Krupanidhi College of Pharmacy, Bangalore

Dr. Laxminarayana Baiyy

Kasturba Medical College, Manipal

Dr. Madhavan V.

M.S.Ramaiah College of Pharmacy, Bangalore

Dr. Manik Ghosh

Birla Institute of Technology, Mesra, Ranchi

Dr. Manjunath Shetty

Manipal College of Pharmaceutical Sciences, Manipal University, Manipal

Dr. Mohd. Yaseen

Central Institute of Medicinal and Aromatic Plants(CIMAP), Lucknow

Dr. Mruthunjaya K.

JSS College of Pharmacy, Mysore

Dr. Nagarajan A.

PSG College of Pharmacy, Coimbatore

Dr. Peter Achunike Akah

University of Nigeria, Nsukka, Nigeria

Dr. Rajendra Gyawali

Department of Pharmacy, Kathmandu

Dr. Sathianarayanan S.

Amrita School of Pharmacy Amrita Vishwavidyapeetram University, Kochi

Dr. Shailendra S. Gurav

Goa College of Pharmacy, Panaji, Goa

Dr. Sinha B. N.

Birla Institute of Technology, Mesra, Ranchi

Dr. Subhash C Mandal

Jadavpur University, Kolkata

Dr. Vasantha Kumar

ICAR, Indian Institute of Horticultural Research, Bengaluru

Dr. Vidhu Aeri

Jamia Hamdard Deemed University, New Delhi

Dr. Vikram Chauhan

Planet Ayurveda Clinic and Krishna Herbal Company, Mohali, India

Dr. Vimal Kumar

Institute of Pharmacy, Nirme University, Ahmedabad

Dr. Padmaa M Paarakh

Pharmacognosical, Phytochemical and Pharmacological studies, Banaras Hindu University, Varanasi

Prof.(Dr.) R. S. Tomar

Director, Amity Institute of Biotechnology, Madhya Pradesh

Manish K. R. Singh

Department of Poultry Science, Duvasu, Mathura, Uttar Pradesh

Dr. N. Kasthuri Bai

Interdisciplinary Institute of Indian System of Medicine, Tamil Nadu, India

Dr. Mani Ganesh

Hansoo University, Chungcheongnam-do, South Korea

Editorial Manager

Kavitha Kumar

Informatics Publishing Ltd.

Journal Coordinator

Priyadarshini C

Informatics Publishing Ltd.

Quick Links

- [Abstracting and Indexing](#)
- [Instructions to Authors](#)
- [Pre-submission checklist](#)
- [Publication Fees](#)
- [Journal Policies and Ethics](#)
- [Submission Process](#)
- [Information for Reviewers](#)
- [Email Alerts](#)
- [CrossMark Policy](#)

Archives

[Expand All]

[2019](#)

[2018](#)

[2017](#)

[2016](#)

[2015](#)

[2014](#)

[2013](#)

[2012](#)

[More..](#)


 All Search

- Home
- Current
- Archives
- Authors
- Institutions
- Editorial Board
- Submit Article
- Online First

Volume 16, Issue 3, July 2016

Table of Contents

Review Article

Paper Flower, *Bougainvillea spectabilis*: Update Properties of Traditional Medicinal Plant

Anisa Ghogar, Kanitta Jiraungkoorskul, Wanee Jiraungkoorskul

DOI: 10.18311/jnr/2016/5703, Pagination: 82-87



Abstract Views: 889 | Pdf Views: 3183

Articles

Antioxidant and Anti Aging Assays of *Oryza sativa* Extracts, Vanillin and Coumaric Acid

Wahyu Widowati, Nurul Fauziah, Heddy Herdiman, Merry Afni, Ervi Afifah, Hanna Sari W. Kusuma, Hayatun Nufus, Seila Arumwardana, Dwi Davidson Rihibiha

DOI: 10.18311/jnr/2016/7220, Pagination: 88-99



Abstract Views: 1421 | Pdf Views: 1823

Hepatoprotective and Antioxidant Activity of Ethanol Extracts of *Dipterocarpus turbinatus* (Dipterocarpaceae) from Tripura

Diptanu Biswas, T. Shivraj Gouda, Narayanswamy Lachmanan Gowrishankar

DOI: 10.18311/jnr/2016/7525, Pagination: 100-107



Abstract Views: 547 | Pdf Views: 401

A Doubled-Blind, Crossover-RCT in T2DM for Evaluating Hypoglycemic Effect of *P. indicus*, *M. charantia*, *P. vulgaris* and *A. paniculata* in Central Java

Mahalul Azam, Charles Ong Saerang, Sri Ratna Rahayu, Fitri Indrawati, Irwan Budiono, Arulita Ika Fibriana, Muhammad Azinar, Nur Anna C. Sa'dyah, Erwin Budi Cahyono, Ronny Lesmana

DOI: 10.18311/jnr/2016/7539, Pagination: 108-114



Abstract Views: 826 | Pdf Views: 472

Administration of Purwoceng (*Pimpinella alpina* Molk) Improves Oxidative Stress Biomarker Following UVC Irradiation in Spargue-Dawley Male Rats

Taufiqurrachman Nasihun, Eni Widayati

DOI: 10.18311/jnr/2016/7684, Pagination: 115-124



Abstract Views: 539 | Pdf Views: 516

Quick Links

- Abstracting and Indexing
- Instructions to Authors
- Pre-submission checklist
- Publication Fees
- Journal Policies and Ethics
- Submission Process
- Information for Reviewers
- Email Alerts
- CrossMark Policy

Archives

[Expand All]

- + 2019
- + 2018
- + 2017
- + 2016
- + 2015
- + 2014
- + 2013
- + 2012

More...

Current Issue

IF 10Y	1.0
RSC	2.0
RFC	1.0



Antioxidant and Anti Aging Assays of *Oryza Sativa* Extracts, Vanillin and Coumaric Acid

Wahyu Widowati^{1*}, Nurul Fauziah¹, Heddy Herdiman¹, Merry Afni², Ervi Afifah², Hanna Sari W. Kusuma², Hayatun Nufus², Seila Arumwardana² and Dwi Davidson Rihibiha²

¹Medical Research Center, Faculty of Medicine, Maranatha Christian University, Jl Prof Drg Surya Sumantri No 65 Bandung 40164, West Java, Indonesia

²Aretha Medika Utama, Biomolecular and Biomedical Research Center, Jl Babakan Jeruk 2, No 9, Bandung 40163, West Java, Indonesia

Abstract

Aging is a natural process in humans as accumulation of oxygen-derived free radicals which leads to the activation of hyaluronidase, collagenase and elastase, that can further contribute to cellular and tissue damage. Bioactive compounds from plants have been used as antioxidant that might inhibit aging processes as well. This study aimed to determine antioxidant and anti aging properties of *Oryza sativa* Extract (OSE), and its compounds, vanillin and coumaric acid. The phytochemical analysis of OSE was performed with Farnsworth modified method. Antioxidant activities were performed by measurement of 2,2-diphenyl 1-picrylhydrazyl (DPPH) free radical scavenger, Ferric Reducing Antioxidant Power (FRAP), and 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid) (ABTS) reducing activity, while anti aging assay were observed through inhibitory of elastase, collagenase, and hyaluronidase activities. Phytochemical analysis showed the presence of terpenoids and saponins in high level. OSE showed lowest DPPH activity ($IC_{50} = 314.51 \mu\text{g}/\text{mL}$) compared to vanillin ($IC_{50} = 283 \mu\text{g}/\text{mL}$) and coumaric acid ($IC_{50} = 255.69 \mu\text{g}/\text{mL}$). In ABTS assay, OSE resulted lowest activity ($IC_{50} = 145.67 \mu\text{g}/\text{mL}$), compared to vanillin ($IC_{50} = 4.96 \mu\text{g}/\text{mL}$) and coumaric acid ($IC_{50} = 1.67 \mu\text{g}/\text{mL}$). OSE also showed the lowest FRAP activity ($21.26 \mu\text{M Fe(II)}/\mu\text{g}$), compared to vanillin ($35.05 \mu\text{M Fe(II)}/\mu\text{g}$) and coumaric acid ($48.52 \mu\text{M Fe(II)}/\mu\text{g}$). OSE showed the lowest collagenase, elastase, and hyaluronidase inhibitory activities ($IC_{50} = 816.78, 107.51, \text{ and } 203.13 \mu\text{g}/\text{mL}$), compared to vanillin ($IC_{50} = 16.27, 14.46, 45.23 \mu\text{g}/\text{mL}$ respectively) and coumaric acid ($IC_{50} = 146.89, 25.38, 8.21 \mu\text{g}/\text{mL}$ respectively). In summary, OSE possess the lowest antioxidant and anti aging activities compared to vanillin and coumaric acid.

Keywords: Antioxidant, Anti aging, Coumaric Acid, *Oryza sativa*, Vanillin

1. Introduction

Skin aging is the natural process due to photo aging by environmental factors such as chronic UV radiation. The repetitive exposure to UV radiation cause accelerated physical changes in the skin and connective tissue through the formation of lipid peroxides, the cell contents and Reactive Oxygen Species (ROS)¹. It leads

to loss of skin elasticity implicated in formation of wrinkle, uneven pigmentation, brown spots, laxity and leathery appearance, solar elastosis, actinic purpura, precancerous lesions, skin cancer, and melanoma²⁻⁴.

During aging process, collagen, elastin, and hyaluronic acid decrease, that causes loss of strength and flexibility in the skin, resulting in visible wrinkles. It is also related to increasing enzymes activity including

*Author for correspondence

Email: wahyu_w60@yahoo.com

collagenase, elastase and hyaluronidase. Collagenase is known as an enzyme that plays role in the degradation of collagen. Collagen is the main component with percentage of 70-80 % of the total skin weight, the increasing degradation of collagen is significant in the photo aging process^{5,6}. Hyaluronan or hyaluronic acid is one of important components of the tissue matrix substance and has a role in the development, growth, and repair of damaged tissue⁶. Meanwhile, elastin play a role in the maintenance of skin elasticity, but elastase can degrade it⁷. Degradation of the Extracellular Matrix (ECM) has been directly linked to skin aging and is correlated with an increase in activity of certain enzymes involved in skin aging^{8,9}. Inhibition of these enzymes is crucial in anti aging prevention¹⁰.

It has been reported that skin aging occurs in the presence of cumulative endogenous damage due to Reactive Oxygen Species (ROS)¹¹. ROS are defined as oxygen-containing, highly reactive species. ROS are generated constantly during normal cellular metabolism which is essential for biological functions. Excessive ROS causes oxidative stress and damage of biological molecules^{12,13}. Previous studies have investigated that continuous ROS exposure can stimulate skin aging through antioxidant system destruction, wrinkle formation, and melanogenesis¹². ROS are usually eliminated from the body through antioxidant defense system¹⁴. Thus, maintaining antioxidant homeostasis is an appropriate strategy to prevent skin aging.




Fig. 1. Chemical structure of vanillin and coumaric acid.

Antioxidant properties derived from natural sources have been proposed for aging prevention¹⁵. Bioactive compounds contained in plants such as isoflavones, anthocyanins, and catechins may have promising antioxidant activity against ROS¹⁶. *Oryza sativa* L. is one of the most produced and consumed cereals in the world that contain phenolic compounds, tocopherols,

tocotrienols, and g-oryzanol. Phenolic acids were identified in the lignin fraction of rice bran (*O. sativa*) such as caffeic, chlorogenic, p-coumaric, ferulic, gallicacids, p-hydroxybenzoic, protocatechuic, syringic and vanillin¹⁷⁻¹⁹. Vanillin and coumaric acid have antioxidants activity that can inhibit aging processes^{20,21}. In the present study, free radical scavenging activity of *O. sativa* Extract (OSE) and its compounds, vanillin and coumaric acid (Figure 1) were evaluated as well as inhibitory activities of collagenase, elastase, and hyaluronidase.

2. Materials and Methods

2.1 Preparation of *O. sativa* Extracts

The plants of *O. sativa* were collected from the plantation in Ciharang, Subang, West Java. The plants were identified by herbarium staff, Department of Biology, School of Life Science and Technology, Bandung Institute of Technology, Bandung, West Java, Indonesia. The grain of *O. sativa* (600 g) were mashed, extracted using distilled ethanol 70% (2,750 mL) by a maceration method. Every 24 h the ethanol was filtered and the wastes were re-macerated in triplicate. The ethanol filtrate collected was condensed using 50°C rota vapor to obtain OSE. The extract in pasta form (5.64 g) was stored at -20 °C, and used for further assay²². Standards compounds used in this study were vanillin with 99% purity [Sigma V1104, USA] and coumaric acid (CA) with 98% purity [Biopurify Phytochemical 14111707, China].

2.2 Qualitative Phytochemical Screening Assay

The phytochemical assay was conducted on *O. sativa* Extracts (OSE) using modified Farnsworth method to qualitatively identify presence of phenols, steroid/triterpenoids, saponins, tannins, terpenoids, flavonoids, and alkaloids as listed below²³⁻²⁵.

2.2.1 Phenol Identification

Around 10 mg of sample was placed on a dropping plate, then 1% FeCl₃ [Merck 1.03861.0250, USA] was added

into the sample. The color formation of green/red/purple/blue/black shows presence of phenol²³⁻²⁵.

2.2.2 Steroid /Triterpenoid Identification

Approximately 10 mg of sample was placed on a dropping plate, then soaked with acetate acid until the sample was covered. After 10-15 min, one drop of absolute sulfate acid (H₂SO₄) [Merck 109073, USA] was added to the sample. The formation of green/blue color indicates the presence of steroid while red/orange sediment indicates the presence of triterpenoid²³⁻²⁵.

2.2.3 Saponin Identification

Approximately 10 mg of sample was put into the test tube with some water and boiled for 5 min. It was shaken vigorously and saponin content was indicated by persistence of froth on the surface²³⁻²⁵.

2.2.4 Tannin Identification

Approximately 10 mg of samples was added with 2 mL of HCl 2N [Merck 1003171000] in the test tube, then heated on a the water bath for 30 min. The mixture was cooled down and filtered, the filtrate was added with amyl alcohol [Merck 10979, USA]. Purple colour formation indicates positive reaction for tannins²³⁻²⁵.

2.2.5 Terpenoid Identification

Around 10 mg of sample was added into a dropping plate, then vanillin and H₂SO₄ was added to the sample. Terpenoid presence was indicated by the formation of purple color on the mixture²³⁻²⁵.

2.2.6 Flavonoid Identification

About 10 mg of sample was added into a test tube, then Mg [Merck EM105815, USA] and HCl 2N was added to the sample. The mixture sample was heated for 5 to 10 min, then filtered after it was cooled down. Subsequently, amyl alcohol was added into the filtrate. The positive reaction was shown by the formation of red or orange color²³⁻²⁵.

2.2.7 Alkaloid Identification

The small amount of sample (10 mg) was introduced into a test tube, then 10% ammonia was added into the sample. After chloroform added to the mixture, two

layers of liquid was formed and the bottom layer was collected. HCl 1N was added to the liquid, forming two layers. The upper layer collected and added with 1-2 drops of Dragendorff solution. The presence of yellow colour indicated positive result²³⁻²⁵.

2.3 2,2-Diphenyl-1-picrylhydrazil (DPPH) Assay

The DPPH assay was conducted using the method from Widowati *et al.*, study²⁶. The method is based on the reduction of alcoholic DPPH solution in the presence of a hydrogen-donating antioxidant due to the formation of the non-radical 2,2-diphenyl-1-picrylhydrazine (DPPH-H)²⁷. Briefly, 50 µL of various level of samples (50-400 µg/mL for extract and 50-400 µM for compounds in the DMSO) were added to each well in a 96-well micro plate. It was then followed by addition of 200 µl of 2,2-Diphenyl-1-picrylhydrazil (DPPH) [Sigma D9132, USA] solution (0.077 mmol/L in methanol) into the well. The mixture was then incubated in the dark for 30 min at room temperature. Afterward, the absorbance was read using a microplate reader (Multiskan™ GO Microplate Spectrophotometer, Thermo Scientific, Waltham, MA, USA) at 517 nm wave length. The radical scavenging activity was measured using the following formula:

$$\% \text{ Scavenging} = (A_c - A_s) / A_c \times 100$$

A_c = negative control absorbance (without sample).

A_s = sample absorbance.

2.4 ABTS-Reducing Activity Assay

The antioxidant capacity of OSE, vanillin, and CA were measured using 2,2'-Azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS^{•+}) [Sigma A1888-2G, USA] diammonium salt-free radical assay²⁸. ABTS^{•+} solution was produced by reacting 14 mM ABTS and 4.9 mM potassium persulfate [Merck EM105091, USA] (1:1 volume ratio) for 16 h in dark condition at room temperature, then the mixture was diluted with 5.5 mM PBS (pH 7.4) until the absorbance of the solution was 0.70 ± 0.02 at wavelength 745 nm. In brief, 2 µl of various level of samples were added to each well at 96-well microplate, then to the samples the fresh 198 µl ABTS^{•+} solution were added. The absorbance was

measured at 745 nm after the plate incubated for 6 min at 30°C. The percentage inhibition of ABTS radical (%) was determined by the ratio of reducing of ABTS^{•+} absorbance in the presence of the sample relative to the absorbance in the absence of the sample (negative control). The median Inhibitory Concentration (IC₅₀) were also calculated^{28,29}.

2.5 Ferric Reducing Antioxidant Power (FRAP) Assay

The Ferric Reducing Antioxidant Power Assay (FRAP) was estimated using modified method from Mishra *et al.*, and Widowati^{28,30}. The FRAP reagent was prepared freshly by mixing 10 mL of acetate buffer 300 mM, 1 mL of ferric chloride hexahydrate [Merck 1.03943.0250, USA] 20 mM dissolved in distilled water, and 1 mL of 2,4,6-Tris(2-pyridyl)-s-triazine (TPTZ) [Sigma 3682-35-7, USA] 10 mM dissolved in HCl 40 mM. In 96-well micro plate, 7.5 µL of various level of samples (12.5-100 µg/mL for OSE and 12.5-100 µM for compounds) were mixed with 142.5 µL FRAP reagent then incubated for 30 min at 37°C. The absorbance value was measured at 593 nm with a micro plate reader. The standard curve was made using FeSO₄, between 0.019 and 95 µg/mL FeSO₄. The results of samples were expressed in µM Fe (II)/µg extract²⁸.

2.6 Collagenase Inhibitory Activity Assay

Collagenase inhibitory activity was measured according to modified method of Sigma-Aldrich dan Thring *et al*⁷. Mixed solution included 10 µL Collagenase from *Clostridium histolyticum* [Sigma C8051, USA] (0.01 U/mL in the cool aquades), 60 µL *Tricine* buffer (50mM, pH 7.5, content of 10mM CaCl₂ dan 400mM NaCl), 30 µL of various level of sample (0-250 µg/mL for OSE and 0-250 µM for compounds in the DMSO), then incubated at 37 °C for 20 min. After incubated added 20 µL N-[3-(2-Furyl)acryloyl]-leu-gly-Pro-Ala [Sigma F5135, USA] (1mM in the *Tricine* buffer) substrate. Absorbance was measured at 335 nm wave length.

$$\% \text{ Collagenase inhibition} = (1-B/A) \times 100\%$$

A = sample absorbance

B = control absorbance

2.7 Elastase Inhibitory Activity Assay

Elastase inhibitory activity was measured by modified method of Sigma Aldrich and Thring *et al*⁷. A mixture of 10 µL of various level of samples (0-66.67 µg/mL for OSE and 0 - 66.67 µM for compounds), 5 µL elastase from porcine pancreas [Sigma 45124, USA] (0.5 mU/mL in the cool aquades) and 125 µL Tris buffer was pre-incubated for 15 min at 25°C. Mixed solution was added N-Sucanyl-Ala-Ala-Ala-p-Nitroanilide substrate [Sigma 54760, USA] and then incubated for 15 min at 25°C. Absorbance was measured by 410 nm wavelengths.

$$\% \text{ Elastase inhibition} = (1-B/A) \times 100\%$$

A = sample absorbance

B = control absorbance

2.8 Hyaluronidase Inhibitory Activity Assay

Hyaluronidase inhibitory of activity was measured by modified method of Sigma Aldrich and Tu and Tawata³¹. A mix of 25 µL of various level of samples (0-166.67 µg/mL for OSE and 0 - 166.67 µM for compounds) and 3 µL hyaluronidase from bovine testes type I-S [Sigma H3506, USA] was pre-incubated for 10 min at 37°C and then added 12 µL phosphate buffer (300mM, pH 5.35) for 10 min at 37°C. Afterward 10 µL hyaluronic acid substrate [Sigma H5542, USA] was added and incubated for 45 min at 37°C. Decomposition reaction of hyaluronic acid was stopped by adding 100 µL acidic albumin acid. Mixed solution incubated at room temperature for 10 min, then absorbance was measured at 600 nm wavelengths.

Quantification of inhibition activity by formula:

$$\% \text{ Hyaluronidase inhibition} = (1-B/A) \times 100\%$$

A = sample absorbance

B = control absorbance

3. Results

3.1 Phytochemical Screening of OSE

Phytochemical screening of the plants showed the presence of phenols, flavonoids, terpenoids, saponins,

tannins and alkaloids. The result of OSE phytochemical screening can be seen in Table 1.

Table 1: The result of qualitative phytochemical screening of OSE

Phytochemical content	OSE
Phenols	+
Steroids/Triterpenoids	-/+
Terpenoids	+++
Saponins	+++
Flavonoids	-
Tannins	-
Alkaloids	-

*++++: very high content; +++: high content; ++: moderate content; +: low content; -: not detected

Phytochemical screening of OSE aimed to detect presence of phenols, steroids, saponins, flavonoids, and tannins in OSE. Terpenoids and saponins was detected in high content (+++), phenols and triterpenoids were low content (+), while steroids, flavonoids, tannins and alkaloids were not detected (-) (Table 1).

3.2 2,2-Diphenyl-1-picrylhydrazil (DPPH) Assay

DPPH is a reagent for investigating the free radical scavenging activities of compounds. In the DPPH test, the extracts were able to reduce the stable radical DPPH to the yellow coloured diphenylpicrylhydrazine (DPPH-H)^{32,33}. The percentage of DPPH scavenging activity of OSE, vanillin, and coumaric acid can be seen in Figure 2 and the median Inhibitory Concentration (IC₅₀) of samples toward DPPH free radical scavenging activity can be seen in Table 2.




Fig. 2. DPPH scavenging activity of OSE, vanillin, and coumaric acid. OSE, vanillin, and coumaric acid were diluted in DMSO to reach the final concentration of 50.00; 100.00; 200.00; 400.00 (µg/mL; µM).

Table 2: IC₅₀ value DPPH scavenging activity of OSE, vanillin, and coumaric acid

Samples	Equation	R ²	IC ₅₀ (µM)	IC ₅₀ (µg/mL)
OSE	Y= 0.13x+8.36	0.98	-	314.51
Vanillin	Y=0.02x+16.05	0.93	1865.16	283.76
Coumaric Acid	Y= 0.02x+23.36	0.95	1557.78	255.69

*Linear equations, coefficient of regression (R²) and IC₅₀ of each sample were calculated. IC₅₀ of OSE was presented in µg/mL, while vanillin and coumaric acid were presented in µM and µg/mL.

Based on Figure 2, DPPH radical scavenging activity was concentration-dependent manner, in which higher concentration increased DPPH scavenging activity. At the highest concentration (400 µg/mL), OSE has the highest DPPH scavenging activity compared to vanillin and coumaric acid (59.62 ± 5.81%, 23.86 ± 0.57%, and 17.39 ± 0.16%, respectively) (Figure 2). However, IC₅₀ value of coumaric acid has the lowest value (255.69 µg/mL) compared to OSE (314.51 µg/mL) and vanillin (283.76 µg/mL) (Table 2). These results indicate OSE has low DPPH-scavenging activity among treatments.

3.3 ABTS - Reducing Activity Assay

ABTS-reducing activity assay measures the relative ability of antioxidant to scavenge the ABTS generated. The ABTS is generated by reacting a strong oxidizing agent (potassium permanganate/potassium persulfate) with the ABTS salt. Reduction of blue-green ABTS radical colored solution by hydrogen-donating antioxidant is measured by the long wave absorption spectrum³⁴. The result ABTS-reducing activity of OSE, vanillin, and coumaric acid based on IC₅₀ value can be seen in Table 3.

Table 3: IC₅₀ value ABTS-reducing activity of OSE, vanillin, and coumaric acid

Samples	Equation	R ²	IC ₅₀ (µM)	IC ₅₀ (µg/mL)
OSE	Y= 0.33x + 2.41	0.99	-	145.67
Vanillin	Y= 1.39x + 4.72	0.99	32.63	4.96
Coumaric Acid	Y= 3.17x+ 17.76	0.95	10.18	1.67

*Linear equations, coefficient of regression (R²) and IC₅₀ of each sample were calculated. IC₅₀ of OSE was presented in µg/mL, while vanillin and coumaric acid were presented in µM and µg/mL.

The antioxidant activity of OSE, vanillin, and coumaric acid were evaluated by ABTS-reducing activity assay. The IC_{50} of ABTS-reducing activity of OSE, vanillin, and coumaric acid can be seen in Table 3. OSE has the lowest ABTS-reducing activity as indicated by highest IC_{50} (145.67 $\mu\text{g}/\text{mL}$) compared to vanillin (4.96 $\mu\text{g}/\text{mL}$) and coumaric acid (1.67 $\mu\text{g}/\text{mL}$). The result indicated OSE has weak activity antioxidant compared to these compounds.

3.4 Ferric Reducing Antioxidant Power (FRAP) Assay

The Ferric Reducing Antioxidant Power (FRAP) method is based on the reduction of a ferroin analog, the Fe^{3+} complex of tripyridyltriazine $\text{Fe}(\text{TPTZ})^{3+}$ to the intensely blue coloured Fe^{2+} complex $\text{Fe}(\text{TPTZ})^{2+}$ by antioxidants in acidic medium. Absorbance of $\text{Fe}(\text{II})$ complex at 593 nm produced by antioxidant reduction of corresponding tripyridyltriazine $\text{Fe}(\text{III})$ complex³⁵. FRAP activity of OSE, vanillin, and coumaric acid can be seen in Figure 3.




Fig. 3. FRAP Activity of OSE, vanillin, and coumaric acid. OSE, vanillin, and coumaric acid were diluted in DMSO to reach the final concentration of 12.50; 25.00; 50.00; 100.00 ($\mu\text{g}/\text{mL}$; μM).

The result of the present study showed FRAP activity in concentration-dependent manner, in which higher concentration increased FRAP activity (Figure 3). At the highest concentration (100 $\mu\text{g}/\text{mL}$) OSE has a lowest FRAP with value (21.26 \pm 0.21 $\mu\text{M Fe(II)}/\mu\text{g}$) compared to vanillin (35.05 \pm 0.80 $\mu\text{M Fe(II)}/\mu\text{g}$) and coumaric acid (48.52 \pm 0.24 $\mu\text{M Fe(II)}/\mu\text{g}$). That indicates OSE has the lowest antioxidant activity in the FRAP assay compared to vanillin and coumaric acid.

3.5 Collagenase Inhibitory Activity

A spectrophotometric method was used to collagenase activity assay, and to detect potential collagenase inhibitor³⁶. The collagenase inhibitory activity of OSE, vanillin, and coumaric acid can be seen in Figure 4.




Fig. 4. Collagenase inhibitory activity of OSE, vanillin and coumaric acid. OSE, vanillin, and coumaric acid were diluted in DMSO to reach the final concentration of 31.25; 62.50; 125.00; 250.00 ($\mu\text{g}/\text{mL}$; μM).

Collagenase inhibitory activity of OSE, vanillin, and coumaric acid based on IC_{50} value can be seen in Table 4.

Table 4: IC_{50} value collagenase inhibitory activity of OSE, vanillin, and coumaric acid

Samples	Equation	R^2	IC_{50} (μM)	IC_{50} ($\mu\text{g}/\text{mL}$)
OSE	$y = 0.06x + 0.42$	0.99	-	816.78
Vanillin	$y = 0.15x + 34.16$	0.92	106.99	16.27
Coumaric Acid	$y = 0.05x + 3.47$	0.93	894.83	146.89

*Linear equations, coefficient of regression (R^2) and IC_{50} of each sample were calculated. IC_{50} of OSE was presented in $\mu\text{g}/\text{mL}$, while vanillin and coumaric acid were presented in μM and $\mu\text{g}/\text{mL}$.

Figure 4 shows collagenase activities of OSE coumaric acid and vanillin in concentration-dependent manner. Collagenase activities of OSE (15.46 \pm 1.12%) and coumaric acid (15.68 \pm 1.67%) were comparable, but lower than vanillin (68.91 \pm 1.56%). However, IC_{50} value of OSE was the highest (816.78 $\mu\text{g}/\text{mL}$) compared to coumaric acid (146.89 $\mu\text{g}/\text{mL}$) and vanillin (16.27 $\mu\text{g}/\text{mL}$) (Table 4). These findings indicate OSE has very low collagenase inhibitory activity, compared to coumaric acid and vanillin.

3.6 Elastase Inhibitory Activity

The elastase inhibitory activity of OSE, vanillin and coumaric acid were measured. The percentation elastase

inhibitory activity of OSE, vanillin and coumaric acid can be seen in Figure 5.




Fig. 5. Elastase inhibitory activity of OSE, vanillin, and coumaric acid. OSE, vanillin, and coumaric acid were diluted in DMSO to reach the final concentration of 8.33; 16.67; 33.33; 66.67 ($\mu\text{g/mL}$; μM).

The IC_{50} value an elastase inhibitory activity of OSE, vanillin, and coumaric acid based on IC_{50} value can be seen in Table 5.

Table 5: IC_{50} value elastase inhibitory activity of OSE, vanillin, and coumaric acid

Samples	Equation	R^2	IC_{50} (μM)	IC_{50} ($\mu\text{g/mL}$)
OSE	$Y=0.29x+19.07$	0.91	-	107.51
Vanillin	$Y=0.45x+7.31$	0.98	95.07	14.46
Coumaric Acid	$Y=0.22x+16.13$	0.94	154.66	25.38

*Linear equations, coefficient of regression (R^2) and IC_{50} of each sample were calculated. IC_{50} of OSE was presented in $\mu\text{g/mL}$, while vanillin and coumaric acid were presented in μM and $\mu\text{g/mL}$.

Elastase inhibitory activity of OSE, vanillin, and coumaric acid showed the highest inhibition percentage at the highest concentration ($36.82 \pm 0.40\%$, $36.28 \pm 0.20\%$, $29.86 \pm 0.45\%$ respectively) (Figure 5). However, vanillin showed the highest activity in elastase inhibition with IC_{50} value $14.46 \mu\text{g/mL}$. OSE and coumaric acid have IC_{50} value $107.62 \mu\text{g/mL}$ and $25.38 \mu\text{g/mL}$, respectively (Table 5). The result showed that OSE posses low elastase inhibition compared to vanillin and coumaric acid.

3.7 Hyaluronidase Inhibitory Activity

Hyaluronidase was assayed by a highly sensitive spectrophotometric method, based on precipitation of HA with cetylpyridinium chloride, which is used for high throughout screening for hyaluronidase inhibitors³⁷.

Hyaluronidase inhibitory activity can be used to evaluate anti aging activity of OSE, vanillin, coumaric acid. The percentation hyaluronidase inhibition activity of OSE, vanillin, and coumaric acid can be seen in Figure 6.




Fig. 6. Hyaluronidase inhibitory activity of OSE, vanillin, and coumaric acid. OSE, vanillin and coumaric acid were diluted in DMSO to reach the final concentration of 5.21; 10.42; 20.83; 41.67; 83.33; 166.67 ($\mu\text{g/mL}$; μM).

The IC_{50} value for hyaluronidase inhibitory activity of OSE, vanillin, and coumaric acid based on IC_{50} value can be seen in Table 6.

Table 6: IC_{50} value hyaluronidase inhibitory activity of OSE, vanillin, and coumaric acid

Samples	Equation	R^2	IC_{50} (μM)	IC_{50} ($\mu\text{g/mL}$)
OSE	$y=0.24x+0.34$	0.91	-	203.13
Vanillin	$y=0.15x+5.47$	0.99	297.33	45.23
Coumaric Acid	$y=0.89x+5.01$	0.94	50.11	8.21

*Linear equations, coefficient of regression (R^2) and IC_{50} of each sample were calculated. IC_{50} of OSE was presented in $\mu\text{g/mL}$, while vanillin and coumaric acid were presented in μM and $\mu\text{g/mL}$.

Based on Figure 6, OSE showed the moderate activity with percentage $36.86 \pm 2.28\%$ compared to coumaric acid ($48.48 \pm 0.40\%$) and vanillin ($20.84 \pm 4.78\%$). Coumaric acid has the lowest IC_{50} value ($8.22 \mu\text{g/mL}$) compared to OSE ($203.13 \mu\text{g/mL}$) and vanillin ($45.23 \mu\text{g/mL}$) (Table 6). These results show OSE has low hyaluronidase inhibitory activity compared to vanillin and coumaric acid.

4. Discussion

Ayurveda, which means science of long life, is a 5,000-year-old system of Indian medicine (1500–1000 BC) designed to promote good health and longevity

rather than to fight disease and was practiced by physicians and surgeons. Ayurveda usually recommends the use of several plant extracts in combination. Toxic effects of modern medicine are insufficient due to a lack of other components derived from the plants. Each herbal formulation contains multiple active compounds that may operate synergistically, producing therapeutic benefits and lowering the risks on adverse effects³⁸. Rice (*O. sativa*), a herb belongs to Poaceae, has been known as a potential source of antioxidants in food, pharmaceutical and cosmetic industries. It has been utilized in the treatment of cancer and aging³⁹. *O. sativa* has bioactive compounds, including many phenolic compounds as an antioxidant¹⁷. Five phenyl compounds isolated from the roots of *O. sativa* L. are vanillin, methyl trans-ferulate, trans-p-coumaric acid methyl ester, N-benzoyltryptamine, and N-(trans-cinnamoyl) tryptamine⁴⁰. Phenols can be divided into two subgroups according to their structure, p-hydroxybenzoic acid derivatives such as gallic, protocatechuic and syringic acids, and hydroxycinnamic derivatives such as caffeic, ferulic, p-coumaric and chlorogenic acids¹⁹. The present study showed that OSE contained low phenols and triterpenoids, high terpenoids, and saponins. Phytochemical analysis of *O. sativa* was conducted in the previous study that showed 11-13 % protein, approximately 11% fiber and 20% of its weight in oil, as well as functional compounds and antioxidants⁴¹. Phenolic compounds, such as benzoic or cinnamic acid derivatives, have been documented in Poaceae plants such as *Sorghum* sp^{42,43}.

UV irradiation generates free radicals that can induce expression of certain members of the Matrix Metalloproteinase (MMP) family, which degrades collagen and other Extracellular Matrix (ECM) proteins that consist of the dermal connective tissue⁴⁴. In this study, OSE had the lowest scavenging free radical activity as indicated by reduced ABTS and FRAP activities and DPPH scavenging activity compared to vanillin and coumaric acid. A recent study on *O. sativa* in different solvents showed that *O. sativa* ethanol extract have the lowest antioxidants (IC_{50} = 4.78 μ g/mL) compared to *O. sativa* methanol extract (IC_{50} = 2.6 μ g/mL) and *O. sativa* isopropanol extract (IC_{50} = 2.4 μ g/mL)⁴⁵. *O. sativa* methanol and isopropanol extracts exhibited better free radical scavenging activity, which is associated

with increasing phenolic compounds that increases antioxidant activity⁴⁶. Coumaric acid has been reported to protect against oxidative stress and play a key role as an antioxidant and anti-inflammatory^{47,48}. *O. sativa* oils such as coumaric acid and vanillin exhibited very good oxidative state with range oxidative stability 5.99-7.40⁴⁹.

OSE has the lowest DPPH scavenging activity compared to vanillin and coumaric acid, this data was not validated with previous study showed that *O. sativa* extract has rich secondary metabolites, especially alkaloids and phenolic acids that have strong antioxidant activity⁵⁰. Phenolic and flavonoid contents are associated with strong antioxidant activity and especially in terms of DPPH and nitric oxide free radical scavenging⁵¹. The phenolic extracts from fermented *O. sativa* showed slow inhibition kinetics of the DPPH radical, as IC_{50} value obtained was 250 mg/g compared to unfermented *O. sativa* with value 213 mg/g, while ferulic acid 235.00 mg/g⁵². In another study, Poaceae plants such as *Z. mays* ethanolic extract has low antioxidant activity with IC_{50} value of DPPH 163.45 ± 6.34 μ g/mL, while *T. aestivum* bran ethanol extract (0.5 mg/kg) has DPPH value $58.00 \pm 1.1\%$ ^{53,54}. *T. aestivum* methanol extract showed good radical scavengers with the inhibition of 12% at 1 mL/mL concentration⁵⁵. *T. aestivum* has shown potential anti-inflammatory, antioxidant, and anti aging properties⁵⁶.

Collagen, elastin, and hyaluronic acid are the skin main components and have an important role in maintaining its structure and hydration⁵⁷. These enzymes cause repetitive collagen fibers breakdown and responsible for structural defect in dermis and wrinkle development⁵⁸. Collagenase and elastase contribute in production and degradation of these fibers, which is also induced by free radical oxidative stress⁵⁹. In the present study, *O. sativa* had the lowest activity in collagenase inhibition compared to coumaric acid and vanillin. *O. sativa* has been studied in anti aging treatment. Water-soluble enzymatic extract from *O. sativa* served as a protective screen against UV-B radiation on cultivated keratinocytes and in vitro reconstructed skin⁶⁰. In the other study, the extract of *O. sativa* plants callus improve the human-skin barrier function⁶¹. Mechanisms of anti aging are via up-regulation of collagen synthesis in normal human dermal fibroblast and down-regulation of matrix metalloproteinase⁶². Vanillin content was also reported to inhibit matrix metalloproteinase-9 expression through

down-regulation of nuclear factor-kB signaling pathway in human hepatocellular carcinoma cells⁶³.

Elastin is a fibrose protein that compose 2–4% of the ECM and involved in the hydration of the skin⁶⁴. Elastase is responsible for increased tissue permeability, inflammation progress and delayed wound healing⁶⁵. Elastase is also the key enzyme that attacks all the major connective tissue matrix protein⁶⁶. In elastase assay, OSE showed the lowest activity compared to coumaric acid and vanillin. Low activity of OSE in the present study might be due to low content of phenol and absence of flavonoids in phytochemical screening. Several studies showed that phenol and flavonoid content of *O. sativa* possess anti-elastase activity^{51,67,68}. However, vanillin has the highest elastase inhibitory activity.

Our present study showed that OSE had the lowest hyaluronidase inhibitory activity compared to CA and vanillin. These results might be correlated with an absence of tannin in OSE. Tannin-rich plants have been reported to inhibit hyaluronidase and elastase release from stimulated neutrophils in vitro⁶⁹. Hyaluronidase (Haases) selectively degrade Hyaluronic Acid (HA), which is a megadalton acidic structural polysaccharide found exclusively in the Extracellular Matrix (ECM). Haase inhibitors are thus potent regulators that maintain HA homeostasis and they might serve as antiinflammatory, anti aging, antimicrobial, anticancer and antitoxin and contraceptive agents⁷⁰. Hyaluronidase degrades hyaluronic acid by lowering its viscosity and increasing the permeability⁷¹.

5. Conclusion

Vanillin and coumaric acid have higher antioxidant activity through DPPH scavenging, ABTS-reducing activities and FRAP, and anti-aging through inhibitory activity of collagenase, elastase and hyaluronidase than *O. sativa* extract.

6. Acknowledgment

This research was supported by Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung, Indonesia for a research grant, laboratory facilities, and research methodology. We are thankful to Neng Sri Mulkiyatul M. A, Kartiko Adi Poerbo and

Muhammad BiharJafarian as the interns at Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung, Indonesia for their valuable assistance.

7. References

1. Rattan SIS, Sodagam L. Gerontomodulatory and youth-preserving effects of zeatin on human skin fibroblasts undergoing aging in vitro. *Rejuvenation Res.* 2005; 8(1):46–57.
2. Kim DW, Hwang I, Kim D. Coenzyme Q10 effects on manganese superoxide dismutase and glutathione peroxidase in the hairless mouse skin induced by ultraviolet b irradiation. *Bio Factors.* 2007; 30(3):139–47.
3. Berneburg M, Plattenberg H, Medve-Konig K, Pfahlberg A, Gers-Barlag H, Gefeller O, *et al.* Induction of the photoaging-associated mitochondrial common deletion in vivo in normal human skin. *J Invest Dermatol.* 2004; 122(5):1277–83.
4. Yaar M, Eller MS, Gilchrist BA. Fifty years of skin aging. *J Invest Dermatol Symp Proc.* 2002; 7(1):51–8.
5. Demina N, Lysenko S. Collagenolytic enzymes synthesized by microorganisms. *Mikrobiologiya.* 1996; 65:293–304.
6. Pogrel M, Lowe M, Stern R. Hyaluronan (hyaluronic acid) in human saliva. *Arch Oral Biol.* 1996; 41(7):667–71.
7. Thring TS, Hili P, Naughton DP. Anti-collagenase, anti-elastase and anti-oxidant activities of extracts from 21 plants. *BMC Compl Alter Med.* 2009; 9(27):1–11.
8. Longo V, Finch C. Evolutionary medicine: From dwarf model systems to healthy centenarians? *Sci.* 2003; 299:1342–6.
9. Makrantonaki E, Brink T, Zampeli V, Elewa R, Mlody B, Hossini A. Identification of biomarkers of human skin ageing in both genders. Wnt signalling- a label of skin ageing? *PLoS ONE.* 2010; 7(11).
10. Ndlovu G, Fouche G, Tselanyane M, Cordier W, Steenkamp V. In vitro determination of the anti-aging potential of four Southern African medicinal plants. *BMC Complement Atern Med.* 2013; 13(304):1–7.
11. Yasui H, Sakurai H. Chemiluminescent detection and imaging of reactive oxygen species in live mouse skin exposed to UVA. *Biochem Biophys Res Commun.* 2000; 269(1):131–6.
12. Kim H, Yun J, Lee J, Hong H, Jeong J, Kim E, *et al.* SUMO1 attenuates stress-induced ROS generation by inhibiting NADPH oxidase 2. *Biochem Biophys Res Commun.* 2011; 410(3):555–62.

13. Mruka D, Silvestrini B, Mo My, Cheng C. Antioxidant superoxide dismutase- A review, its function, regulation in the testis, and role in male fertility. *Contracept.* 2002; 65:305–11.
14. Moini H, Packer L, Saris N. Antioxidant and prooxidant activities of alpha-lipoic acid and dihydrolipoic acid. *Toxicol Appl Pharmacol.* 2002; 182(1):84–90.
15. Cai Y, Luo Q, Sun M, Corke H. Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sci.* 2004; 74(17):2157–84.
16. Kaur C, Kapoor H. Anti-oxidant activity and total phenolic content of some Asian vegetables. *Int J Food Sci Technol.* 2002; 37(2):153–61.
17. Mira N, Barros R, Schiocchet M, Noldin J, Lanfer-Marquez U. Extraction, analysis and distribution of phenolic acids in pigmented and non-pigmented genotypes of rice (*Oryza sativa* L.). *Clencia e Tecnologia de Alimentos.* 2008; 28:994–1002.
18. Pourali O, Asghari FS, Yoshida H. Production of phenolic compounds from rice bran biomass under subcritical water conditions. *Chem Eng J.* 2010; 160(1):259–66.
19. Martins S, Mussatto S, Martinez-Avila G, Montanez-Saenz J, Aguilar C, Teixeira J. Bioactive phenolic compounds: production and extraction by solid-state fermentation. *Biotechnol Adv.* 2011; 29:365–73.
20. Yang HJ, Yue Q, Cao YC, Zhang DF, Wang JQ. Effects of crude feruloyl and acetyl esterase solutions of *Neocallimastix* sp. YQ1 and *Anaeromyces* sp. YQ3 isolated from holstein steers on hydrolysis of chinese wild rye grasshay, wheat bran, maize bran, wheat straw and corn stalks. *Anim Feed Sci Technol.* 2009; 154:218–27.
21. Kilic I, Yesiloglu Y. Spectroscopic studies on the antioxidant activity of p-coumaric acid. *Biomol Spectrosc.* 2013; 115:719–24.
22. Widowati W, Wijaya L, Wargasetia T, Bachtiar I, Yellianty Y, Laksmiawati D. Antioxidant, anticancer, and apoptosis-inducing effects of piper extracts in hela cells. *J Exp Integr Med.* 2013; 3(3):225–30.
23. Widowati W, Ratnawati H, Rusdi U, Winarno W, Imanuel V. Phytochemical assay and antiplatelet activity of fractions of velvet been seeds (*Mucuna pruriens*). *HAYATI J Biosci.* 2010; 17(2):85–90.
24. Bera T, Chatterjee K, Ghosh D. In vitro antioxidant properties of the hydro-methanol extract of the seeds of *Switenia mahagoni* (L.) Jacq. *BGM.* 2015; 7(1):18–24.
25. Adnyana I, Abuzaid A, Iskandar E, Kurniati N. Pancreatic lipase and a-amylase inhibitory potential of mangosteen (*Garcinia mangostana* Linn.) pericarp extract. *Int J Med Res Health Sci.* 2016; 5(1):23–8.
26. Widowati W, Maesaroh M, Fauziah N, Erawijantari PP, Sandra F. Free radical scavenging and a-/b-glucosidase inhibitory activities of rambutan (*Nephelium lappaceum* L.) peel extract. *Indones Biomed J.* 2015; 7(3):157–62.
27. Sohn D, Kim Y, Oh S, Park E, Li X, Lee B. Hepatoprotective and free radical scavenging effects of *Nelumbo nucifera*. *Phytomed.* 2003; 10(2-3):165–9.
28. Widowati W. Green tea extract protects endothelial progenitor cells from oxidative insult through reduction of intracellular reactive oxygen species activity. *J Basic Med Sci.* 2014; 17(9):702–9.
29. Etoundi C, Kuat D, Ngondi J, Oben J. Anti-amylase Anti-lipase and antioxidant effects of aqueous extracts of some cameroonian spices. *J Nat Products.* 2010; 3:165–71.
30. Mishra A, Bapat M, Tilak J, Devasagayam T. Antioxidant activity of *Garcinia indica* (kokam) and its syrup. *Curr Sci.* 2006; 91:90–3.
31. Tu P, Tawata S. Antioxidant, antiaging, and anti-melanogenic properties of the essential oil from two varieties of *Alpinia zerumbet*. *Molecules.* 2015; 20:16723–40.
32. Kumar S, Sandhir R, Ojha S. Evaluation of antioxidant activity and total phenol in different varieties of *Lantana camara* leaves. *BMC Res Notes.* 2014; 7(560):1–9.
33. Widowati W, Ratnawati H, Husin W, Maesaroh M. Antioxidant properties of spice extracts. *Biomed Eng.* 2015; 1(1):24–9.
34. Miller NJ, Rice-Evans CA. Factors influencing the antioxidant activity determined by the ABTS+ radical cation assay. *Free Radical Res.* 1997; 26:195–9.
35. Antolovich M, Prenzler PD, Patsalides E, McDonald S, Robards K. Methods for testing antioxidant activity. *Analyst.* 2002; 127:183–98.
36. Sumantran VN, Kulkarni AA, Harsulkar A, Wele A, Koppikar SJ, Chandwaskar R, et al. Hyaluronidase and collagenase inhibitory activities of the herbal formulation Triphala guggulu. *J Biosci.* 2007; 32(4):755–61.
37. Tung JS, Mark GE, Hollis GF. A microplate assay for hyaluronidase and hyaluronidase inhibitors. *Anal Biochem.* 1994; 223:149–52.
38. Garodia P, Ichikawa H, Malani N, Sethi G, Aggarwal BB. From ancient medicine to modern medicine: Ayurvedic concepts of health and their role in inflammation and cancer. *J Soc Integr Oncol.* 2007 Mar 21; 5(1):25–37.

39. Iqbal S, Bhangar M, Anwar F. Antioxidant properties and components of some commercially available varieties of rice bran in Pakistan. *Food Chem.* 2005; 93:265–72.
40. Cho JG, Huh J, Jeong RH, Baek NI. Inhibition effect of phenyl compounds from the *Oryza sativa* roots on melanin production in mMne B16-F10 melanoma cells. *Nat Prod Res.* 2014; 29(11):1–3.
41. Oliveira M, Feddern V, Kupski L, Cipolatti E, Badiale-Furlong E, Souza-Soares L. Changes in lipid, fatty acids and phospholipids composition of whole rice bran after solid-state fungal fermentation. *Bioresour Technol.* 2011; 102:8335–8.
42. Awika J, Rooney L. Sorghum phytochemicals and their potential impact on human health. *Phytochem.* 2004; 65:1199–221.
43. Dykes L, Rooney L. Sorghum and millet phenols and antioxidants. *J Cer Sci.* 2006; 44:236–51.
44. Quan T, Qin Z, Xia W, Shao Y, Voorhees J, Fisher GJ. Matrix-degrading metalloproteinases in photoaging. *J Investig.* 2009; 14(1):20–4.
45. Zubair M, Anwar F, Shahid SA. Effect of extraction solvents on phenolics and antioxidant activity of selected varieties of pakistani rice (*Oryza sativa*). *Int J Agric Biol.* 2012; 14:935–40.
46. Roginsky V, Lissi EA. Review of methods to determine chain breaking antioxidant activity in food. *Food Chem.* 2005; 92:235–54.
47. Ferguson L, Zhu S, Harris P. Antioxidant and antigenotoxic effects of plant cell wall hydroxycinnamic acids in cultured HT-29 cells. *Mol Nutr Food Res.* 2005; 49:585–93.
48. Svobodova A, Psotova J, Walterova D. Natural phenolics in the prevention of UV-induced skin damage. *Biomed Papers.* 2003; 147(2):137–45.
49. Anwar F, Bhangar M, Kazi T. Relationships of rancimat and AOM values at varying temperatures for several oils and fats. *J Am Oil Chem Sci.* 2003; 80:151–5.
50. Chung HS, Shin JC. Characterization of antioxidant alkaloids and phenolic acids from anthocyanin-pigmented rice (*Oryza sativa* cv. Heugjinjubyeo). *Food Chem.* 2007; 104:1670–7.
51. Karimi E, Mehrabanjoubani P, Keshavarzian M, Oskouelan E, Jaafar HZ, Abdolzadeh A. Identification and quantification of phenolic and flavonoid components in straw and seed husk of some rice varieties (*Oryza sativa* L.) and their antioxidant properties. *J Sci Food Agric.* 2014; 94:2324–30.
52. Schmidt CG, Goncalves LM, Prietto L, Hackbart HS, Furlong EB. Antioxidant activity and enzyme inhibition of phenolic acids from fermented rice bran with fungus *Rhizopus oryzae*. *Food Chem.* 2014; 146:371–7.
53. Liyana-Pathirana CM, Shahidi F. Antioxidant properties of commercial soft and hard winter wheats (*Triticum aestivum* L.) and their milling fractions. *J Sci Food Agric.* 2006; 86:477–85.
54. Liu J, Wang C, Wang Z, Zhang C, Lu S, Liu J. The antioxidant and free-radical scavenging activities of extract and fractions from corn silk (*Zea mays* L.) and related flavone glycosides. *Food Chem.* 2011; 126:261–9.
55. Arya P, Kumar M. Chemoprevention by *Triticum aestivum* of mouse skin carcinogenesis induced by DMBA and croton oil-association with oxidative status. *Asian Pacific J Cancer Prev.* 2011; 12(12):143–8.
56. Smith B. Generalization of spatially variant apodization to nonintegral nyquist sampling rats. *IEEE Trans Image Process.* 2000; 9:1088–93.
57. Hooda R. Antiwrinkle herbal drugs-an update. *J Pharmacog Phytochem.* 2015; 4(4):277–81.
58. Shehada A. A review on natural bioactive compounds as potential antiwrinkle agents. *World J Pharm.* 2014; 3:528–44.
59. Mukherjee P, Maity N, Nema N, Sarkar B. Bioactive compounds from natural resources against skin aging. *Phytomed.* 2011; 19:64–73.
60. Santa-Maria C, Revilla E, Miramontes E, Bautista J, Garcinia-Martinez A, Romero E, *et al.* Protection against free radicals (UVB irradiation) of a water-soluble enzymatic extract from rice bran. study using human keratinocyte monolayer and reconstructed human epidermis. *Food Chem Toxicol.* 2010; 48(1):83–8.
61. Ludwig P, Bennets S, Gruber J. Rice meristem stimulates epigenetic rejuvenation: New technologies. *S Afr Pharm Cosmet Rev.* 2011; 38:34–7.
62. Mitra R. Medicinal plants of Indonesia. *Asia Pac Biotech News.* 2007; 11:727–35.
63. Liang JA, Wu SL, Lo HY, Hsiang CY, Ho TY. Vanillin inhibits matrix metalloproteinase-9 expression through down-regulation of nuclear factor-kb signaling pathway in human hepatocellular carcinoma cells. *Mol Pharmacol.* 2009; 75(1):151–7.
64. Jenkins G. Molecular mechanisms of skin ageing. *Mech Ageing Dev.* 2002; 123:801–810.
65. Siedle B, Hrenn A, Merfort I. Natural compounds as inhibitors of human neutrophil elastase. *Planta Med.* 2007; 73:401–20.
66. Azmi N, Hashim P, Hashim DM, Halimoon N, Nik Majid NM. Anti-elastase, anti-tyrosinase and matrix

- metalloproteinase-1 inhibitory activity of earthworm extracts as potential new anti-aging agent. *Asian Pac J Trop Biomed.* 2014; 4(1):348–52.
67. Hrenn A, Steinbrecher T, Labahn A, Schwager J, Schempp C, Merfort I. Plant phenolics inhibit neutrophil elastase. *Planta Med.* 2006; 72:1127–31.
68. Kanashiro A, Souza J, Kabeya L, Azzolini A, Lucisano-Valim Y. Elastase release by stimulated neutrophils inhibited by flavonoids: Importance of the catechol group. *Z Naturforsch C.* 2007; 62(5-6):357–61.
69. Piwowarski J, Kiss AK, Kozłowska-Wojciechowska M. Anti-hyaluronidase and anti-elastase activity screening of tannin-rich plant material used in traditional polish medicine for external treatment of diseases with inflammatory background. *J Ethnopharmacol.* 2011; 137:937–41.
70. Girish K, Kemparaju K, Nagaraju S, Vishwanath B. Hyaluronidase inhibitors: A biological and therapeutic perspective. *Curr Med Chem.* 2009; 16:2261–88.
71. Girish K, Kemparaju K. The Magic Glue Hyaluronan its Eraser Hyaluronidase: A biological overview. *Sci.* 2007; 80:1921–43.

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau PEER REVIEW

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Antioxidant and Anti Aging Assays of Oryza sativa Extracts, Vanillin and Coumaric Acid

Jumlah Penulis : 9 Orang

Nama-nama Penulis : **Wahyu Widowati**, Nurul Fauziah, Heddy Herdiman, Merry Afni, Ervi Afifah, Hanna Sari W. Kusuma, Hayatun Nufus, Seila Arumwardana, Dwi Davidson Rihibiha

Status Penulis : Penulis Pertama / ~~Penulis ke-....~~ /-Penulis Korespondensi **)

Identitas Jurnal Ilmiah : a. Nama jurnal : Journal of Natural Remedies
b. Nomor ISSN : 2320-3358
c. Vol., No., Bulan, Tahun : Vol 16; No 3; July 2016
d. Penerbit : Informatics Publishing Limited
e. DOI Artikel (jika ada) : 10.18311/jnr/2016/7220
f. Alamat Web Jurnal : www.informaticsjournals.com
g. Terindeks di : Scopus Q4, SJR 0.154

Kategori Publikasi Jurnal Ilmiah: ~~Jurnal Ilmiah Internasional~~ / Internasional Bereputasi **)
(beri tanda \checkmark yang dipilih)

Jurnal Ilmiah Nasional Terakreditasi

Jurnal Ilmiah Nasional / Nasional terindeks ***)

HASIL PENILAIAN (Peer Review) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi <input checked="" type="checkbox"/>	Nasional Terakreditasi <input type="checkbox"/>	Nasional ***) <input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,6
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			11,5
c.	Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12			11,6
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			11,7
	Total	40			38,4

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur... Penulisan memenuhi standard karya ilmiah akademik, metode yang digunakan sesuai dan tepat digunakan pada penelitian ini
- b. Ruang lingkup & kedalaman pembahasan... Penelitian tentang potensi ekstrak beras putih sebagai antioksidan dan antiaging secara in vitro
- c. Kecukupan & kemutakhiran data serta metodologi... Ide dasar penelitian baik membahas ekstrak beras putih antioksidan metode DPPH, FRAP, ABTS, Antiaging, anti kolagenase, anti hyaluronidase, antielastase.
- d. Kelengkapan unsur dan kualitas penerbit... Jurnal JNR terindeks Scopus Q4 SJR 0,154. Penerbit Informatics Publishing Limited

c. Indikasi plagiasi
Tidak terdapat indikasi plagiarisme atau self plagiarisme

f. Kesesuaian bidang ilmu
Paper bidang biokimia, biomedik sesuai dengan bidang ilmu penulis

REVIEWER 1



(Prof. Dr. Chrismis Novalinda Ginting, M.Kes)

NIK : 0115127801

UNIVERSITAS PRIMA INDONESIA

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau *PEER REVIEW*

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Antioxidant and Anti Aging Assays of Oryza sativa Extracts, Vanillin and Coumaric Acid

Jumlah Penulis : 9 Orang

Nama-nama Penulis : **Wahyu Widowati**, Nurul Fauziah, Heddy Herdiman, Merry Afni, Ervi Afifah, Hanna Sari W. Kusuma, Hayatun Nufus, Seila Arumwardana, Dwi Davidson Rihibiha

Status Penulis : Penulis Pertama / ~~Penulis ke-....~~ / -Penulis Korespondensi **)

Identitas Jurnal Ilmiah :

- a. Nama jurnal : Journal of Natural Remedies
- b. Nomor ISSN : 2320-3358
- c. Vol., No., Bulan, Tahun : Vol 16; No 3; July 2016
- d. Penerbit : Informatics Publishing Limited
- e. DOI Artikel (jika ada) : 10.18311/jnr/2016/7220
- f. Alamat Web Jurnal : www.informaticsjournals.com
- g. Terindeks di : Scopus Q4, SJR 0.154

Kategori Publikasi Jurnal Ilmiah : ~~Jurnal Ilmiah Internasional~~ / Internasional Bereputasi **)

(beri tanda yang dipilih)

Jurnal Ilmiah Nasional Terakreditasi

Jurnal Ilmiah Nasional / Nasional terindeks (***)

HASIL PENILAIAN (*Peer Review*) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi <input checked="" type="checkbox"/>	Nasional Terakreditasi <input type="checkbox"/>	Nasional (***) <input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,5
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			11,5
c.	Kecukupan dan kemutakhiran data/ informasi dan metodologi (30%)	12			11,7
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			11,6
	Total	100%	40		38,5

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur
Jurnal ini sudah cukup lengkap dan ditemukan kesesuaian antara unsur dan lainnya
- b. Ruang lingkup & kedalaman pembahasan
Cakupan pembahasan sudah mendalam dan terluput juga kedalaman analisa data dan pembahasan
- c. Kecukupan & kemutakhiran data serta metodologi
Secara umum metodologi sudah memadai dan lengkap serta kemutakhiran data sudah terpenuhi
- d. Kelengkapan unsur dan kualitas penerbit
Kualitas penerbit sudah terdeteksi Internasional bereputasi Scopus Q4 SJR dan masuk kategori yang baik

c. Indikasi plagiasi
belum ditemukan adanya indikasi plagiasi

f. Kesesuaian bidang ilmu

Artikel ini sudah sesuai dengan bidang ilmu yang diteliti oleh penulis

REVIEWER 2



(Prof. Dr. Ermi Girsang, M. Kes)

NIK : 0117057501

UNIVERSITAS PRIMA INDONESIA

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau *PEER REVIEW*

KARYA ILMIAH : JURNAL ILMIAH

- Judul Karya Ilmiah (Artikel) : Antioxidant and Anti Aging Assays of *Oryza sativa* Extracts, Vanillin and Coumaric Acid
- Jumlah Penulis : 9 Orang
- Nama-nama Penulis : **Wahyu Widowati**, Nurul Fauziah, Heddy Herdiman, Merry Afni, Ervi Afifah, Hanna Sari W. Kusuma, Hayatun Nufus, Seila Arumwardana, Dwi Davidson Rihibiha
- Status Penulis : Penulis Pertama / ~~Penulis ke-....~~ / -Penulis Korespondensi **)
- Identitas Jurnal Ilmiah : a. Nama jurnal : Journal of Natural Remedies
 b. Nomor ISSN : 2320-3358
 c. Vol., No., Bulan, Tahun : Vol 16; No 3; July 2016
 d. Penerbit : Informatics Publishing Limited
 e. DOI Artikel (jika ada) : 10.18311/jnr/2016/7220
 f. Alamat Web Jurnal : www.informaticsjournals.com
 g. Terindeks di : Scopus Q4, SJR 0.154
- Kategori Publikasi Jurnal Ilmiah: Jurnal Ilmiah Internasional / Internasional Bereputasi **)
 (beri tanda \checkmark yang dipilih)
 Jurnal Ilmiah Nasional Terakreditasi
 Jurnal Ilmiah Nasional / Nasional terindeks ***)

HASIL PENILAIAN (Peer Review) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi	Nasional Terakreditasi	Nasional ***)	
		<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,55
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			11,5
c.	Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12			11,65
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			11,75
	Total	100%	40		38,45

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur. *Penulisan memenuhi standar karya ilmiah akademis, metode yang digunakan sesuai dan tepat digunakan pada penelitian ini.*
- Jurnal ini sudah cukup lengkap dan ditemukan kesesuaian antara unsur dan isinya.*
- b. Ruang lingkup & kedalaman pembahasan *Penelitian tentang potensi ekstrak beras putih sebagai antioksidan dan antiaging secara in vitro.*

Cakupan pembahasan sudah mendalam dan terdapat juga kedalaman analisa data dan pembahasan

c. Kecukupan & kemutakhiran data serta metodologi. Ide penelitian baik membahas ekstrak beras putih, antioksidan metode DPPH, FRAP, ABTS, antiaging, anti kolagenase, anti hyalurodi- nase dan anti elastase.

Secara umum metodologi sudah memadai dan lengkap serta kemutakhiran data sudah terpenuhi

d. Kelengkapan unsur dan kualitas penerbit. Penerbit Informatika Publishing Limited. Jurnal JNA terindeks Scopus Q4 SJR 0,154

Kualitas penerbit sudah terdeteksi Internasional bereputasi Scopus Q4 SJR dan masuk kategori yang baik

e. Indikasi plagiasi. Tidak terdapat indikasi plagiarisme atau self plagiarisme

Belum ditemukan adanya indikasi plagiasi

f. Kesesuaian bidang ilmu

Paper bidang biokimia (farmasi) sesuai dengan bidang ilmu penulis


Artikel ini sudah sesuai dengan bidang ilmu yang diteliti penulis

Medan,
Reviewer 2



(Prof. Dr. Ermi Girsang, M.Kes)
NIK : 0117057501
UNIVERSITAS PRIMA INDONESIA

Medan,
Reviewer 1



(Prof. Dr. Chrismis Novalinda Ginting, M.Kes)
NIK : 0115127801
UNIVERSITAS PRIMA INDONESIA