



ISSN: 2008-5874(Online)

ISSN: 2666-5866(Print)

IRANIAN JOURNAL
of
BASIC MEDICAL SCIENCES

IJBMS

Editor-in-Chief



Fazly Bazzaz, Bibi Sedigheh

Mashhad University of Medical Sciences, Mashhad, Iran
Specialty: **Pharmaceutical Microbiology**
Home Page: www.scopus.com/authid/detail.uri?authorId=8430945600
Email Address: fazlis@mums.ac.ir
Phone: 051-31801130

Assistant Editor



Malaekheh-Nikouei, Bizhan

Mashhad University of Medical Sciences, Mashhad, Iran
Specialty: **Pharmaceutics**
Home Page: www.scopus.com/authid/detail.uri?authorId=56012720400
Email Address: malaekheh@mums.ac.ir
Phone: 051-31801325



Roohbakhsh, Ali

Mashhad University of Medical Sciences, Mashhad, Iran
ORCID 0000-0001-5032-4263
Specialty: **Pharmacodynamics & Toxicology**
Home Page: www.scopus.com/authid/detail.uri?authorId=9335615700
Email Address: roohbakhsha@mums.ac.ir
Phone: 051-31801180



Arabi, Leila

Department of Pharmaceutical Nanotechnology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad
Specialty: **Pharmaceutical Nanotechnology**
Home Page: pharmacy.mums.ac.ir/index.php/43-persian-category/1337-drarabtlcv
Email Address: arabi@mums.ac.ir
Phone: 51-31801320

Editorial Board



Abbaszadegan, Mohammad Reza

Professor of Medical Genetics, Mashhad University of Medical Sciences
Specialty: **Genetics**
Home Page: rsc-medgenetics.mums.ac.ir/index.php/2015-03-11-10-59-12/43-persian-category/141-dr-abbaszadegan?format=pdf
Email Address: abbaszadeganmr@mums.ac.ir
Phone: 51-38002414



Abdollahi, Mohammad

Tehran University of Medical Sciences, Tehran, Iran
ORCID 0000-0003-0123-1209
Specialty: **Toxicology**
Home Page: www.scopus.com/authid/detail.uri?authorId=57200537974
Email Address: mohammad@tums.ac.ir
Phone: +98 21 64122319



Bazarbachi, Ali A.

American University of Beirut
Specialty: **Oncology-Hematology**
Home Page: www.scopus.com/authid/detail.uri?authorId=7006517199
Email Address: bazarbac@aub.edu.lb
Phone: (961)- 3-612434



Behravan, Javad

Mashhad University of Medical Sciences, Mashhad, Iran
Specialty: **Biotechnology**
Home Page: www.scopus.com/authid/detail.uri?authorId=11339398900
Email Address: behravanj@mums.ac.ir



Boskabady, Mohammad Hosein

Mashhad University of Medical Sciences, Mashhad, Iran
Specialty: **Physiology**
Home Page: www.scopus.com/authid/detail.uri?authorId=7004360435
Email Address: boskabadyhm@mums.ac.ir



Chuen Neng, LEE

University Surgical Cluster, National University Health System, Singapore 119228
Specialty: **Professor in Surgery, Engineering**
Home Page: paracrinetherapeutics.com/index.php/prof-lee-chuen-neng
Email Address: surlcn@nus.edu.sg



Eisenberg, Robert A.

University of Pennsylvania, Department of Medicine, Rheumatology Division, Philadelphia
Specialty: **Immunology**
Home Page: www.med.upenn.edu/apps/faculty/index.php/g275/p17860
Email Address: raemd@mail.med.upenn.edu
Phone: 215-573-9681



Fazel, Alireza

Mashhad University of Medical Sciences, Mashhad, Iran
ORCID 0000-0002-1470-619X
Specialty: **Anatomy and Histology**
Home Page: www.scopus.com/authid/detail.uri?authorId=7003843018
Email Address: fazela@mums.ac.ir

	<p>FERNs, Gordon A A Brighton and Sussex Medical School, Division of Medical Education, Brighton, United Kingdom ORCID  0000-0002-0957-8349 Specialty: Molecular & Metabolic Medicine Home Page: www.scopus.com/authorid/detail.uri?authorid=7005674972 Email Address: g.ferns@surrey.ac.uk Phone: 01483-688606</p>
	<p>Gorji, Ali Epilepsy Research Center, Robert-Koch-Str. 45, 48149 Münster Specialty: Physiology Home Page: www.scopus.com/authorid/detail.uri?authorid=7003473874 Email Address: gorjial@uni-muenster.de</p>
	<p>Hosseinzadeh, Hossein Mashhad University of Medical Sciences, Mashhad, Iran ORCID  0000-0002-3483-851X Specialty: Pharmacodynamics and Toxicology Home Page: www.scopus.com/authorid/detail.uri?authorid=56212454800 Email Address: hosseinzadehh@mums.ac.ir Phone: 51-38823252</p>
	<p>Mahmoudi, Mahmoud Mashhad University of Medical Sciences, Mashhad, Iran Specialty: Immunology Home Page: www.scopus.com/authorid/detail.uri?authorid=23050973200 Email Address: mahoudim@mums.ac.ir</p>
	<p>Malaekheh-Nikouei, Bizhan Mashhad University of Medical Sciences Specialty: Pharmaceutics Home Page: www.scopus.com/authorid/detail.uri?authorid=56012720400 Email Address: malaekhehb@mums.ac.ir Phone: 051-38823255</p>
	<p>Meshkani, Reza Department of Biochemistry, Faculty of Medicine, Tehran University of Medical Sciences Specialty: Clinical Biochemistry Home Page: www.scopus.com/authorid/detail.uri?authorid=14056524200 Email Address: rmeshkani@tums.ac.ir Phone: +98 21 64053426</p>
	<p>Monabati, Ahmad Shiraz University of Medical Sciences Specialty: Pathology Home Page: www.scopus.com/authorid/detail.uri?authorid=55393919600 Email Address: monabatia@sums.ac.ir Phone: 0711-2301784</p>
	<p>Nokhodchi, Ali JMS Building, School of Life Sciences University of Sussex, Brighton BN1 9QG, UK ORCID  0000-0002-3244-2482 Specialty: Pharmaceutics Home Page: www.scopus.com/authorid/detail.uri?authorid=7003819347 Email Address: a.nokhodchi@sussex.ac.uk</p>
	<p>Omid, Abbasali Mashhad University of Medical Sciences, Mashhad, Iran Specialty: Pathology Home Page: www.scopus.com/authorid/detail.uri?authorid=6507843666 Email Address: omidia@mums.ac.ir</p>
	<p>Ramezani, Mohammad Mashhad University of Medical Sciences, Mashhad, Iran ORCID  0000-0001-5888-6703 Specialty: Pharmacogenosy and Biotechnology Home Page: www.scopus.com/authorid/detail.uri?authorid=6701412858 Email Address: ramezanim@mums.ac.ir Phone: 051-37112470</p>
	<p>Sadeghi, Fatemeh Mashhad University of Medical Sciences, Mashhad, Iran Specialty: Pharmaceutics Home Page: www.scopus.com/authorid/detail.uri?authorid=7006326307 Email Address: sadeghi@f@mums.ac.ir</p>
	<p>Shakeeb Hasan, Moosavi Department of Biological and Medical Sciences, Faculty of Health and Life Sciences, Oxford Brookes University Specialty: Respiratory Medicine Home Page: www.brookes.ac.uk/bms/about/staff/?wid=BMS%20staff%20directory&op=full&uid=p0076413 Email Address: smoosavi@brookes.ac.uk</p>
	<p>Soleimani Rad, Jafar Tabriz University of Medical Sciences, Tabriz, Iran Specialty: Anatomy Home Page: www.scopus.com/authorid/detail.uri?authorid=41361975500 Email Address: soleimanij@tbzmed.ac.ir Phone: 0411-3313053</p>


[Articles in Press](#)
[Current Issue](#)
[Journal Archive](#)
[Volume 22 \(2019\)](#)
[Volume 21 \(2018\)](#)
[Issue 12](#)
[Issue 11](#)
[Issue 10](#)
[Issue 9](#)
[Issue 8](#)
[Issue 7](#)
[Issue 6](#)
[Issue 5](#)
[Issue 4](#)
[Issue 3](#)
[Issue 2](#)
[Issue 1](#)
[Volume 20 \(2017\)](#)
[Volume 19 \(2016\)](#)
[Volume 18 \(2015\)](#)
[Volume 17 \(2014\)](#)
[Volume 16 \(2013\)](#)
[Volume 15 \(2012\)](#)
[Volume 14 \(2011\)](#)
[Volume 13 \(2010\)](#)
[Volume 12 \(2009\)](#)
[Volume 11 \(2008\)](#)
[Volume 10 \(2007\)](#)

**Iranian Journal of
Basic Medical Sciences**

Rejection Rate	78.33%
----------------	--------

Average time to first decision (days)	19.31
---------------------------------------	-------

Volume & Issue: Volume 21, Issue 7, July 2018, Page 745-752, Page 651-769 [XML](#)

Review Article

1 ☐ Interaction of viral oncogenic proteins with the Wnt signaling pathway

Page 651-659

Sayyad Khanizadeh; Banafshe Hasanvand; Hamed Esmaeil Lashgarian; Mohammad Almasian; Gholamreza Goudarzi

[View Article](#) | [PDF \(630 K\)](#)

Original Article

2 ☐ Spermatogenesis after transplantation of adipose tissue-derived mesenchymal stem cells in busulfan-induced azoospermic hamster

Page 660-667

Negar Karimaghahi; Amin Tamadon; Farhad Rahmanifar; Davood Mehrabani; Alireza Raayat Jahromi; Shahrokh Zare; Zahra Khodabandeh; Iman Razeghian Jahromi; Omid Koohi-Hosseinabadi; Mehdi Dianatpour

[View Article](#) | [PDF \(1037 K\)](#)

3 ☐ The effect of vitamin C on mice hemolytic anemia induced by phenylhydrazine: an animal model study using histological changes in testis, pre-implantation embryo development and biochemical changes

Page 668-677

Hojat Anbara; Rasoul Shahrooz; Mazdak Razi; Hassan Malekinejad; GholamReza Najafi

[View Article](#) | [PDF \(1009 K\)](#)

4 ☐ AGS cell line xenograft tumor as a suitable gastric adenocarcinoma model: growth kinetic characterization and immunohistochemistry analysis

Page 678-681

Tahereh Barati; Mahnaz Haddadi; Fatemeh Sadeghi; Samad Muhammadnejad; Ahad Muhammadnejad; Ronak Heidarian; Motahareh Arjomandnejad; Saeid Amanpour

[View Article](#) | [PDF \(449 K\)](#)

5 ☐ Protective effects of troxerutin on maternal high-fat diet-induced impairments of spatial memory and apelin in the male offspring

Page 682-687

Rogaye Diba; Gisou Mohaddes; Fariba Mirzaie Bavi; Fereshteh Farajdokht; Parvin Bayandor; Maryam Hosseindoost; Keivan Mehri; Zohreh Zavvari Oskuye; Shirin Babri

[View Article](#) | [PDF \(492 K\)](#)

6 ☐ Human wild-type superoxide dismutase 1 gene delivery to rat bone marrow stromal cells: its importance and potential future trends

Page 688-694

Mohsen Abedi; Seyed Alireza Mesbah-Namin; Ali Noori-Zadeh; Taki Tiraihi; Taher Taheri

[View Article](#) | [PDF \(721 K\)](#)

7 ☐ Examination of the immunohistochemical localization and gene expression by RT-PCR of the oxytocin receptor in diabetic and non-diabetic mouse testis

Page 695-700

Ayşe AYDOĞAN; Seyit Ali BİNGÖL

[View Article](#) | [PDF \(1024 K\)](#)

8 ☐ Lipoprotein lipase gene polymorphisms as risk factors for stroke: a computational and meta-analysis

- 9 ☐ **Mild hypoxia and human bone marrow mesenchymal stem cells synergistically enhance expansion and homing capacity of human cord blood CD34+ stem cells**

Page 709-716

Fatemeh Mohammadali; saeid Abroun; Amir Atashi

[View Article](#) | [PDF \(1112 K\)](#)

- 10 ☐ **Gambogic acid inhibits LPS-induced macrophage pro-inflammatory cytokine production mainly through suppression of the p38 pathway**

Page 717-723

Ma Jianjun; Kangmao Huang; Yan Ma; Shuai Chen; Chao Liu; Zhi Shan; Xiangqian Fang

[View Article](#) | [PDF \(860 K\)](#)

- 11 ☐ **CX691 as an AMPA receptor positive modulator, improves the learning and memory in a rat model of Alzheimer's disease**

Page 724-730

Nazanin Mozafari; Ali Shamsizadeh; Iman Fatemi; Mohammad Allahtavakoli; Amir Moghadam-Ahmadi; Elham Kaviani; Ayat Kaeidi

[View Article](#) | [PDF \(1000 K\)](#)

- 12 ☐ **The effects of ginsenoside Rb1 on fatty acid β -oxidation, mediated by AMPK, in the failing heart**

Page 731-737

Hong-liang Kong; Ai-jie Hou; Ning-ning Liu; Bo-han Chen; Hua-ting Huang; Sheng-nan Dai

[View Article](#) | [PDF \(627 K\)](#)

- 13 ☐ **Design and synthesis of new esters of terpenoid alcohols as 15-lipoxygenase inhibitors**

Page 738-744

Hamid Sadeghian; Seyed Mohammad Seyed; Zeinab Jafari

[View Article](#) | [PDF \(592 K\)](#)

- 14 ☐ **Effects of insulin-like growth factor-induced Wharton jelly mesenchymal stem cells toward chondrogenesis in an osteoarthritis model**

Page 745-752

Wahyu Widowati; Ervi Afifah; Tjandrawati Mozef; Ferry Sandra; Rizal Rizal; Annisa Amalia; Yukko Arinta; Indra Bachtiar; Harry Murti

[View Article](#) | [PDF \(815 K\)](#)

- 15 ☐ **Subcutaneous administration of a fusion protein composed of pertussis toxin and filamentous hemagglutinin from *Bordetella pertussis* induces mucosal and systemic immune responses**

Page 753-759

Ali Torkashvand; Fariborz Bahrami; Minoo Adib; Soheila Ajdary

[View Article](#) | [PDF \(660 K\)](#)

- 16 ☐ **Relationship of cell surface hydrophobicity with biofilm formation and growth rate: A study on *Pseudomonas aeruginosa*, *Staphylococcus aureus*, and *Escherichia coli***

Page 760-769

Zulfiqar Ali Mirani; Aiman Fatima; Shaista Urooj; Mubashir Aziz; Muhammad Khan; Tanveer Abbas

[View Article](#) | [PDF \(739 K\)](#)

Effects of insulin-like growth factor-induced Wharton jelly mesenchymal stem cells toward chondrogenesis in an osteoarthritis model

Wahyu Widowati^{1*}, Ervi Afifah², Tjandrawati Mozef³, Ferry Sandra⁴, Rizal Rizal², Annisa Amalia², Yukko Arinta², Indra Bachtiar⁵, Harry Murti⁵

¹ Medical Research Center, Faculty of Medicine, Maranatha Christian University, Bandung 40164, West Java, Indonesia

² Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung 40163, West Java, Indonesia

³ Research Center for Chemistry, Indonesian Institute of Sciences, Serpong, Indonesia

⁴ Department of Biochemistry and Molecular Biology, Faculty of Dentistry, Trisakti University, Jakarta, Indonesia

⁵ Stem Cell and Cancer Institute, Jakarta 13210, Indonesia

ARTICLE INFO

Article type:
Original article

Article history:
Received: Dec 5, 2017
Accepted: Apr 9, 2018

Keywords:
Chondrogenesis
IGF-1
Mesenchymal stem cell
MMPs
Osteoarthritis

ABSTRACT

Objective(s): This study aimed to determine the collagen type II (COL2) and SOX9 expression in interleukin growth factor (IGF-1)-induced Wharton's Jelly mesenchymal stem cells (WJMSCs) and the level of chondrogenic markers in co-culture IGF1-WJMSCs and IL1 β -CHON002 as osteoarthritis (OA) cells model.

Materials and Methods: WJMSCs were induced with IGF1 (75, 150, and 300 ng/ml) to enhance their chondrogenesis capability. The gene expression of SOX9 and COL2 was evaluated with quantitative RT-PCR. Furthermore, IGF1-WJMSCs were co-cultured with IL1 β -CHON002 cells in varied ratios (1:2, 1:1, 2:1). Chondrogenic markers ADAMTS1, ADAMTS5, MMP3, MMP1, and RANKL were measured with ELISA.

Results: The IGF1-WJMSCs had an increased expression of COL2 and SOX9. ADAMTS1, ADAMTS5, MMP1, MMP3, and RANKL levels were decreased in the co-culture IGF1-WJMSCs and IL1 β -CHON002.

Conclusion: The IGF1-induced WJMSCs were capable to enhance chondrogenesis, indicated by increased expression of SOX9 and COL2 and decreased expression of ADAMTS1, ADAMTS5, MMP3, MMP1, and RANKL. These findings can be further used in the osteoarthritis treatment.

► Please cite this article as:

Widowati W, Afifah E, Mozef T, Sandra F, Rizal R, Amalia A, Arinta Y, Bachtiar I, Murti H. Effects of insulin-like growth factor-induced Wharton jelly mesenchymal stem cells toward chondrogenesis in an osteoarthritis model. Iran J Basic Med Sci 2018; 21:745-752. doi:10.22038/IJBMS.2018.28205.6840.

Introduction

Osteoarthritis (OA) is one of the common joint disorders worldwide and the main cause of body-support disability. The characteristics of OA include phenotypic changes in the superficial layer cells of the articular cartilage (AC), chondrocyte hypertrophy and apoptosis, progressive formation of osteophyte, fibrillation of the AC, sclerosis of subchondral bone, and increased remodeling of the periarticular bone (1-3).

The chondrocyte phenotype is characterized by specific genes expressions, i.e., collagen type II and the transcription factor SOX9 (4, 5). COL2 is an essential abundant component in the cartilage extra cellular matrix (ECM). Therefore, the COL2 disruption and loss of other cartilage ECM components during degenerative joint diseases such as OA will lead to severe disability and aging-related health problems (6). This may be stimulated by complex pathogenic mechanisms that decreased matrix synthesis and upregulated pathways of tissue degradation (7).

Stem cells take a role in novel treatment strategies for both clinical situations. Johnson *et al.* discovered a drug candidate by screening the small molecules that induced mesenchymal stem cells (MSCs) chondrogenic differentiation (8). MSCs were studied for cartilage development (9), which may be helpful for the developmental programs in OA.

MSCs were collected from cartilage (bone marrow mesenchymal stem cells/BM-MSCs) and subsequently from adipose tissue, the placenta, dental pulp, umbilical cord, amnion (10), and Wharton's Jelly (11). Adipose tissue-MSCs (AD-MSCs) have lower chondrogenesis ability than BM-MSCs. Induction of TGF β 2 and IGF1 in AD-MSCs may produce chondrocytes that are slightly inferior to BM-MSCs chondrocytes, as measured using chondrocytes markers including COL1A, COL2A1, and SOX-9 (12).

The insulin-like growth factor (IGF1) is an enhancer that is responsible for the rate of gene expression (13-16), and IGF-2 plays a role as a growth stimulant in a non-differentiated state and as a regulator for glucose in all stages of differentiation (17, 18). A study reported that plasmid-based upregulation of IGF-1 in rabbit chondrocytes encapsulated using alginate *in vivo* showed an ability to repair cartilage and accelerated subchondral bone reformation in osteochondral disorders (19). Thus, we aimed to observe IGF-1 induction effect on the gene expression of chondrogenic markers, SOX9 and COL2, in Wharton's Jelly MSCs.

Materials and Methods

Cell culture preparation

The human Wharton's Jelly mesenchymal stem cells (hWJMSCs) of passage 4 (P4) were collected from the

*Corresponding author: Wahyu Widowati. Faculty of Medicine, Maranatha Christian University, Bandung 40164, Indonesia. Tel: +62-819-1004-0010; Email: wahyu_w60@yahoo.com

Stem Cell and Cancer Institute (Jakarta, Indonesia). The cells had been characterized by the cell multipotent differentiation and surface phenotype (11, 20). Informed consent was obtained from the Institutional Ethics Committee at the Stem Cell and Cancer Institute (Jakarta, Indonesia).

The hWJMSCs at a density of 5×10^5 /well were cultured in minimum essential medium- α (α -MEM) (Gibco, 12561056), supplemented with fetal bovine serum (20%) (FBS, Gibco, 10270106) and 1% antibiotic and anti-mycotic (Gibco, 15240062). They were incubated in a humidified atmosphere with 5% CO₂ at 37 °C for 24 hr. The medium was discarded and washed with Phosphate Buffered Saline (PBS).

hWJMSCs at density 1×10^6 cells/well was maintained in a complete medium. The cells were treated with IGF-1 (Biolegend, 590904) at concentrations of 75, 150, and 300 ng/ml, and incubated at 5% CO₂, 37 °C for 7 days, to obtain IGF1-induced WJMSCs cells for measuring SOX9 and COL2 gene expression (11, 20).

Co-culture of IGF1-WJMSCs and IL1 β -CHON002

IGF1 75 ng/ml-induced WJMSCs, IGF1 120 ng/ml-induced WJMSCs (IGF1-WJMSCs), IL1 β 5 ng/ml-induced CHON002, and IL1 β 10 ng/ml-induced CHON002 (IL1 β -CHON002) cells were collected with PBS containing 0.15% (w/v) trypsin (2000 units/g) and 0.02% EDTA. The cell suspensions of IGF1-WJMSCs and IL1 β -CHON002 were mixed at five ratios including WJMSCs alone, IGF1-WJMSCs and IL1 β -CHON002 1:2, 1:1, 2:1, IGF1-WJMSCs alone, IL1 β -CHON002 alone, and CHON002 alone. The cells were stored at 37 °C in 5% CO₂, with medium changes three times per week. The co-culture was stained with alcian blue, after 2 weeks of incubation. Meanwhile, the conditioned medium (CM) was collected from the co-culture and evaluated by ELISA assay (21).

Quantification of COL2 and SOX9

RNA was extracted using the Aurum total RNA kit (Bio-Rad, 7326820) and checked its concentration. The RNA was used for cDNA synthesis using iScript cDNA synthesis kit (Bio-Rad, 1708890) at 25 °C temperature for 5 min, 42 °C for 30 min, and 85 °C for 5 min for the final step. The end-product was stored at -20 °C.

Quantitative gene expression was conducted using Thermo Scientific PikoReal Real-time PCR System (Thermo Fisher). PCR followed these condition: pre-incubation cycle (95 °C for 5 min), 40 cycles of denaturation (95 °C for 1 min), annealing (47 °C for 40 sec for SOX9, and 52 °C for 40 sec for COL2), and extension (72 °C for 1 min). The reaction mix that was used to perform qPCR was from an Evagreen master mix (Bio-Rad, 1725200). Table 1 shows the primers used in this research.

Table 1. Primer sequences

Primer	Forward	Reverse
Beta actin	5'-TCTGGCACCACACCTTCTACAATG-3'	5'-AGCACAGCCTGGATAGCAACG-3'
SOX-9	5'-TTCGGTTATTTTAGGATCATCTCG-3'	5'-CACACAGCTCACTCGACCTTG-3'
COL-2	5'-GGCAATAGCAGGTTACAGTACA-3'	5'-CGATAACAGTCTTGCCCCACTT-3'

Levels of ADAMTS1, ADAMTS5, MMP1, MMP3, and RANKL

Levels of ADAMTS1, ADAMTS5, MMP1, MMP3, and RANKL were assessed in accordance with manufacturer's protocols (Elabscience, H5539, H5590, H1441, H1446, and H2400, respectively). Briefly, 100 μ l samples or standards were added into wells and incubated at 37 °C for 90 min before they were discarded. 100 μ l of biotinylated was added, stored at 37 °C for 60 min, and washed with 350 μ l wash buffer three times. Afterwards, the 100 μ l of HRP conjugate was added, then incubated at 37 °C for 30 sec and washed with 350 μ l wash buffer five times. 90 μ l of substrate was added and incubated at 37 °C for 15 min until it turned blue. Then, 50 μ l of stop solution was added until yellow changes appeared. Absorbance test was carried out at 450 nm.

Results

Levels of SOX9 and COL2

SOX9 is present in differentiated chondrocytes and all chondrocyte progenitors during chondrogenesis. However, the expression is completely turned off in hypertrophic chondrocytes (22). SOX9 expression parallels that of the gene coding for COL2A1, a chondrocyte differentiation specific marker (23). SOX9 and COL2 were measured in WJMSCs induced by IGF1 (Figure 1).

Figure 1 shows that the expression of SOX9 and COL2 at IGF1 at concentration 150 ng/ml was higher, with values of (1.15 ± 0.07) and (8.44 ± 0.44) , respectively compared to IGF1 300 ng/ml, IGF1 75 ng/ml, and control. This indicates that IGF1-induced WJMSCs cells can improve chondrogenesis in controlling differentiation of chondrogenic cells, which can repair chondrocyte damage in OA. For co-culture of IL1 β -induced CHON002 and IGF1-induced WJMSCs, we used IGF1 75 ng/ml and




Figure 1. Gene expression levels of SOX9 and COL2 on IGF1-induced WJMSCs

*The data was presented as a histogram of mean \pm standard deviation. Different letters (a,b) are significant among various concentrations of IGF1 (0, 75, 150, and 300 ng/ml) toward level of SOX9 gene expression, different letters (a,b,c,d) are significant among concentrations of IGF toward COL2 gene expression based on Duncan's *post hoc* test ($P < 0.05$)




Figure 2. Levels of ADAMTS1 in IL1 β -CHON002 and IGF1-WJMSCs co-cultures

A. Effect varied ratios of non-induced WJMSCs and non-induced CHON002 toward ADAMTS1 level. Different letters (a,ab,abc,bc,c) are significant among ratios of WJMSCs and CHON002 (1:0, 1:2, 2:1, 1:1, 0:1) toward ADAMTS1 level based on Duncan's *post hoc* test ($P < 0.05$)

B. Effect varied ratios of IGF1 75 ng/ml-induced WJMSCs and IL1 β 5 ng/ml-induced CHON002 toward ADAMTS1 level. Different letters (a,b,c) are significant among ratios of IGF1 75 ng/ml induced-WJMSCs and IL1 β 5 ng/ml-induced CHON002 (1:0, 1:2, 2:1, 1:1, 0:1) toward ADAMTS1 level based on Duncan's *post hoc* test ($P < 0.05$)

C. Effect varied ratios of IGF1 120 ng/ml-induced WJMSCs and IL1 β 10 ng/ml-induced CHON002 toward ADAMTS1 level. Different letters (a,b,c) are significant among ratios of IGF1 120 ng/ml induced-WJMSCs and IL1 β 10 ng/ml-induced CHON002 (1:0, 1:2, 2:1, 1:1, 0:1) toward ADAMTS1 level based on Duncan's *post hoc* test ($P < 0.05$)




Figure 3. Levels of ADAMTS5 in IL1 β -CHON002 and IGF1-WJMSCs co-cultures

A. Effect of varied ratios of non-induced WJMSCs and non-induced CHON002 toward ADAMTS5 level. Different letters (a,ab,b) are significant among ratios of WJMSCs and CHON002 (1:0, 1:2, 2:1, 1:1, 0:1) toward ADAMTS5 level based on Duncan's *post hoc* test ($P < 0.05$)

B. Effect of varied ratios of IGF1 75 ng/ml-induced WJMSCs and IL1 β 5 ng/ml-induced CHON002 toward ADAMTS5 level. Different letters (a,ab,bc,c) are significant among ratios of IGF1 75 ng/ml induced-WJMSCs and IL1 β 5 ng/ml-induced CHON002 (1:0, 1:2, 2:1, 1:1, 0:1) toward ADAMTS5 level based on Duncan's *post hoc* test ($P < 0.05$)

C. Effect varied ratios of IGF1 120 ng/ml-induced WJMSCs and IL1 β 10 ng/ml-induced CHON002 toward ADAMTS5 level. Different letters (a,ab,bc,c,d) are significant among ratios of IGF1 120 ng/ml induced-WJMSCs and IL1 β 10 ng/ml-induced CHON002 (1:0, 1:2, 2:1, 1:1, 0:1) toward ADAMTS5 level based on Duncan's *post hoc* test ($P < 0.05$)

120 ng/ml for inducing WJMSCs.

Level of ADAMTS1 and ADAMTS5

ADAMTS1 is presented within cartilage and the synovium (24) and its expression is significantly upregulated in OA cartilage (25). ADAMTS1 and ADAMTS5 levels are shown in Figures 2 and 3, respectively.

In general, co-cultures of IGF1-induced WJMSCs and IL1 β -induced CHON002 with ratio 1:2 showed lowest ADAMTS1 among treatments (Figure 2); co-culture of IGF 75 ng/ml-induced WJMSCs and IL1 β 5 ng/ml-induced CHON002 (1:2) showed lowest ADAMTS1

level (3.15 ng/ml), and it was significantly different compared to the positive control, IL1 β 5 ng/ml-induced CHON002, and IL1 β 10 ng/ml-induced CHON002 (10.74 and 14.64 ng/ml, respectively). However, there was no marked difference between IGF1-induced WJMSCs alone, in which ADAMTS1 levels in both treatments (IGF1 75 ng/ml, 120 ng/ml) were 11.44 and 10.31 ng/ml, respectively.

On the other hand, co-culture IGF1 120 ng/ml-induced WJMSCs and IL1 β 10 ng/ml-induced CHON002 (2:1) showed the lowest ADAMTS5 level (24.83 ng/ml), which was significantly different compared to positive control, IL1 β 5 ng/ml-induced CHON002, and IL1 β 10

ng/ml-induced CHON002 (39.48 and 45.58 ng/ml, respectively). The result was comparable to negative control, CHON002 alone without induction (25.36 ng/ml). Single culture and IGF1-induced WJMSCs alone (IGF1 75 ng/ml, 120 ng/ml) also presented the low levels of ADAMTS5 (25.89 and 26.95 ng/ml, respectively).

Level of MMP1 and MMP3

The levels of MMP1 and MMP3 in co-culture IGF1-WJMSCs and IL1 β -CHON002 are shown in Figures 4 and 5, respectively.

The results of the present study show that both IGF1-WJMSCs itself and co-culture with IL1 β -CHON002 significantly decreased the level of MMP1 (Figure

4) compared to positive control namely, IL1 β 5 ng/ml-induced CHON002 and IL1 β 10 ng/ml-induced CHON002 (7.19 and 8.73 ng/ml, respectively). These results were comparable to negative control, CHON002 alone without induction (6.49 ng/ml). Similar results were also found in MMP3 levels in which both IGF1-induced WJMSCs alone and co-culture significantly decreased the MMP3 level compared to IL1 β -induced CHON002 (Figure 5).

Level of RANKL

The TNF molecules called RANKL (receptor activator of NF κ B ligand) is the main regulator of bone remodeling




Figure 4. Levels of MMP1 in IL1 β -CHON002 and IGF1-WJMSC co-cultures

A. Effect of varied ratios of non-induced WJMSCs and non-induced CHON002 toward MMP1 level. Different letters (a,b,c) are significant among ratios WJMSCs and CHON002 (1:0, 1:2, 2:1, 1:1, 0:1) toward MMP1 level based on Duncan's *post hoc* test ($P<0.05$)

B. Effect of varied ratios of IGF1 75 ng/ml-induced WJMSCs and IL1 β 5 ng/ml-induced CHON002 toward MMP1 level. Different letters (a,ab,b,c) are significant among ratios of IGF1 75 ng/ml induced-WJMSCs and IL1 β 5 ng/ml-induced CHON002 (1:0, 1:2, 2:1, 1:1, 0:1) toward MMP1 level based on Duncan's *post hoc* test ($P<0.05$)

C. Effect of varied ratios of IGF1 120 ng/ml-induced WJMSCs and IL1 β 10 ng/ml-induced CHON002 toward MMP1 level. Different letters (a,ab,b,c) are significant among ratios of IGF1 120 ng/ml induced-WJMSCs and IL1 β 10 ng/ml-induced CHON002 (1:0, 1:2, 2:1, 1:1, 0:1) toward MMP1 level based on Duncan's *post hoc* test ($P<0.05$)




Figure 5. Levels of MMP3 in IL1 β -CHON002 and IGF1-WJMSCs co-cultures

A. Effect of varied ratios of non-induced WJMSCs and non-induced CHON002 toward MMP3 level. Different letters (a,b) are significant among ratios of WJMSCs and CHON002 (1:0, 1:2, 2:1, 1:1, 0:1) toward MMP3 level based on Duncan's *post hoc* test ($P<0.05$)

B. Effect varied ratios of IGF1 75 ng/ml-induced WJMSCs and IL1 β 5 ng/ml-induced CHON002 toward MMP3 level. Different letters (a,b) are significant among ratios of IGF1 75 ng/ml induced-WJMSCs and IL1 β 5 ng/ml-induced CHON002 (1:0, 1:2, 2:1, 1:1, 0:1) toward MMP3 level based on Duncan's *post hoc* test ($P<0.05$)

C. Effect of varied ratios of IGF1 120 ng/ml-induced WJMSCs and IL1 β 10 ng/ml-induced CHON002 toward MMP3 level. Different letters (a,ab,b,c) are significant among ratios of IGF1 120 ng/ml induced-WJMSCs and IL1 β 10 ng/ml-induced CHON002 (1:0, 1:2, 2:1, 1:1, 0:1) toward MMP3 level based on Duncan's *post hoc* test ($P<0.05$)




Figure 6. Level of RANKL in IL1 β -CHON002 and IGF1-WJMSCs co-cultures

A. Effect of varied ratios of non-induced WJMSCs and non-induced CHON002 toward RANKL level. Different letters (a, ab, bc, c) are significant among ratios WJMSCs and CHON002 (1:0, 1:2, 2:1, 1:1, 0:1) toward RANKL level based on Duncan's *post hoc* test ($P < 0.05$)

B. Effect of varied ratios of IGF1 75 ng/ml-induced WJMSCs and IL1 β 5 ng/ml-induced CHON002 toward RANKL level. Different letters (a,b,c,d) are significant among ratios of IGF1 75 ng/ml induced-WJMSCs and IL1 β 5 ng/ml-induced CHON002 (1:0, 1:2, 2:1, 1:1, 0:1) toward RANKL level based on Duncan's *post hoc* test ($P < 0.05$)

C. Effect of varied ratios of IGF1 120 ng/ml-induced WJMSCs and IL1 β 10 ng/ml-induced CHON002 toward RANKL level. Different letters (a,b,c,d) are significant among ratios of IGF1 120 ng/ml induced-WJMSCs and IL1 β 10 ng/ml-induced CHON002 (1:0, 1:2, 2:1, 1:1, 0:1) toward RANKL level based on Duncan's *post hoc* test ($P < 0.05$)

and development and activation of osteoclasts. The levels of RANKL can be seen in Figure 6.

The results obtained show that there was a significant difference in RANKL levels between the treatment of IGF1-WJMSCs and IL1 β -CHON002 (Figure 6). Level of RANKL was reduced by both IGF1-WJMSCs alone and co-culture, and it was significant compared to the negative control, IL1 β 5 ng/ml-induced CHON002, and IL1 β 10 ng/ml-induced CHON002. Co-culture IGF1 75 ng/ml-induced WJMSCs and IL1 β 5 ng/ml-induced CHON002 (2:1) showed the lowest RANKL levels among co-cultures.

Discussion

MSCs have been employed as one particular sector of tissue engineering that involves the repair, replacement, or regeneration of cartilage tissue, due to their superior proliferative and differentiation capacities (26). It has been reported that WJMSCs may differentiate into chondrocytes, skeletal muscle cells, cardiac muscle cells, osteoblasts, adipocytes, β cells in the islets of Langerhans, or endothelial cells *in vitro* (27). Hence, these cells may be applied in the treatment of chronic degenerative disorders and prevent cartilage degradation in patients with OA through their trophic/regenerative potential.

In the present study, treating WJMSCs with IGF1 increased expression of SOX9 (Figure 1). Many cartilage matrix genes have been indicated to be under the regulation of transcriptional control of SOX9. They include COL2A1, COL9A1, COL11A2, aggrecan, and cartilage link protein (CRTL1) genes (28–31), all of which are involved in articular cartilage structure and function. Furthermore, SOX9 is presented and present

in presumptive cartilage during embryo development. The mutations in human SOX9 gene leads to campomelic dysplasia with skeletal malformation and dwarfism (32). Thus, downregulation of SOX9 in OA is clearly likely to contribute to cartilage pathology.

Moreover, induction of IGF1 150 ng/ml in WJMSCs also increased expression of COL2 (Figure 1). Referring to previous studies, SOX9 overexpression in human chondrocytes increases COL2A1 expression, as well as their capacity to reform a cartilage ECM (33–35). Collagenase-1, -2, and -3 are all synthesized by chondrocytes and have been considered as the rate-limiting enzymes in collagen degradation (36–38). Collagenase levels in the synovial fluid and serum correlate with cartilage destruction in OA (39). An imbalance between collagenases and their endogenous inhibitors has also been suggested to result in cartilage collagenolysis (40).

The results obtained in this study are in line with previous studies that proteoglycan core protein and collagen type II are induced by IGF1 and that it stabilizes the chondrocyte phenotype in pathological conditions (41–43). IGF-1 is fairly mitogenic in human adult articular cartilage and highly stimulates the production of chondrocyte extracellular matrix components (13, 14).

Furthermore, IGF1 with concentrations of 75 ng/ml and 120 ng/ml was used to induce WJMSCs which were co-cultured with IL1 β (5 ng/ml, 10 ng/ml)-induced CHON002. These IGF1 concentrations were used in this study because SOX9 expression is not appropriate. According to Kim and Im, 2009, supplementation of mesenchymal stem cells with IGF, did not induce

SOX9 significantly (44). Parameters measured were chondrogenic markers that included ADAMTS-5, ADAMTS-1, MMP-3, MMP-1, and RANKL. Elevation of MMP-1 (collagenase-1) and MMP-3 (stromelysin-1) have been documented in osteoarthritic cartilage (45, 46) and in the synovial fluid of osteoarthritic joints (47, 48). The present study has recognized the messenger RNA (mRNA) presence for some MMPs, i.e., MMP13, MMP9, MMP3, and MMP1, in human OA cartilage (49, 50), and other studies have reported specific MMP proteins and collagenase-mediated type II collagen degradation products (51, 52). These enzymes are involved in intrinsic chondrocyte-mediated degenerative changes of the cartilage matrix in OA.

In the present study, IGF-induced WJMSCs decreased the ADAMTS1 level. ADAMTS1 has shown in cartilage and the synovium (24) to cleave aggrecan and versican (53). Some studies reported that the expression of ADAMTS1 is significantly elevated in OA cartilage (25, 54–58), however, some studies also indicated a decreased expression in late-stage human OA (59–61).

The TNF family of molecules called RANKL (receptor activator of NF κ B ligand, also known as osteoprotegerin ligand), osteoclast differentiation factor (ODF), TNF related activation-induced cytokine (TRANCE), and TNFSF11 and its receptor RANK (TNFRSF11A) is the main regulator of bone remodeling and activation of osteoclasts (62, 63–66). In this study, IGF-induced WJMSCs also reduced the RANKL level. Production of RANKL activated by T-cells directly controls osteoclastogenesis, bone remodeling, and also associated with autoimmune diseases, cancers, leukemias, asthma, chronic viral infections, and periodontal disease (63). In particular, RANKL is more likely to be the pathogenetic principle that results in the destruction of bone and cartilage in arthritis. RANKL is highly presented in osteoblast/stromal cells, primitive mesenchymal cells surrounding the cartilaginous anlagen, and hypertrophying chondrocytes (62). RANKL mRNA has also been observed in hypertrophic and prehypotrophic chondrocytes at day 15 of embryogenesis and extraskeletal tissues such as the brain, heart, kidneys, skeletal muscles, and skin throughout mouse development (67). RANKL expression can be upregulated by bone-resorbing factors i.e., vitamin D3, glucocorticoids, IL1, IL6, IL11, IL17, TNF α , PGE $_2$, and PTH (68, 62, 65).

These findings are supported by several studies. A study done by Ahmed *et al.* showed that both rat model and co-culture between MSCs and cartilage chips involve MMP-13 and tissue inhibitor of MMP1 and MMP2 as factors in hypertrophy (69). Moreover, some studies also reported enhanced chondrogenesis of MSCs co-cultured with chondrocytes, which shows higher cartilage-specific marker expression in the co-culture compared with monocultures as well as reduced expression of hypertrophic markers such as MMP3 (70, 71). IGF1-WJMSCs is, therefore, promising for use as medicine in the treatment of OA.

Conclusion

The IGF1-induced WJMSCs increased expression of COL2 and SOX9 compared to controls, which indicates IGF1-WJMSCs are capable of enhancing chondrogenesis

and can be further used in OA treatment. Validation of IGF1-WJMSCs in animal models should eventually follow as further study.

Acknowledgment

This study was supported by the Grants-in-Aid from Insinas Riset Pratama 2017 and 2018, the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia. The authors like to thank Hanna Sari W Kusuma, Rahmawati Rahmawati, Risma Laila Qodariah, and Fajar Sukma Perdana from Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung, West Java, Indonesia for their technical assistance.

References

1. Bijlsma JW, Berenbaum F, Lafeber FP. Osteoarthritis: an update with relevance for clinical practice. *Lancet* 2010; 377: 2115-2126.
2. Anderson DD, Chubinskaya S, Guilak F. Post-traumatic osteoarthritis: improved understanding and opportunities for early intervention. *J Orthop Res* 2011; 29: 802-809.
3. Van den Berg WB. Osteoarthritis year 2010 in review: pathomechanisms. *Osteoarthr Cartilage* 2010; 19: 338-341.
4. Aigner T, Gebhard PM, Schmid E, Bau B, Harley V, Poschl E. SOX9 expression does not correlate with type II collagen expression in adult articular chondrocytes. *Matrix Biol* 2003; 22: 363-372.
5. Ashraf A, Cha BH, Kim JS, Ahn J, Han I, Park H, Lee SH. Osteoarthritis and Cartilage. *Osteoarthritis Cartilage* 2016; 24:196-205.
6. Buckwalter JA, Saltzman C, Brown T. The impact of osteoarthritis: implications for research. *Clin Orthop Relat Res* 2004; S6: 15.
7. Aigner T, McKenna L. Molecular pathology and pathobiology of osteoarthritic cartilage. *Cell Mol Life Sci* 2012; 59: 5-18.
8. Johnson K, Zhu S, Tremblay MS. A stem cell-based approach to cartilage repair. *Science* 2012; 336: 717-721.
9. Haleem-Smith H, Calderon R, Song Y. Cartilage oligomeric matrix protein enhances matrix assembly during chondrogenesis of human mesenchymal stem cells. *J Cell Biochem* 2012; 113: 1245-1252.
10. Hass R, Kasper C, Bohm S, Jacobs R. Different populations and sources of human mesenchymal stem cells (MSC): a comparison of adult and neonatal tissue-derived MSC. *Cell Commun Signal* 2011; 9: 12-25.
11. Widowati W, Laura W, Bachtar I. Effect of oxygen tension towards proliferation and characteristics of wharton's jelly-derived mesenchymal stem cells. *BGM* 2014; 7: 1-8.
12. Kim HJ, Im GI. Chondrogenic differentiation of adipose tissue-derived mesenchymal stem cells: Greater doses of growth factor are necessary. *J Orthop Res* 2009; 27: 612-619.
13. Guenther HL, Guenther HE, Froesch ER, Fleisch H. Effect of insulin-like growth factor on collagen and glycosaminoglycan synthesis by rabbit articular chondrocytes in culture. *Experientia* 1982; 38: 979-981.
14. McQuillan DJ, Handley CJ, Campbell MA, Bolis S, Milway VE, Herington AC. Stimulation of proteoglycan biosynthesis by serum and insulin-like growth factor-I in cultured bovine articular cartilage. *Biochem J* 1986; 240: 423-430.
15. Schoenle E, Zapf J, Humbel RE, Froesch ER. Insulin-like growth factor I stimulates growth in hypophysectomized rats. *Nature* 1982; 296: 252-253.

16. Trippel SB, Corvol MT, Dumontier MF, Rappaport R, Hung HH, Mankin HJ. Effect of somatomedin-C/insulin-like growth factor I and growth hormone on cultured growth plate and articular chondrocytes. *Pediatr Res* 1989; 25: 76-82.
17. Bhaumick B, Bala RM. Differential effects of insulin-like growth factors I and II on growth, differentiation and glucoregulation in differentiating chondrocyte cells in culture. *Acta Endocrinol* 1991; 125: 201-211.
18. Bhaumick B. Insulin-like growth factor (IGF) binding proteins and insulin-like growth factor secretion by cultured chondrocyte cells: Identification, characterization and ontogeny during cell differentiation. *Regul Pept* 1993; 48: 113-122.
19. Madry H, Kaul G, Gucchiarini M, Stein U. Enhanced repair of articular cartilage defects *in vivo* by transplanted chondrocytes overexpressing insulin-like growth factor I (IGF-I). *Gene Ther* 2005; 12: 1171-1179.
20. Widowati W, Wijaya L, Murti H, Widyastuti H, Agustina D, Laksmiawati DR, et al. Conditioned medium from normoxia (WJMSCs-norCM) and hypoxia-treated WJMSCs (WJMSCs-hypoCM) in inhibiting cancer cell proliferation. *Biomarkers and Genomic Medicine* 2015; 7: 8-17.
21. Tsuchiya K, Chen G, Ushida T, Matsuno T, Tateishi T. The effect of coculture of chondrocytes with mesenchymal stem cells on their cartilaginous phenotype *in vitro*. *Materials Science and Engineering: C* 2004; 24: 391-396.
22. Ng LJ, Wheatley S, Muscat GE, Conway-Campbell J, Bowles J, Wright E. SOX9 binds DNA, activates transcription, and coexpresses with type II collagen during chondrogenesis in the mouse. *Dev Biol* 1997; 183: 108e21.
23. Zhao Q, Eberspaecher H, Lefebvre V, deCrombrughe B. Parallel expression of Sox9 and Col2a1 in cells undergoing chondrogenesis. *Dev Dyn* 1997; 209: 377e86.
24. Kelwick R, Desantis I, Wheeler GN, Edwards DR. The ADAMTS (A Disintegrin and Metalloproteinase with Thrombospondin motifs) family. *Genome Biol* 2015; 16: 113-128.
25. Gardiner MD, Vincent TL, Driscoll C, Burleigh A, Bou-Gharios G, Saklatvala J. Transcriptional analysis of microdissected articular cartilage in post-traumatic murine osteoarthritis. *Osteoarthritis and Cartilage* 2015; 23: 616-628.
26. Seo S, Na K. Mesenchymal stem cell-based tissue engineering for chondrogenesis. *J Biomed Biotechnol* 2011: 1-8.
27. Can A, Karahuseynoglu S. Concise review: Human umbilical cord stroma with regard to the source of fetus – derived stem cells. *Stem Cells* 2007; 25: 2886-2895.
28. Sekiya I, Tsuji K, Koopman P, Watanabe H, Yamada Y, Shinomiya K, et al. SOX9 enhances aggrecan gene promoter/enhancer activity and is up-regulated by retinoic acid in a cartilage-derived cell line, TC6. *J Biol Chem* 2000; 275: 10738-10744.
29. Zhang P, Jimenez SA, Stokes DG. Regulation of human COL9A1 gene expression. Activation of the proximal promoter region by SOX9. *J Biol Chem* 2003; 278: 117-123.
30. Bridgewater LC, Lefebvre V, de Crombrughe B. Chondrocyte-specific enhancer elements in the Col11a2 gene resemble the Col2a1 tissue-specific enhancer. *J Biol Chem* 1998; 273: 14998-15006.
31. Kou I, Ikegawa S. SOX9-dependent and -independent transcriptional regulation of human cartilage link protein. *J Biol Chem* 2004; 279: 50942-50948.
32. Wagner T, Wirth J, Meyer J, Zabel B, Held M, Zimmer J, et al. Autosomal sex reversal and campomelic dysplasia are caused by mutations in and around the SRY-related gene SOX9. *Cell* 1994; 79: 1111-1120.
33. Li Y, Tew SR, Russell AM, Gonzalez K, Hardingham TE, Hawkins RE. Transduction of human articular chondrocytes with adenoviral, retroviral and lentiviral vectors and the effects of enhanced expression of SOX9. *Tissue Eng* 2004; 10: 575-584.
34. Tew SR, Li Y, Pothacharoen P, Tweats LM, Hawkins RE, Hardingham TE. Retroviral transduction with SOX9 enhances re-expression of the chondrocyte phenotype in passaged osteoarthritic human articular chondrocytes. *Osteoarthritis Cartilage* 2005; 13: 80-89.
35. Lefebvre V, Huang W, Harley VR, Goodfellow PN, de Crombrughe B. SOX9 is a potent activator of the chondrocyte-specific enhancer of the pro alpha1 (II) collagen gene. *Mol Cell Biol* 1997; 17: 2336-2346.
36. Catterall JB, Carrere S, Koshy PJ. Synergistic induction of matrix metalloproteinase 1 by interleukin-1alpha and oncostatin M in human chondrocytes involves signal transducer and activator of transcription and activator protein 1 transcription factors via a novel mechanism. *Arthritis Rheum* 2001; 44: 2296-2310.
37. Stremme S, Duerr S, Bau B, Schmid E, Aigner T. MMP-8 is only a minor gene product of human adult articular chondrocytes of the knee. *Clin Exp Rheumatol* 2003; 21: 205-209.
38. Neuhold LA, Killar L, Zhao W. Postnatal expression in hyaline cartilage of constitutively active human collagenase-3 (MMP-13) induces osteoarthritis in mice. *J Clin Invest* 2001; 107: 35-44.
39. Maiotti M, Monteleone G, Tarantino U, Fasciglioni EG, Marini S, Coletta M. Correlation between osteoarthritic cartilage damage and levels of proteinases and proteinase inhibitors in synovial fluid from the knee joint. *Arthroscopy* 2000; 16: 522-526.
40. Martel-Pelletier J, McCollum R, Fujimoto N, Obata K, Cloutier JM, Pelletier JP. Excess of metalloproteinases over tissue inhibitor of metalloproteinase may contribute to cartilage degradation in osteoarthritis and rheumatoid arthritis. *Lab Invest* 1994; 70: 807-815.
41. Franchimont P, Bassler C. Effects of hormones and local growth factors on articular chondrocyte metabolism. *J Rheumatol* 1991; 18 (Suppl 27): 68-70.
42. Sandell LJ, Dudek EJ. Insulin-like growth factor I stimulates type II collagen gene expression in cultured chondrocytes. *Trans Orthop Res Soc* 1988; 35: 300.
43. Tesch GH, Handley CJ, Cornell HJ, Herington AC. Effects of free and bound insulin-like growth factors on proteoglycan metabolism in articular cartilage explants. *J Orthop Res* 1992; 10: 14-22.
44. Kim HJ, Im GL. Chondrogenic Differentiation of Adipose Tissue-Derived Mesenchymal Stem Cells: Greater Doses of Growth Factor are Necessary. *J Orthop Res* 2009; 27: 612-619.
45. Dean PW, Nelson JK, Schumacher J. Effects of age and prosthesis material on *in vitro* cartilage retention of laryngoplasty prostheses in horses. *American Journal of Veterinary Research* 1990; 51: 114-117.
46. Su S, Grover J, Roughley PJ, DiBattista JA, Martel-Pelletier J, Pelletier JP, et al. Expression of the tissue inhibitor of metalloproteinases (TIMP) gene family in normal and osteoarthritic joints. *Rheumatology International* 1999; 18: 183-191.
47. Clark IM, Powell LK, Ramsey S, Hazleman BL, Cawston TE. The measurement of collagenase, tissue inhibitor of metalloproteinases (timp), and collagenase—timp complex in synovial fluids from patients with osteoarthritis and

- rheumatoid arthritis. *Arthritis and Rheumatology* 1993; 36: 372-379.
48. Cawston TE. Proteinases and inhibitors. *British medical bulletin* 1995; 51: 385-401.
49. Freemont AJ, Hampson V, Tilman R, Goupille P, Taiwo Y, Hoyland JA. Gene expression of matrix metalloproteinases 1, 3, and 9 by chondrocytes in osteoarthritic human knee articular cartilage is zone and grade specific. *Annals of The Rheumatic Diseases* 1997; 56: 542-548.
50. Shlopov BV, Gumanovskaya ML, Hasty KA. Autocrine regulation of collagenase 3 (matrix metalloproteinase 13) during osteoarthritis. *Arthritis and Rheumatism* 2000; 43: 195-205.
51. Freemont AJ, Byers RJ, Taiwo YO, Hoyland JA. In situ zymographic localisation of type II collagen degrading activity in osteoarthritic human articular cartilage. *Annals of the rheumatic diseases* 1999; 58: 357-365.
52. Billingham RC, Dahlberg L, Lonescu M, Reiner A, Bourne R, Rorabeck C. Enhanced cleavage of type II collagen by collagenases in osteoarthritic articular cartilage. *J Clin Invest* 1997; 99: 1534-1545.
53. Rodríguez-Manzaneque JC, Westling J, Thai SN, Luque A, Knauper V, Murphy G. ADAMTS1 cleaves aggrecan at multiple sites and is differentially inhibited by metalloproteinase inhibitors. *Biochem Biophys Res Commun* 2002; 293: 501-508.
54. Geyer M, Grassel S, Straub RH, Schett G, Dinser R, Grifka J. Differential transcriptome analysis of intraarticular lesional vs intact cartilage reveals new candidate genes in osteoarthritis pathophysiology. *Osteoarthritis and Cartilage*, 2009; 17: 328-335.
55. Geyer M, Grassel S, Straub RH, Schett G, Dinser R, Grifka J. Differential transcriptome analysis of intraarticular lesional vs intact cartilage reveals new candidate genes in osteoarthritis pathophysiology. *Osteoarthritis and Cartilage*, 2009; 17: 328-335.
56. Karlsson C, Dehne T, Lindahl A, Brittberg M, Pruss A, Sittertinger M. Genome-wide expression profiling reveals new candidate genes associated with osteoarthritis. *Osteoarthritis and Cartilage*, 2010; 18: 581-592.
57. Ramos YF, den Hollander W, Bovee JV, Bomer N, Van der Breggen R, Lakenberg N. Genes involved in the osteoarthritis process identified through genome wide expression analysis in articular cartilage; the RAAK study. *PLoS One*, 2014; 9: e103056.
58. Wachsmuth L, Bau B, Fan Z, Pecht A, Gerwin N, Aigner T. ADAMTS-1, a gene product of articular chondrocytes *in vivo* and *in vitro*, is downregulated by interleukin 1beta. *J Rheumatol* 2004; 31: 315-320.
59. Davidson RK, Waters JG, Kevorkian L, Darrah C, Cooper A, Donell ST. Expression profiling of metalloproteinases and their inhibitors in synovium and cartilage. *Arthritis Res Ther* 2006; 8: R124-R133.
60. Kevorkian L, Young DA, Darrah C, Donell ST, Shepstone L, Porter S. Expression profiling of metalloproteinases and their inhibitors in cartilage. *Arthritis Rheum* 2004; 50: 131-141.
61. Swingle TE, Waters JG, Davidson RK, Pennington CJ, Puente XS, Darrah C. Degradome expression profiling in human articular cartilage. *Arthritis Res Ther* 2009; 11: R96.
62. Lacey DL, Timms E, Tan HL, Kelley MJ, Dunstan CR, Burgess T. Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. *Cell* 1998; 93: 165-176.
63. Kong YY, Feige U, Sarosi I, Bolon B, Tafuri A, Morony S. Activated T cells regulate bone loss and joint destruction in adjuvant arthritis through osteoprotegerin ligand. *Nature* 1999; 402: 304-309.
64. Simonet WS, Lacey DL, Dunstan CR, Kelley M, Chang MS, Luthy R. Osteoprotegerin: a novel secreted protein involved in the regulation of bone density. *Cell* 1997; 89: 309-319.
65. Yasuda H, Shima N, Nakagawa N, Yamaguchi K, Kinosaki M, Mochizuki S. Osteoclast differentiation factor is a ligand for osteoprotegerin/osteoclastogenesis-inhibitory factor and is identical to TRANCE/RANKL. *Proc Natl Acad Sci USA* 1998; 95: 3597-3602.
66. Wong BR, Rho J, Arron J, Robinson E, Orlinick J, Chao M. TRANCE is a novel ligand of the tumor necrosis factor receptor family that activates c-Jun N-terminal kinase in T cells. *J Biol Chem* 1997; 272: 25190-25194.
67. Kartsogiannis V, Zhou H, Horwood NJ, Thomas RJ, Hards DK, Quinn JM. Localization of RANKL (receptor activator of NFkB ligand) mRNA and protein in skeletal and extracellular tissues. *Bone* 1999; 25: 525-534.
68. Roodman GD. Cell biology of the osteoclast. *Exp Hematol*. 1999; 27: p. 1229-1241.
69. Ahmed N, Dreier R, Göpferich A, Grifka J, Grassel S. Soluble signalling factors derived from differentiated cartilage tissue affect chondrogenic differentiation of rat adult marrow stromal cells. *Cell Physiol Biochem* 2007; 20: 665-678.
70. Bian L, Zhai DY, Mauck RL, Burdick JA. Coculture of human mesenchymal stem cells and articular chondrocytes reduces hypertrophy and enhances functional properties of engineered cartilage. *Tissue Eng A* 2011; 17: 1137-1145.
71. Babur BK, Kabiri M, Klein TJ, Lott WB, Doran MR. The rapid manufacture of uniform composite multicellular-biomaterial micropellets, their assembly into macroscopic organized tissues, and potential applications in cartilage tissue engineering. *PLoS One* 2015; 10: e0122250.

Manuscript ID: IJBMS-1712-6840

Manuscript Title: **Effects of Insulin Like Growth Factor (IGF)-Induced Wharton Jelly Mesenchymal Stem Cells (WJMSCs) toward Chondrogenesis in Osteoarthritis Model**

Authors: Wahyu Widowati, Ervi Afifah, Tjandrawati Mozef, Ferry Sandra, Rizal Rizal, Annisa Amalia, Yukko Arinta, Indra Bachtiar, Harry Murti

Dear Dr **Dr. Wahyu Widowati**

I wish to acknowledge the receipt of the above mentioned manuscript.

Please be sure that the submitted manuscript has not been published previously and will not be submitted elsewhere prior to our decision. Accepted manuscripts cannot be withdrawn, once this decision has been reached.

Open access publishing is not without costs. IJBMS, therefore, levies an article-processing charge of 4.000.000 Rls. for each original and short communication article accepted for publication. Please note that these charges apply only to accepted articles of Iranian authors. All other articles are exempt from these charges.

Our editorial decision will be brought to your attention once the paper has been reviewed.

Regards

Bibi Sedigheh Fazly Bazzaz, Ph.D.

Editor-in-Chief
Iranian Journal of Basic Medical Sciences

<http://ijbms.mums.ac.ir>

Tel: +98-5138823265 Fax: +98-5138823251

E-mail: jbms@mums.ac.ir

Dear Madam/Sir

The manuscript ID No.6840 has been checked for plagiarism. The evaluation shows that your article has some similarity in text with previous published paper(s). We can not accept your manuscript in its current form. Please find the attached report, correct the manuscript and send back it as soon as possible.

Manuscript ID: IJBMS-1712-6840

Manuscript Title: **Effects of Insulin Like Growth Factor (IGF)-Induced Wharton Jelly Mesenchymal Stem Cells (WJMSCs) toward Chondrogenesis in Osteoarthritis Model**

Authors: Wahyu Widowati, Ervi Afifah, Tjandrawati Mozef, Ferry Sandra, Rizal Rizal, Annisa Amalia, Yukko Arinta, Indra Bachtiar, Harry Murti

Dear Dr. **Dr. Wahyu Widowati**

Your manuscript entitled: **Effects of Insulin Like Growth Factor (IGF)-Induced Wharton Jelly Mesenchymal Stem Cells (WJMSCs) toward Chondrogenesis in Osteoarthritis Model**

which you submitted to The Iranian Journal of Basic Medical Sciences, has been reviewed. The reviewer(s) would like to see some revisions made to your manuscript before publication. Therefore, I invite you to respond to the reviewer(s)' comments and revise your manuscript.

When you revise your manuscript please highlight the changes you make in the manuscript by using bold and underline text.

A revised version of your manuscript that takes into account the comments of the reviewer(s) will be reconsidered for publication. Please note that revising your manuscript does not guarantee eventual acceptance, and that your revision will be subject to re-review before a decision is rendered.

Because we are trying to facilitate timely publication of manuscripts submitted to The Iranian Journal of Basic Medical Sciences, your revised manuscript should be uploaded within one month. If it is not possible for you to submit your revision in a reasonable amount of time, we may have to consider your paper as a new submission.

To see reviewers' comments please log in to the journal's website.

When submitting your revised manuscript, please respond to each of the comments made by the reviewer(s) in a separate file.

IMPORTANT:

1) If this study is part of student's thesis, please add this sentence" The results described in this paper were part of student thesis" to the Acknowledge part.

Once again, thank you for submitting your manuscript to The Iranian Journal of Basic Medical Sciences, and I look forward to receiving your revision.

Sincerely,
B.S. Fazly Bazzaz, Ph.D.

Editor-in-Chief
Iranian Journal of Basic Medical Sciences

Editor's Comments to Author:

Please also make any or all of the following revisions as needed:

- 1- Separate discussion from results.
- 2- Abstracts should be structured and limited to 250 words.
- 3- References should conform to the journal's format.
- 4- Conclusions are required.
- 5- Declare financial source in acknowledgements.
- 6-Running title is included and has 50 characters at the most.
- 7-A list of 5-8 keywords, chosen from the Medical Subject Headings (MeSH) list, is to be provided directly below the abstract.

Manuscript ID: IJBMS-1712-6840 (R1)

Manuscript Title: **Effects of Insulin Like Growth Factor (IGF)-Induced Wharton Jelly Mesenchymal Stem Cells (WJMSCs) toward Chondrogenesis in Osteoarthritis Model**

Authors: Wahyu Widowati, Ervi Afifah, Tjandrawati Mozef, Ferry Sandra, Rizal Rizal, Annisa Amalia, Yukko Arinta, Indra Bachtiar, Harry Murti

Date: 2017-12-05

Dear **Dr. Wahyu Widowati**

Thank you for submitting the revised file of your manuscript to the **Iranian Journal of Basic Medical Sciences**

The Editorial Office will proceed on your manuscript and inform you in the earliest time.

If there is anything else, please do not hesitate to contact us.

Truly yours,

Executive Managing Director of **Iranian Journal of Basic Medical Sciences**

Manuscript ID: IJBMS-1712-6840 (R1)

Authors: Wahyu Widowati, Ervi Afifah, Tjandrawati Mozef, Ferry Sandra, Rizal Rizal, Annisa Amalia, Yukko Arinta, Indra Bachtiar, Harry Murti

Dear **Dr. Wahyu Widowati**

I am pleased to inform you that your paper entitled:

Manuscript Title: **Effects of Insulin Like Growth Factor (IGF)-Induced Wharton Jelly Mesenchymal Stem Cells (WJMSCs) toward Chondrogenesis in Osteoarthritis Model**

has been accepted for publication in the Iranian Journal of Basic Medical Sciences. Thank you for submitting your work to the Iranian Journal of Basic Medical Sciences.

With kind regards,

Bibi Sedigheh Fazly Bazzaz, Ph.D.

Editor-in-Chief

Iranian Journal of Basic Medical Sciences

<http://ijbms.mums.ac.ir>

Tel: +98-5138823265 Fax: +98-5138823251

E-mail: jbms@mums.ac.ir

Should be:	
Italic	<i>In vitro- In vivo</i>
Only the first letter of the title should be capitalized. Other letters should be typed in lower case. Also within the tables and figures. DO NOT USED ABBREVIATIONS IN TITLES. No frame or background or extra lines.	Title of Figures and Tables
Italic without period	<i>et al</i>
Italic	<i>Post hoc- ad hoc –ad libitum</i>
min	Minute
hr	hour
sec	second
g	gram
Italic. First time mentioned in the manuscript should be typed complete. For the second time etc, the first letter or the first letter of the first word should be capitalized, then one space and then the second word should be typed with lowercase letter -. e.g.: <i>Escherichia coli</i> <i>E. coli</i>	names of microbes, bacteria, plants
Capitalized, no period	IP-IV-IM
Capitalized and complete.	The word Figure-Table
Italic, Capitalized	<i>P</i> -value
Vancouver Style	References
Should be structures as:	Abstract format
objectives, Material&Methods, results, conclusion	
-Running title is included and has 50 characters at the most. -Discussion is separated from results. -Introduction, Conclusion, Materials and Methods are included. -Tables and figures are integrated into the manuscript and not submitted separately	manuscript structure
Cannot be MSc student. Necessarily a Professor	Corresponding author

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau *PEER REVIEW*

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Effects of insulin-like growth factor-induced Wharton jelly mesenchymal stem cells toward chondrogenesis in an osteoarthritis model

Jumlah Penulis : 9 orang

Nama-nama Penulis : **Wahyu Widowati**, Ervi Afifah, Tjandrawati Mozef, Ferry Sandra, Rizal Rizal, Annisa Amalia, Yukko Arinta, Indra Bachtiar, Harry Murti

Status Penulis : Penulis Pertama / ~~Penulis ke-....~~ / Penulis Korespondensi **)

Identitas Jurnal Ilmiah : a. Nama jurnal : Iranian Journal of Basic Medical Sciences
b. Nomor ISSN : 2008-3874
c. Vol., No., Bulan, Tahun : Vol.21., No.7, July 2018
d. Penerbit : Mashhad University of Medical Sciences
e. DOI Artikel (jika ada) : 10.22038/IJBMS.2018.28205.6840
f. Alamat Web Jurnal : <https://pubmed.ncbi.nlm.nih.gov/30140415/>
g. Terindeks di : Scopus Q2, SJR 0.611

Kategori Publikasi Jurnal Ilmiah : ☒ Jurnal Ilmiah Internasional / Internasional Bereputasi **)
(beri tanda \checkmark yang dipilih)

☐ Jurnal Ilmiah Nasional Terakreditasi

☐ Jurnal Ilmiah Nasional / Nasional terindeks ***)

HASIL PENILAIAN (Peer Review) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi <input checked="" type="checkbox"/>	Nasional Terakreditasi <input type="checkbox"/>	Nasional ***) <input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,7
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			11,5
c.	Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12			11,7
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			11,8
	Total	40			38,7

Catatan Penilaian ARTIKEL oleh Reviewer :


- a. Kelengkapan dan kesesuaian unsur.....
Unsur lengkap, hasil penelitian sesuai dengan judul penelitian, mengkuh kardah penelitian
- b. Ruang lingkup & kedalaman pembahasan.....
Data sangat memadai meliputi induksi WJMSCs menggunakan lGF untuk kondrogenesis pada sel model Osteoarthritis
- c. Kecukupan & kemutakhiran data serta metodologi.....
Sumber pustaka mutakhir, metode co-culture WJMSCs yang diinduksi lGF dan sel kondroit menunjukkan lGF meningkatkan ekspresi COL-2, SOX-9 lebih baik dibanding WJMSCs tanpa induksi lGF
- d. Kelengkapan unsur dan kualitas penerbit.....
Jurnal IJBMS terindeks Scopus Q2 SJR 0,611. Penerbit Mashad University of Medical Sciences
- e. Indikasi plagiasi

Similarity index sebesar : 16%, tidak terdapat indikasi plagiarisme atau self-plagiarism.

f. Kesesuaian bidang ilmu

Paper bidang biomedik sangat sesuai dengan bidang ilmu penulis.

REVIEWER 1



(Prof. Dr. Chrismis Novalinda Ginting, M.Kes)
NIK : 0115127801
UNIVERSITAS PRIMA INDONESIA

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau PEER REVIEW

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Effects of insulin-like growth factor-induced Wharton jelly mesenchymal stem cells toward chondrogenesis in an osteoarthritis model

Jumlah Penulis : 9 orang

Nama-nama Penulis : **Wahyu Widowati**, Ervi Afifah, Tjandrawati Mozef, Ferry Sandra , Rizal Rizal , Annisa Amalia, Yukko Arinta , Indra Bachtiar, Harry Murti

Status Penulis : Penulis Pertama ~~/ Penulis ke~~ / Penulis Korespondensi **)

Identitas Jurnal Ilmiah : a. Nama jurnal : Iranian Journal of Basic Medical Sciences
b. Nomor ISSN : 2008-3874
c. Vol., No., Bulan, Tahun : Vol.21 ., No.7, July 2018
d. Penerbit : Mashhad University of Medical Sciences
e. DOI Artikel (jika ada) : 10.22038/IJBMS.2018.28205.6840
f. Alamat Web Jurnal : <https://pubmed.ncbi.nlm.nih.gov/30140415/>
g. Terindeks di : Scopus Q2, SJR 0.611

Kategori Publikasi Jurnal Ilmiah: ☒ Jurnal Ilmiah Internasional / Internasional Bereputasi **)

(beri tanda ✓ yang dipilih)

☐ Jurnal Ilmiah Nasional Terakreditasi

☐ Jurnal Ilmiah Nasional / Nasional terindeks ***)

HASIL PENILAIAN (Peer Review) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi <input checked="" type="checkbox"/>	Nasional Terakreditasi <input type="checkbox"/>	Nasional ***) <input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,4
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			11,7
c.	Kecukupan dan kemutakhiran data/ informasi dan metodologi (30%)	12			11,5
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			11,8
Total 100%		40			38,4

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur.....
Jurnal ini sudah cukup lengkap dan terdapat kesesuaian antara unsur dan isinya
- b. Ruang lingkup & kedalaman pembahasan
Ruang lingkup bahasan sudah memadai dan terdapat kedalaman analisa data dan juga pembahasannya.
- c. Kecukupan & kemutakhiran data serta metodologi.....
Secara umum metodologinya masih terbatas dan belum sepenuhnya
- d. Kelengkapan unsur dan kualitas penerbit
Kualitas penerbit sudah memadai dan memenuhi standar dan kualitas jurnal sudah masuk kategori Internasional Bereputasi yang baik
- e. Indikasi plagiasi

Setara umum belum terlihat adanya unsur plagiasi

f. Kesesuaian bidang ilmu

Jurnal ini sudah sesuai dengan bidang ilmu yang diteliti

REVIEWER 2



(Prof. Dr. Ermi Girsang, M. Kes)

NIK : 0117057501

UNIVERSITAS PRIMA INDONESIA

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau *PEER REVIEW*

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Effects of insulin-like growth factor-induced Wharton jelly mesenchymal stem cells toward chondrogenesis in an osteoarthritis model

Jumlah Penulis : 9 orang

Nama-nama Penulis : **Wahyu Widowati**, Ervi Afifah, Tjandrawati Mozef, Ferry Sandra , Rizal Rizal , Annisa Amalia, Yukko Arinta , Indra Bachtiar, Harry Murti

Status Penulis : Penulis Pertama / ~~Penulis ke-....~~ / Penulis Korespondensi **)

Identitas Jurnal Ilmiah : a. Nama jurnal : Iranian Journal of Basic Medical Sciences
b. Nomor ISSN : 2008-3874
c. Vol., No., Bulan, Tahun : Vol.21 ., No.7, July 2018
d. Penerbit : Mashhad University of Medical Sciences
e. DOI Artikel (jika ada) : 10.22038/IJBMS.2018.28205.6840
f. Alamat Web Jurnal : <https://pubmed.ncbi.nlm.nih.gov/30140415/>
g. Terindeks di : Scopus Q2, SJR 0.611

Kategori Publikasi Jurnal Ilmiah: ☒ Jurnal Ilmiah Internasional / Internasional Bereputasi **)
(beri tanda ✓ yang dipilih)

☐ Jurnal Ilmiah Nasional Terakreditasi

☐ Jurnal Ilmiah Nasional / Nasional terindeks ***)

HASIL PENILAIAN (Peer Review) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi <input checked="" type="checkbox"/>	Nasional Terakreditasi <input type="checkbox"/>	Nasional ***) <input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,55
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			11,6
c.	Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12			11,6
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			11,8
	Total	40			38,55

Catatan Penilaian ARTIKEL oleh Reviewer :

a. Kelengkapan dan kesesuaian unsur.....

Unsur lengkap, hasil penelitian sesuai dengan judul penelitian, mengikuti kaidah penelitian

Jurnal ini sudah cukup lengkap dan terdapat kesesuaian antara unsur dan bingkai

b. Ruang lingkup & kedalaman pembahasan

Data sangat memadai multiple induksi WJMSCs menggunakan LGF untuk kondrogenesis pada sel osteoarthritis

Ruang lingkup bahasan sudah memadai dan terdapat kedalaman analisa data dan juga pembahasannya

c. Kecukupan & kemutakhiran data serta metodologi

Sumber pustaka mutakhir, metode *in-vitro*

WJMSCs yang diinduksi IGF dan sel kondrosit menunjukkan IGF meningkatkan ekspresi COL 2 & COX 9 lebih baik dibanding WJMSCs tanpa induksi IGF

Secara umum metodologi masih terbatas dan belum sepenuhnya

d. Kelengkapan unsur dan kualitas penerbit

Jurnal ITBMS terindeks Scopus Q2 SJR 0,611. Penerbit Marshall University of Medical Sciences

Kualitas penerbit sudah memadai dan memenuhi kaidah dan kualitas jurnal sudah masuk kategori internasional bereputasi yang baik

e. Indikasi plagiasi

Similaritas index sebesar 16% , tidak terdapat indikator plagiarisme atau self plagiarisme

Secara umum belum terlihat adanya unsur plagiasi

f. Kesesuaian bidang ilmu

Paper bidang biomedik sangat sesuai dengan bidang Ilmu penulis

Jurnal ini sudah sesuai dengan bidang Ilmu yang ditekuni

Medan,
Reviewer 2

(Prof. Dr. Ermi Girsang, M.Kes)

NIK : 0117057501

UNIVERSITAS PRIMA INDONESIA

Medan,
Reviewer 1

(Prof. Dr. Chrismis Novalinda Ginting, M.Kes)

NIK : 0115127801

UNIVERSITAS PRIMA INDONESIA