



Printed: 2016-12-01
E-ISSN: 2278-9978

Volume 5 | Issue 1 | January-June 2016

Journal of Reports in Pharmaceutical Sciences

An Official Publication of Karnataka University of Medical Sciences

www.kumscjournal.com



Editorial Board

Chairman

Reza Khodarahmi, Department of Pharmacognosy & Biotechnology, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, PO box: 67145-1673, Iran.

Editor in Chief

Gholamreza Bahrani, Department of Pharmacology and Toxicology, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, PO box: 67145-1673, Iran.

Executive Manager

Mohsen Shahlaei, Department of Medicinal Chemistry, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, PO box: 67145-1673, AND Nanosciences and Technology Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran.

Office Manager

Elaheh Alizadeh, Pharmaceutical Sciences Research Center, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, PO box: 67145-1673, Iran.

Editorial Board

Avat Taherpoor, Department of Organic Chemistry, Chemistry Faculty, Razi University P.O.Box: 67149-67346, Kermanshah, Iran.

Omید Tavallaei, Department of Pharmacognosy & Biotechnology, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, PO box: 67145-1673, Iran.

Katayoun Derakhshandeh, Department of Pharmacology and Pharmaceutics, University of Medical Science, Hamedan, Iran.

Laila Hosseinzadeh, Department of Pharmacology and Toxicology, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, PO box: 67145-1673, Iran.

Yalda Shokoolinia, Department of Pharmacognosy & Biotechnology, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, PO box: 67145-1673, Iran.

Ahmad Mohammadi-Farani, PharmD, PhD Assistant Professor of pharmacology, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, PO box: 67145-1673, Iran.

Amir Kiani, Department of Pharmacology and Toxicology, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, PO box: 67145-1673, Iran.

Lotfollah Saghie, Department of Medicinal Chemistry - Faculty of Pharmacy and Pharmaceutical sciences, Isfahan University of Medical Sciences, Isfahan, Iran.

Afshin Fassih, Medicinal Chemistry department, School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Sciences, Isfahan, Iran.

Mojtaba Amani, Department of Basic Sciences, Faculty of Medicine, Ardabil University of Medical Sciences, Ardabil, Iran.

Alireza Allahbadi, Department of Medicinal Chemistry, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, PO box: 67145-1673, Iran.

Ghobad Mohammadi, Department of Pharmaceutics, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, PO box: 67145-1673, Iran.

Valiollah Hajhashemi, Department of Pharmacology, School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Sciences, Isfahan, Iran.

Mohammad Barzegar-Jalali, Department of Pharmaceutics, School of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran.

Alireza Foroumadi, Faculty of Pharmacy and Pharmaceutical Sciences Research Center, Tehran University of Medical Sciences, Tehran, Iran.

All Pourmotabbed, Department of physiology, Kermanshah University of Medical Sciences, Kermanshah 67145-1673, Iran.

Farhad Ahmadi, Department of Medicinal Chemistry, Faculty of Pharmacy, School of Pharmacy, Iran University of Medical Sciences, Iran.

Hadi Adibi, Department of Medicinal Chemistry, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, PO box: 67145-1673, Iran.

All Mostafaei, Department of Immunology and Biology Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran.

Ramin Miri, Faculty of Pharmacy, Shiraz University of Medical Sciences, PO Box: 71145-1149, Shiraz, and Medicinal & Natural Products Chemistry Research Center, Shiraz University of Medical Sciences, PO Box: 71345-3388, Shiraz, Iran.

Seyed Ebrahim Sajjadi Jazi, Professor of Pharmacognosy, School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Sciences, Isfahan, Iran.

Mohsen Shahlaei, Department of Medicinal Chemistry, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, PO box: 67145-1673, AND Nanosciences and Technology Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran.

Shahla Mirzaei, Department of Pharmaceutics, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, PO box: 67145-1673, Iran.

Eliham Arkan, Department of Medicinal Nanobiotechnology, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, PO box: 67145-1673, Iran, AND Nanosciences and Technology Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran.

Mohammad Hosein Farzaei, Department of Traditional Medicine, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, PO box: 67145-1673, Iran.

Abolghasem Jouyban, Department of Medicinal Chemistry, Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran.

Seyed Ahmad Emami, Department of Pharmacognosy, School of Pharmacy, Mashhad University of Medical Sciences, AND Medical Toxicology Research Center, Mashhad University of Medical Sciences, Mashhad, Iran.

Abbas Delazar, Faculty of Pharmacy and Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

Sako Mirzaei, Islamic Azad University Of Sanandaj, Department of Biochemistry, Sanandaj, Iran.

International Editorial Board

Keykavous Parang, Department of Biomedical and Pharmaceutical Sciences, University of Rhode Island, Kingston, Rhode Island, USA.

All Nokhodchi, Chemistry and Drug Delivery Group, Medway School of Pharmacy, University of Kent, UK.

Gert Storm, Department of Pharmaceutics, Utrecht Institute of Pharmaceutical Sciences, Utrecht University, Utrecht, The Netherlands.

Elena Papaleo, Structural Biology and NMR Laboratory, Department of Biology, University of Copenhagen, Denmark.

Boris Ivanovich Kurganov, A.N.Bach Institute of Biochemistry, Russian Academy of Sciences, Leninsky prospekt 33, Moscow 119071, RUSSIA.

Faris Farasaati, The University of Kansas Medical Center, Department of Medicine, Kansas city, KS, 66160.

Vikas Kumar, National Center for Natural Products Research (NCNPR), School of Pharmacy, University of Mississippi.

Saqain Haider, Postdoctoral Research Associate – Medicinal Chemistry (November 2014 – Currently working) - National Center for Natural Products Research, School of Pharmacy, The University of Mississippi, Oxford-MS, USA.

André M. Marques, Ph.D In Chemistry of Natural Products, Laboratory of Natural Product (PN3), Natural Products Department, Institute of Pharmaceutical Technology - Farmanguinhos (FIOCRUZ), Rio de Janeiro, Brazil. CEP: 21041-250. Tel. +5521-3977-2492.

Tarun Belwal, Post-Doc Fellow College of Biosystems Engineering and Food Science, Zhejiang University, People's Republic of China, Contact No.: +86 14758166086.

Shankar D. Birajdar, Assistant Professor, Department of Basic Science and Humanities, Marathwada Institute of Technology, Aurangbad, (M. S) India. Mobile: +919822736542.

[About the journal](#)[SUBMIT ARTICLE](#)[SUBSCRIBE](#)[POPULAR ARTICLES](#)[JOIN AS REVIEWER](#)[GET EMAIL ALERTS](#)[RECOMMEND](#)

Table of Contents



September-December 2018

Volume 7 | Issue 3

Page Nos. 222-399

Online since Monday, March 25, 2019

Accessed 119 times.

PDF access policy

Journal allows immediate open access to content in HTML + PDF

View issue as eBook

Issue statistics

RSS

[Next Issue](#)

[Previous Issue](#)

[SUBMIT ARTICLE](#)

[SUBSCRIBE](#)

[POPULAR ARTICLES](#)

[JOIN AS REVIEWER](#)

[GET EMAIL ALERTS](#)

[RECOMMEND](#)

[Show all abstracts](#) [Show selected abstracts](#) [Export selected to](#) [Add to my list](#)

RESEARCH ARTICLES

- Chemical composition and antioxidant activity of *Pistacia Atlantica Subsp. Kurdica* from awraman** p. 222
Seyed Mostafa Nachvak, Seyran Hosseini, Amir Nil-Ahmadabadi, Dara Dastan, Mansour Rezaei
[ABSTRACT] HTML Full Text not available [PDF] [Sword Plugin for Repository]^{Beta}
- The effect of anthraquinones fractions from *Morinda Citrifolia* leaves against pathogenic bacteria with antioxidant activities** p. 231
Abdul Mushin M. Shami
[ABSTRACT] HTML Full Text not available [PDF] [Sword Plugin for Repository]^{Beta}
- Effect of combined exposure to chronic high fat diet and arsenic on kidney function in male mice** p. 240
Akram Ahangarpour, Azin Samimi, Leila Zeidooni, Soheila Alboghobeish, Layasadat Khorsandi, Ali Akbar Oroojan
[ABSTRACT] HTML Full Text not available [PDF] [Sword Plugin for Repository]^{Beta}
- Formulation and evaluation of langsung (*Lansium domesticum* Corr.) peel ethanol extracts lotion as anti-mosquito repellent** p. 250
Rama Hiola, Nurul Rizky, Hamsidar Hasan, Nurain Thomas
[ABSTRACT] HTML Full Text not available [PDF] [Sword Plugin for Repository]^{Beta}
- Spasmolytic effects of hydroalcoholic extract of *Melissa Officinalis* on isolated rat ileum** p. 260
Azam Khalaj, Samira Khani
[ABSTRACT] HTML Full Text not available [PDF] [Sword Plugin for Repository]^{Beta}
- Antioxidant and antiaging activities of *Jasminum Sambac* extract, and its compounds** p. 270
Wahyu Widowati, B Widya Janeva, Sri Nadya, Annisa Amalia, Seila Arumwardana, Hanna Sari W. Kusuma, Yukko Arinta
[ABSTRACT] HTML Full Text not available [PDF] [Sword Plugin for Repository]^{Beta}
- Antitumoral effects of digoxin in the liver cancer cell line is mediated through induction of apoptosis and down regulation of HIF-1 expression** p. 286
Atefeh Tahervand, Minoo Mahmoodi, Mehryar Habibi Roudkenar, Amaneh Mohammadi Roushandeh
[ABSTRACT] HTML Full Text not available [PDF] [Sword Plugin for Repository]^{Beta}
- Anti-inflammatory effect of *Ocimum Basilicum* Linn. seeds hydroalcoholic extract and mucilage on acetic acid-induced colitis in rats** p. 295
Fahimeh Saeidi, Sayed-Ebrahim Sajadi, Mohsen Minaian
[ABSTRACT] HTML Full Text not available [PDF] [Sword Plugin for Repository]^{Beta}
- Characterization of the *Urtica Dioica* Leaf extracts and studies its effects on DNA and sperm in rats** p. 306
Yasamin Khudiar Alghanimi, Najem Abdulhussain Najem, Ali Hamood AL-Saadi
[ABSTRACT] HTML Full Text not available [PDF] [Sword Plugin for Repository]^{Beta}
- Acute and sub-chronic toxicity evaluation of aqueous extract of *Phoenix Dactylifera* seeds in wistar rats** p. 319
Sajad Fakhri, Parisa Shokoohinia, Marzieh Marami, Nastaran Ghiasvand, Leila Hosseinzadeh, Yalda Shokoohinia
[ABSTRACT] HTML Full Text not available [PDF] [Sword Plugin for Repository]^{Beta}
- Active fractions of dichloromethane extract of *Artemisia Aucheri* inhibit proliferation of human breast cancer MCF-7 cells via induction of apoptosis** p. 331
Mahdi Mojarab, Alireza Pourasadollah, Hajar Motamed, Farahnaz Ahmadi, Leila Hosseinzadeh, Marziyeh Hajialyani
[ABSTRACT] HTML Full Text not available [PDF] [Sword Plugin for Repository]^{Beta}
- Protective effect of grape seed extract against acrylamide-induced neurotoxicity in *Vitro* and *In Vivo*** p. 344
Leila Mohammadzadeh, Soghra Mehri, Hossein Hosseinzadeh
[ABSTRACT] HTML Full Text not available [PDF] [Sword Plugin for Repository]^{Beta}
- The anti-atherosclerotic effect of *Prosopis Farcta* is associated with the inhibition of VCAM-1 and ICAM-1** p. 357
Gholamreza Bahrami, Leila Hosseinzadeh, Zahra Azadi, Bahareh Mohammadi, Zahra Mahdijan, Marziyeh Hajialyani, Mohammad Hosein Farzaei
[ABSTRACT] HTML Full Text not available [PDF] [Sword Plugin for Repository]^{Beta}
- In vitro* cytotoxic study of benzylothio-triazol-5-haloisatin scaffolds and their evaluation on supernatants activities and levels of MMP-2 and MMP-9 in MCF-7 cell line** p. 368
Mahdi Rahpeyma, Hadi Adibi, Mohsen Shahlaei, Khadje Najafi, Amir Kiani
[ABSTRACT] HTML Full Text not available [PDF] [Sword Plugin for Repository]^{Beta}
- Antibacterial and antioxidant properties of anthraquinones fractions from *Morinda Citrifolia* fruit** p. 379
Abdul Mushin M. Shami
[ABSTRACT] HTML Full Text not available [PDF] [Sword Plugin for Repository]^{Beta}
- Protective effect hydroalcoholic extract *Allium Jesdianum* Bolss against Ccl4-Induced hepatotoxicity in male mice** p. 389
Heibatullah Kalantari, Esmat Keshavarzi, Parvin Kheradmand, Maedeh Goodarzi, Leila Zeidooni
[ABSTRACT] HTML Full Text not available [PDF] [Sword Plugin for Repository]^{Beta}

Antioxidant and Antiaging Activities of *Jasminum Sambac* Extract, and its Compounds

Wahyu Widowati^{a*}, Widya Janeva Ba^a, Sri Nadya^a, Annisa Amalia^b, Seila Arumwardana^b, Hanna Sari W. Kusuma^b, Yukko Arinta^b

^aMedical Research Center, Faculty of Medicine, Maranatha Christian University, Jl. Prof. Drg. Surya Sumantri No. 65 Bandung 40164, West Java, Indonesia

^bAretha Medika Utama, Biomolecular and Biomedical Research Center, Jl. Babakan Jeruk 2, No. 9, Bandung 40163, West Java, Indonesia

ABSTRACT

Aging is a complex process characterized by a progressive decline in physiological function, followed by dysfunction, and ultimately, death. Increase activity of hyaluronidase, elastase and collagenase, are documented in skin aging. Free radicals can stimulate skin aging through antioxidant system destruction, wrinkle formation, and melanogenesis. Antioxidant and anti-aging agents have been recently developed from herbal plants. In this study, we investigated the antioxidant and anti-aging activities of *Jasminum sambac* extract (JSE). The phytochemical assay was performed with modified Farnsworth method. Antioxidant assays were measured by 2,2-diphenyl-1-picrylhydrazyl (DPPH) scavenger, Ferric Reducing Antioxidant Power (FRAP) and 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS)-reducing activities. Anti-aging properties were measured through inhibitory activities of collagenase, elastase, and hyaluronidase. Phytochemical analysis showed presence of phenols, triterpenoids, and flavonoids in low level, and terpenoids in high level. JSE showed higher DPPH-scavenging activity ($IC_{50}=94.13 \pm 10.54 \mu\text{g/mL}$) than eugenol ($2.28 \pm 0.12 \mu\text{g/mL}$), but lower than hesperidin ($226.34 \pm 4.96 \mu\text{g/mL}$). JSE showed lowest ABTS-activity ($IC_{50}=39.20 \pm 0.45 \mu\text{g/mL}$) compared to hesperidin and eugenol ($IC_{50}=8.10 \pm 0.60$ and $1.56 \pm 0.03 \mu\text{g/mL}$, respectively). The FRAP-reducing activity of JSE, hesperidin, and eugenol showed JSE was the lowest activity at highest concentration (65.46, 178.16 and $402.42 \mu\text{M Fe(II)/}\mu\text{g}$) respectively). JSE showed the lowest anti-collagenase activity ($IC_{50}=339.30 \pm 7.87 \mu\text{g/mL}$), anti-elastase ($IC_{50}=249.94 \pm 16.51 \mu\text{g/mL}$), and anti-hyaluronidase ($IC_{50}=269.26 \pm 90.52 \mu\text{g/mL}$) compared to hesperidin, and eugenol. Overall, JSE has low antioxidant activity compared to hesperidin and eugenol, as well as low anti-collagenase, anti-elastase, and anti-hyaluronidase activities.

Article Type:
Research Article

Article History:

Received: 2018-04-25
Revised: 2018-05-28
Accepted: 2018-06-27
ePublished: 2018-07-03

Keywords:
Jasminum Sambac
Antioxidant
Antiaging
Anti-Collagenase
Anti-Elastase
Anti-Hyaluronidase

*Corresponding Author: Wahyu Widowati, E-mail: wahyu_w60@yahoo.com

Copyright © 2018 by Kermanshah University of Medical Sciences

Introduction

Aging is a complex process characterized by a progressive decline in physiological function, followed by dysfunction, and ultimately death [1]. Aging can be affected by various factors, such as disease, injury, nutrition, exercise, stress, and environmental factors. World Health Organization (WHO) reported that the percentage of the world's population of people over 60 years old will be double from 11% to 22% between 2000 and 2050 [2]. Thus, many recent studies have focused on maintaining a healthy life by postponing aging.

Collagen is one of the major building blocks of the skin that maintains elasticity and strength of the skin. It is the main component of connective tissue, hair and nails [3]. Elastin is a protein found in connective tissue which is responsible for the elasticity of the skin and lungs [4, 5]. Degradation of elastin by intracellular elastase increases with age and/or repeated UV-radiation, leading to skin aging. Hyaluronic acid plays a role in retaining the moisture of the skin, as well as its structure and elasticity. It also facilitates the exchange of nutrients and waste products and is involved in rapid tissue proliferation, regeneration and repair [6]. Collagen, elastin and hyaluronic acid levels decrease during aging process, the increasing enzymes activity including collagenase, elastase and hyaluronidase, resulting loss of strength and flexibility of skin that generates wrinkles [7, 8].

It has been reported that skin aging occurs in presence of cumulative endogenous damage due to free radicals including reactive oxygen species (ROS) [9,10]. ROS are defined as oxygen-containing, highly reactive species. ROS are generated constantly during normal cellular metabolism

which is essential for biological functions. Excessive ROS causes oxidative stress and damage to biological molecules [11, 12]. Previous studies have been investigated that continuous ROS exposure can stimulate skin aging through antioxidant system destruction, wrinkle formation, and melanogenesis [11]. ROS are usually removed from the body through antioxidant defense system [13]. Thus, maintaining antioxidant homeostasis is appropriate strategy to prevent skin aging [14].

Antioxidant properties derived from natural sources have been proposed for aging prevention [14]. Bioactive compounds contained in plants such as isoflavones, anthocyanins, and catechins may have promising antioxidant activity against ROS [15]. It has been reported that *Jasminum sambac* contain iridoidal glycosides [16], linalyl 6-*O*-malonyl- β -D-glucopyranoside, benzyl 6-*O*- β -Dxylopyranosyl- β -D-glucopyranoside (β -primeveroside), 2-phenylethyl β -primeveroside, 2-phenylethyl 6-*O*- α -L-rhamnopyranosyl- β -D-glucopyranoside (β -rutinoside) [17], dotriacontanoic acid, dotriacontanol, oleanolic acid, daucosterol, and hesperidin [18]. The volatile constituents consist of benzyl acetate, indole, E-E- α -farnesene, Z-3-hexenyl benzoate, benzyl alcohol, linalool, and methyl anthranilate [19, 20], eugenol, benzyl benzoate, cis-jasmone, geraniol, farnesol and trace amounts of isohyitol and phytol [19, 20]. Thus, the flowers of *J. sambac* have been utilized as traditional medicines in Asia to treat various diseases including diarrhea, fever, conjunctivitis, abdominal pain, dermatitis, asthma, abscess, breast cancer, uterine bleeding, and toothache. This study we investigated the antioxidant and anti-aging activities of *J. sambac* compared to its compounds namely hesperidin and eugenol (Figure 1).




Fig. 1. Structure of two compounds contained in *J. sambac*; (1) Hesperidin (2) Eugenol

Materials and methods

Extract preparation

Flowers of *J. sambac* were collected from Brebes, Center Java, Indonesia. The plants were identified by herbarium staff, Department of Biology, School of Life Science and Technology, Bandung Institute of Technology, Bandung, West Java, Indonesia. The dried simplicia of flower of *J. sambac* 70 g were extracted with distilled ethanol 70% (500 mL) using maceration technique. Ethanol filtrate was filtered, and wastes were re-macerated in triplicates. Macerates were concentrated using 50 °C rotavapor to obtain pasta form. The yield of *J. sambac* extract (JSE) was 18.28 g, it was stored at -20°C [8, 21, 22, 23, 24]. JSE was used as the experiment. Hesperidin [Chengdu Biopurify Phytochemicals Ltd 14021406] and eugenol [Sigma Aldrich E51791] were used as standard compounds.

Qualitative phytochemical screening assay:

Content of *J. sambac* extract was evaluated by phytochemical screening assay using modified Farnsworth method. Phytochemical screening aim to identify qualitatively presence of phenols, steroids/triterpenoids, saponins, tannins, terpenoids, flavonoids, and alkaloids [8,24,25,26].

High performance liquid chromatography (HPLC)

The analysis of chemical profiling of JSE by HPLC. Semi quantification JSE using the standard hesperidin [Biopurify Phytochemicals Ltd 14021406], and eugenol [Sigma Aldrich E51791]. HPLC analysis used the Hitachi Pump HPLC L-6200, Hitachi L-4000 UV detector and Reverse Phase Column C-18 (Phenosphere ODS-2, Phenomenex, 4.6 mm x 250 mm). Acetonitril 70% [Merck 100030] using to mobile phase (isocratic) with a flow rate of 1.0 mL/min. The samples were dissolved in methanol 70% (1 mg/mL) and filtered through a 0.22 µm syringe and injected 20 µL. UV absorbance was measured at 254 nm [27].

DPPH scavenging activity

Fifty µL samples (0-2000 µg/mL for JSE, 0-2000 µM for compounds) was introduced in 96-well microplate and 200 µL of 0.077 mmol DPPH [Sigma Aldrich D9132] in dimethyl sulfoxide (DMSO) were added. The mixture was shaken vigorously and incubated in a dark room, at room temperature, for 30 min. After that, absorbance was measured at 517 nm using a microplate reader [Multiskan™ GO Microplate Spectrophotometer, Thermo Scientific] was performed. For negative controls, 250 µL DPPH was used. For blank, 250 µL methanol was used

[8,26,28,29]. The DPPH scavenging activity (%) was calculated as follows:

The scavenging activity (%) was then continued to be calculated as median inhibitory concentration (IC₅₀).

$$\text{Scavenging Activity (\%)} = (\text{Ac-As})/\text{Ac} \times 100$$

As: sample absorbance

Ac: negative control absorbance (without sample)

ABTS-reducing activity

The antioxidant capacity JSE, compounds were measured using the 2,2'-Azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) diammonium salt free radical assay [8,26,27,28,29]. ABTS was produced by reacting 14 mM ABTS and equal volume of 4.9 mM potassium persulfate (Merck EM105091) achieved final concentration of 7 mM ABTS in 2.45 mM potassium persulfate. The mixture was incubated in the dark room temperature for 16 h. The ABTS^{•+} solution (Sigma Aldrich A1888) was diluted with 5.5 mM phosphate buffer saline (PBS) pH 7.4 and measured with microplate reader at 745 nm, resulting the absorbance of 0.70 ± 0.02. Briefly 2 µL sample (0-5000 µg/mL for JSE, 0-5000 µM for compounds) was added to 198 µL of ABTS^{•+} solution, incubated for 6 min at 30 °C, and the absorbance was measured at 745 nm (Multiskan™ GO Microplate Spectrophotometer, Thermo Scientific). The percentage inhibition of ABTS radical (%) was expressed by the ratio of the reducing of ABTS^{•+} absorbance in the presence of the test sample relative to the ABTS^{•+} absorbance in the absence of the test sample (negative control). ABTS-reducing activity (%) was then continued to be calculated as median inhibitory concentration (IC₅₀).

$$\text{ABTS-reducing activity \%} = (\text{Ac-As})/\text{Ac} \times 100$$

As: sample absorbance

Ac: negative control absorbance (without sample)

FRAP reducing activity

The ferric reducing antioxidant power assay (FRAP) of each standard solution and sample were measured according to a modified protocol [8,26,28,29]. The FRAP reagent was prepared by adding 2,4,6-tripyridyl-s-triazine (TPTZ) (Sigma 3682-35-7) and ferric chloride hexahydrate (Merck 1.03943.0250), forming the Fe³⁺-TPTZ complex. In 96-well microplate, 7.5 µL of samples (0-2000 µg/mL for JSE, 0-2000 µM for hesperidin, 0-250 µM for eugenol) were mixed with 142.5 µL FRAP reagent then incubated for 30 min at 37 °C. Antioxidant reduced to Fe²⁺-TPTZ at low pH was measured at 595 nm (Multiskan™ GO Microplate Spectrophotometer, Thermo Scientific). The results were expressed in µM Fe (II)/µg extract and compounds [8,26,27,28,29].

Collagenase assay

Collagenase inhibitory activity was measured according to modified method done by Sigma Aldrich Thring et al. (2009) and Widowati et al. (2016, 2017) [7,8,26,30]. Mixture was composed of 10 µL collagenase from *Clostridium histolyticum* [Sigma C8051] (0.01 U/mL in cold aquadest), 60 µL Tricine buffer (50 mM, pH 7.5, containing 10 mM CaCl₂ and 400 mM NaCl), 30 µL sample (0-1000 µg/mL for JSE, 0-1000 µM for compounds), then incubated at 37 °C for 20 min. After incubation, 20 µL of N-[3-(2-Furyl)acryloyl]-leu-gly-Pro-Ala substrate [Sigma F5135] (1 mM in Tricine buffer) was added. Absorbance was measured at 335 nm wavelengths (Multiskan™ GO Microplate Spectrophotometer, Thermo Scientific).

$$\text{Inhibitory activity \%} = (\text{Ac-As})/\text{Ac} \times 100$$

As: sample absorbance

Ac: negative control absorbance (without sample)

Elastase assay

Inhibitory activity of elastase was measured according to modified method by Sigma Aldrich and Thring et al. (2009) and Widowati *et al.* (2016, 2017) [7,8,26,30]. Approximately, 10 μ L sample (0-1000 μ g/mL for JSE, 0-1000 μ M for compounds) was pre-incubated at 25 °C for 15 min with 5 μ L *elastase from porcine pancreas* [Sigma 45124] (0.5 mU/mL in cold aquadest), 125 μ L Tris buffer (100 mM, pH 8). After pre-incubation, mixture was then added with 10 μ L N-Sucanyl-Ala-Ala-Ala-p-Nitroanilide substrate [Sigma 54760] (2 mg/mL in Tris buffer), incubated at 25 °C for 15 min. Absorbance was measured at 410 nm wavelength (Multiskan™ GO Microplate Spectrophotometer, Thermo Scientific, USA).

$$\text{Inhibitory activity \%} = (\text{Ac}-\text{As})/\text{Ac}\times 100$$

As: sample absorbance

Ac: negative control absorbance (without sample)

Hyaluronidase assay

Inhibitory activity of hyaluronidase was measured based on modified method of Sigma Aldrich and Tu and Tawata (2015) and Widowati *et al.* (2016; 2017) [8,26,31]. Briefly 25 μ L samples (0-1000 μ g/mL for JSE, 0-1000 μ M for compounds) was pre-incubated at 37 °C for 10 min with 3 μ L hyaluronidase from bovine testes type I-S [Sigma Aldrich H3506] (0.4 U/mL in 20 mM phosphate buffer, pH 7 containing 77 mM sodium chloride,

and 0.01% bovine serum albumin), and 12 μ L phosphate buffer (300 mM, pH 5.35) at 37 °C for 10 min. After pre-incubation, 10 μ L hyaluronic acid substrate (0.03% in 300 mM phosphate buffer, pH 5.35) [Sigma Aldrich H5542], then incubated at 37 °C for 45 min. Reaction was stopped using 100 μ L acidic albumin acid (24 mM sodium acetate, 79 mM acetate acid, and 0.1% BSA). Mixture was left in room temperature for 10 min, and absorbance was measured at 600 nm wavelengths (Multiskan™ GO Microplate Spectrophotometer, Thermo Scientific).

$$\text{Inhibitory activity \%} = (\text{Ac}-\text{As})/\text{Ac}\times 100$$

As: sample absorbance

Ac: negative control absorbance (without sample)

Results

Phytochemical Analysis

Phytochemical analysis was done to determine content of compounds class in JSE. Compounds of class of JSE can be seen in Table 1. Phytochemical analysis shows the presence of phenols, triterpenoids, and flavonoids in low content, and terpenoids in high content.

HPLC Analysis

HPLC analysis was evaluated to determine the compounds content of JSE. The HPLC standards using hesperidin and eugenol (Figure 2). Hesperidin has a retention time at 1.43 min, and eugenol at 2.15 min. The JSE peak at 1.36 min which was assumed that JSE contains hesperidin and doesn't contain eugenol.

Table 1. The result of qualitative phytochemical assay of JSE

Phytochemical content	Qualitative value of JSE
Phenols	+
Steroids/triterpenoids	-/+
Terpenoids	+++
Saponins	-
Flavonoids	+
Tannins	-
Alkaloids	-




Fig. 2. Chromatogram of JSE, hesperidin, and eugenol from HPLC analysis; (a) JSE (b) Hesperidin (c) Eugenol

DPPH scavenging activity

DPPH free radical scavenging activity can be used to determine antioxidant capacity of JSE and its compounds. This assay based on alcoholic DPPH solution reduction in the presence of a hydrogen-donating antioxidant due to the formation of the non-radical from 2,2-diphenyl-1-picrylhydrazine (DPPH-H). The DPPH samples generally changes

purple colour into a colourless when antioxidant molecules quench DPPH free radicals [8, 28]. The IC₅₀ of scavenging activities of JSE, hesperidin, and eugenol can be seen in Table 2. As shown in Table 2, the IC₅₀ value of JSE was higher (94.13 ± 10.54 µg/mL) than eugenol (2.28 ± 0.12 µg/mL), but lower than hesperidin (226.34 ± 4.96 µg/mL). These results indicate moderate scavenging activity of JSE among samples.

Table 2. The IC₅₀ value of DPPH scavenging activity of JSE, hesperidin, eugenol

Samples	Equation	R ²	IC ₅₀ (µM)	IC ₅₀ (µg/mL)
JSE	y = 0.33x + 18.56	0.94	-	94.13 ± 10.54
Hesperidin	y = 0.07x + 24.05	0.96	370.71 ± 8.13	226.34 ± 4.96
Eugenol	y = 0.84x + 38.25	0.922	13.92 ± 0.74	2.28 ± 0.12

ABTS - reducing activity

ABTS-reducing activity can be used to determine antioxidant capacity of JSE and its compounds. The assay is based on the discoloration of ABTS by antioxidant compounds, the ABTS is generated by reacting a strong oxidizing agent (potassium permanganate/potassium persulfate) with the ABTS salt. The reduction of blue-green ABTS

radical by hydrogen-donating antioxidant compounds [8,26]. The ABTS-reducing activity of JSE, hesperidin, and eugenol can be seen in Table 3. Table 3 shows that the IC₅₀ value of JSE was the highest (39.20 ± 0.45 µg/mL) compared to hesperidin and eugenol (8.10 ± 0.60 and 1.56 ± 0.03 µg/mL, respectively). Based on the results, JSE exhibited lowest ABTS-reducing activity among samples.

Table 3. The IC₅₀ value of ABTS-reducing activity of JSE, hesperidin, eugenol

Samples	Equation	R ²	IC ₅₀ (µM)	IC ₅₀ (µg/mL)
JSE	y = 1.21x + 2.73	0,99	-	39.20 ± 0.45
Hesperidin	y = 3.65x + 1.86	0.99	13.27 ± 0.99	8.10 ± 0.60
Eugenol	y = 4.89x + 3.36	0.97	9.54 ± 0.20	1.56 ± 0.03

Ferric reducing antioxidant power (FRAP) activity

FRAP activity can be used to determine antioxidant capacity of JSE and its compounds. The FRAP activity of JSE, hesperidin, and eugenol can be seen in Table 4. The FRAP activity method is based on the reduction of a ferroin analog, the Fe³⁺ complex of tripyridyltriazine Fe(TPTZ)³⁺ to the intensely blue coloured Fe²⁺ complex

Fe(TPTZ)²⁺ by antioxidant in acidic medium [8,28]. In this study, FRAP activity of JSE, hesperidin and eugenol at the highest concentration are 65.46 ± 1.84, 178.16 ± 2.00 and 402.42 ± 3.46 µM Fe(II)/µg respectively. These results indicate JSE has lowest FRAP activity among samples.

Collagenase assay

Collagen, the major component of the skin, is degraded by the enzyme collagenase. Inhibition of

collagenase activity delays the process of forming pre-collagen fibre and subsequently the wrinkling process [3,8,26]. The inhibitory activity of JSE,

hesperidin, and eugenol can be seen at Figure 3, while the IC_{50} values are shown in Table 5.

Table 4. FRAP activity of JSE, hesperidin, eugenol

Sample	Concentration (μ M)	Concentration (μ g/mL)	FRAP Activity (μ M Fe(II)/ μ g)
JSE	-	2000	65.46 \pm 1.84
		1000	32.11 \pm 0.69
		500	15.22 \pm 0.39
		250	5.94 \pm 0.72
Hesperidin	2000	1221.13	178.16 \pm 2.00
	1000	610.57	103.87 \pm 1.04
	500	305.28	57.81 \pm 0.43
	250	152.64	29.72 \pm 0.77
Eugenol	250	41.05	402.42 \pm 3.46
	125	20.52	249.69 \pm 4.80
	62.50	10.26	155.13 \pm 1.95
	31.25	5.13	82.17 \pm 1.71

Table 5. The IC_{50} value of Collagenase inhibitory activity of JSE, hesperidin, eugenol

Samples	Equation	R^2	IC_{50} (μ M)	IC_{50} (μ g/mL)
JSE	$y = 0.13x + 4.82$	0.95	-	339.30 \pm 7.87
Hesperidin	$y = 0.13x + 9.12$	0.97	326.72 \pm 22.95	198.09 \pm 14.01
Eugenol	$y = 0.03x + 3.39$	0.92	1370.88 \pm 190.53	225.10 \pm 31.20




Fig. 3. Effect of JSE ($\mu\text{g/mL}$), hesperidin, eugenol (μM) on Collagenase inhibitory activity of JSE, hesperidin, eugenol (JSE, hesperidin, and eugenol were diluted in DMSO to reach final concentration of 31.25, 62.50, 125.00, 250.00 ($\mu\text{g/mL}$; μM))

The inhibitory activity of JSE was comparable with hesperidin and eugenol. Figure 3 shows the anti-collagenase activity of JSE, hesperidin, and eugenol. The highest inhibitory activity was hesperidin ($40.72 \pm 2.12\%$). JSE possess moderate anti-collagenase activity among samples ($37.03 \pm$

0.54%). However, as shown in Table 5, hesperidin show the highest anti-collagenase activity with IC_{50} ($198.09 \pm 14.01 \mu\text{g/mL}$), whereas eugenol was moderate ($225.10 \pm 31.20 \mu\text{g/mL}$), and JSE was the lowest ($339.30 \pm 7.87 \mu\text{g/mL}$).




Fig. 4. Effect of JSE (µg/mL), hesperidin, eugenol (µM) on Elastase inhibitory activity of JSE, hesperidin, eugenol

(JSE, hesperidin, and eugenol were diluted in DMSO to reach final concentration of 8.33, 16.67, 33.33, 66.67 (µg/mL; µM))

Elastase assay

Elastin is a protein found in connective tissue which is responsible for the elasticity of the skin and lungs [4,5]. This protein is catalyzed by the enzyme elastase. Degradation of elastin by intracellular elastase increases with age and/or repeated UV-radiation, leading to skin aging [6]. The elastase inhibitory activity of JSE, hesperidin, and eugenol can be seen at Figure 4, while the IC₅₀ values are shown in Table 6.

Figure 4 shows the anti-elastase activity of JSE, hesperidin, and eugenol. The highest activity of JSE at 66.67 µg/mL is 18.60 ± 0.55%, the highest activity of hesperidin, eugenol at 66.67 µM are 39.52 ± 0.71%, 19.83 ± 0.52%, respectively. The IC₅₀ value of samples (Table 6) show eugenol has highest inhibitory activity (IC₅₀= 34.52 ± 1.17 µg/mL), whereas hesperidin was moderate (IC₅₀= 57.46 ± 0.19 µg/mL), and JSE was the lowest (IC₅₀= 249.94 ± 16.51 µg/mL).




Fig. 5. Effect of JSE (µg/mL), hesperidin, eugenol (µM) on Hyaluronidase inhibitory activity of JSE, hesperidin, eugenol

(JSE, hesperidin, and eugenol were diluted in DMSO to reach final concentration of 5.21, 10.42, 20.83, 41.67 (µg/mL; µM))

Hyaluronidase assay

Hyaluronic acid plays a role in retaining the moisture of the skin, as well as its structure and elasticity. Hyaluronic acid levels decrease during aging process [6]. Hyaluronidase inhibitory activity

can be used to determine antiaging capacity of JSE, its compounds. The hyaluronidase inhibitory activity of JSE, hesperidin, and eugenol can be seen in Figure 5, while the IC₅₀ values were shown in Table 7.

Table 6. The IC₅₀ value of Elastase inhibitory activity of JSE, hesperidin, eugenol

Samples	Equation	R ²	IC ₅₀ (µM)	IC ₅₀ (µg/mL)
JSE	y = 0.17x + 8.01	0.94	-	249.94 ± 16.51
Hesperidin	y = 0.39x + 12.92	0.99	94.11 ± 1.63	57.46 ± 0.19
Eugenol	y = 0.20x + 7.11	0.92	210.25 ± 7.12	34.52 ± 1.17

Table 7. The IC₅₀ value of Hyaluronidase inhibitory activity of JSE, hesperidin, eugenol

Samples	Equation	R ²	IC ₅₀ (µM)	IC ₅₀ (µg/mL)
JSE	y = 0.23x - 11.93	0.93	-	269.26 ± 90.52
Hesperidin	y = 0.83x + 32.59	0.90	21.08 ± 0.05	12.87 ± 0.03
Eugenol	y = 0.26x + 6.19	0.93	168.55 ± 19.32	27.67 ± 3.17

Figure 5 shows the anti-hyaluronidase activity of JSE, hesperidin, and eugenol. The highest activity of JSE at 41.67 µg/mL is hesperidin, and eugenol

at concentration 41.67 µM are 64.36 ± 6.03%, -3.49 ± 2.38% respectively. These results are confirmed by IC₅₀ value of JSE and its compounds, as shown in Table 7. Hesperidin showed lowest

IC₅₀ value (12.87 ± 0.03 µg/mL) which indicate highest inhibitory activity, whereas eugenol is moderate (27.67 ± 3.17 µg/mL), and JSE is the lowest 269.26 ± 90.52 µg/mL.

Discussion

Plants have been widely used to treat various diseases due to its medicinal properties. Therefore, phytochemical analysis on plants is required to observe compounds class that might possess medicinal activities. The present study showed JSE contained phenols, triterpenoids, and flavonoids in low content, and terpenoids in high content. In this study, HPLC analysis showed that JSE contain hesperidin. Previous phytochemical analysis showed bioactive compounds in *J. sambac*, namely iridoidal glycosides [16], and hesperidin [18], based on the HPLC result indicate that JSE doesn't contain eugenol, this result was not in line with previous research that *J. sambac* contained eugenol [20].

Antioxidant is used as antiaging prevention due to its involvement ROS in skin aging process. UV radiation absorbed by the skin promotes overproduction of ROS, as well as oxidative stress. Oxidative damage results in lipid peroxide formation, mitochondrial and DNA damage, and protein and gene modification which alter protein structure and function [32]. Excessive ROS leads to the activation of hyaluronidase, collagenase and elastase, which can further contribute to skin aging [3]. In the present study, JSE showed moderate DPPH-scavenging activity (94.13 ± 10.54 µg/mL), and lowest ABTS (39.20 ± 0.45 µg/mL) and FRAP activity in the highest concentration (65.46 ± 1.84 µM Fe (II)/µg) was lower compared to hesperidin and eugenol. Based on other study, if compound has IC₅₀ value < 200 ppm or < 200 µg/mL it has strong antioxidant activity [33]. Based on standards, that indicates JSE has high antioxidant activity toward DPPH-

scavenging activity and ABTS-reducing activity. Referring to previous study, *J. sambac* possess moderate scavenging effect in the order towards the DPPH radicals (122 µg/mL), nitric oxide (173.94 µg/mL) and hydrogen peroxide (125 µg/mL) when compared to ascorbic acid [34]. Following the results, Shekhar and Prasad (2015) found that *Jasminum* species, including *J. sambac*, showed higher FRAP-activity compared to ascorbic acid [35]. Low activity of JSE in the present study might be due to low of flavonoids detected. It has been reported that flavonoids are the largest group among plants with bioactive properties [36]. Flavonoids decrease the early activation of signaling pathway in response to UV induced injury. The quercetin, polyphenol, flavonoid and flavonol, ellagic acids from natural resources, produces the free radicals scavenging activity, inhibition of elastase activity and matrix metalloproteinases (MMPs) expression, and increase expression of procollagen type I [37,38,39,40].

The low antioxidant activity was indeed associated with low anti-aging activity. Based on previous study, antioxidant obtained from plant extract, inhibit elastase, hyaluronidase, collagen synthesis, lipid peroxidation activity, protein expression of procollagen and prevent MMPs gene induction and due to this fibrillin fibre length elongated to maintain elasticity of skin [41,42]. MMPs are part of a group of transmembrane zinc containing endopeptidases which include collagenases and gelatinases. Collagenases are metalloproteinases capable of cleaving other molecules found within the cell for example collagenase-2 (MMP-8) can cleave aggrecan, elastin, fibronectin, gelatine and laminin as well as collagen [43]. Another proteolytic system involved in the degradation of the extracellular matrix (ECM) is that of serine proteases one of which is elastase. Elastase, a member of the chymotrypsin family of proteases, is responsible primarily for the breakdown of elastin which is vital for giving

elasticity to arteries, lungs, ligaments and skin [44, 45, 46, 47]. In terms of anti-aging, finding inhibitors of elastase enzymes can be useful to prevent loss of skin elasticity. In the present study, JSE showed lowest activity of anti-collagenase, anti-elastase and anti-hyaluronidase compared to hesperidin and eugenol. From our best knowledge, there is no studies regarding anti-aging activity of *J. sambac*. However, the activity of JSE in the present study was very low, which might be due to absence of tannins. Tannins is likely found in the most active extract. Recent study has done by Piwowarski et al. (2011) show those twelve tannin-rich plants possess anti-collagenase and anti-hyaluronidase activity [48].

In this study, hesperidin showed highest activity. Hesperidin is a flavanone glycoside (a subclass of flavonoids) that is found abundantly in citrus fruits. Hesperidin possesses vitamin-like activity that decrease capillary permeability (vitamin P), leakiness, and fragility. It also showed antioxidant, anti-inflammatory, anticarcinogenic, and antiallergic properties [49, 50, 51]. The study done by Wilmsen et al. (2005) showed that hesperidin significantly reduced the level of the DPPH which was comparable to antioxidant trolox [52]. It has been reported that hesperidin significantly inhibits the ROS of the yeast [53]. Hesperidin also inhibits the formation of advanced glycation endproducts (AGEs) that possess potentially harmful effects on biological function, and are associated with aging and many degenerative illnesses [54]. Hesperidin derived from Citrus genus extended the lifespan of yeast at doses of 5 and 10 μM as compared to non-treated yeast.

Eugenol in the present study showed moderate antioxidant and anti-aging activity. However, previous studies showed good antioxidant activity of eugenol. Referring to Widowati et al. (2013), eugenol showed good antioxidant activity (IC_{50} of DPPH= 3.8 $\mu\text{g}/\text{mL}$) [22]. Eugenol had the most

powerful antioxidant activity and radical-scavenging activity compared to butylated hydroxyanisole, butylated hydroxytoluene, α -tocopherol, and trolox [55]. Eugenol inhibit lipid peroxidation by trapping active oxygen species such as O_2^- or hydroxyl radicals, rather than by breaking the free radical chain reaction [56].

Conclusion

Overall, JSE showed low antioxidant activity compared to hesperidin and eugenol, as well as low anti-collagenase, anti-elastase, and anti-hyaluronidase.

Acknowledgments

This research was supported by Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung, Indonesia for a research grant, laboratory facilities, and research methodology. We are thankful to Rismawati Laila Qodariah, Rizal Rizal, Fajar Sukma Perdana, Ni Luh Wisma Ekayanti for their valuable assistance.

Conflict of interest

Authors certify that there is no actual or potential conflict of interest in relation to this article.

References

- [1] Raj K, Chanu SI, Sarkar S. Decoding complexity of aging. *Cell Dev Biol*. 2012;1:e117.
- [2] World Health Organization. Interesting facts about ageing. <http://www.who.int/ageing/about/facts/en/> (accessed march 2012). 2012
- [3] Mukherjee PK, Maity N, Nema NK, Sarkarm BK. Bioactive compounds from natural resources against skin aging. *Phytomed*. 2011;19:64–63.
- [4] Fulop T, Khalil A, Larbi A. The role of elastin peptides in modulating the immune response in aging and age-related diseases. *Pathol Biol*. 2012;60:28–33.

- [5] Kurtz A, Oh S-J. Age related changes of the extracellular matrix and stem cell maintenance. *Prev Med.* 2012;54:S50–S56.
- [6] Hsu M-F, Chiang B-H. Stimulating effects of *Bacillus subtilis* natto-fermented *Radix astragali* on hyaluronic acid production in human skin cells. *J Ethnopharmacol.* 2009;125:474–481.
- [7] Thring TSA, Hili P, dan Naughton DP. Anti-collagenase, anti-elastase, and anti-oxidant activities of extract from 21 plants. *BMC Compl Altern Med.* 2009;9:1-11.
- [8] Widowati W, Fauziah N, Herdiman H, Afni M, Afifah E, Kusuma, HSW, et al. Antioxidant and anti-aging assays of *Oryza sativa* extracts, vanillin and coumaric acid. *J Nat Remed.* 2016;16:88-89.
- [9] Yasui K, Sakurai H. Chemiluminescent detection and imaging of reactive oxygen species in live mouse skin exposed to UVA. *Biochem Biophys Res Commun.* 2000;269:131-136.
- [10] Park JS. Walnut husk ethanol extract possess antioxidant activity and inhibitory effect of matrix metalloproteinase-1 expression induced by tumor necrosis factor alpha in human keratinocyte. *Kor J Aesthet Cosmetol.* 2013;11:715-719.
- [11] Kim DS, Jeon BK, Mun YJ, Kim YM, Lee YE, Woo WH. Effect of *Dioscorea aimadoimo* on anti-aging and skin moisture capacity. *J Orient Physiol Pathol.* 2011;25:425-430.
- [12] Mruka DD, Silvestrini B, Mo MY, Cheng CY. Antioxidant superoxide dismutase—a review: its function, regulation in the testis, and role in male fertility. *Contraception.* 2002;65:305-311.
- [13] Moini H, Packer L, Saris NEL. Antioxidant and prooxidant activities of α -lipoic acid and dihydrolipoic acid. *Toxicol Appl Pharmacol.* 2002;182:84-90.
- [14] Cai Y, Luo Q, Sun M, Corke H. Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sci.* 2004;74:2157-2164.
- [15] Kaur C, Kapoor HC. Anti-oxidant activity and total phenolic content of some Asian vegetables. *Int J Food Sci Technol.* 2002;37:153-161.
- [16] Zhang Y-J, Liu Y-Q, Pu X-Y, Yang C-R. Iridoidal glycosides from *Jasminum sambac*. *Phytochem.* 1995;38:899-903.
- [17] Inagaki J, Watanabe N, Moon JH, Yagi A, Sakata K, Ina K, et al. Glycosidic aroma precursors of 2-phenylethyl and benzyl alcohols from *Jasminum sambac* flowers. *Biosci Biotechnol Biochem.* 1995;59:738-739.
- [18] Zhang ZF, Bian BL, Yang J, Tian XF. [Studies on chemical constituents in roots of *Jasminum sambac*]. *Zhongguo Zhong yao za zhi= Zhongguo zhongyao zazhi= China journal of Chinese materia medica.* 2004;29:237-239.
- [19] Younis A, Mehdi A, Riaz A. Supercritical carbon dioxide extraction and gas chromatography analysis of *Jasminum sambac* essential oil. *Pak J Bot.* 2011; 43:163-168.
- [20] Edris A, Franz C, Chizzola R. Isolation and characterization of the volatile aroma compounds from the concrete headspace and the absolute of *Jasminum sambac* (L.) Ait. (Oleaceae) flowers grown in Egypt. *Eur Food Res Technol.* 2008;226:621–626.
- [21] Widowati W, Mozef T, Risdian C, Yellianty Y. Anticancer and free radical scavenging potency of *Catharanthus roseus*, *Dendrophthoe petandra*, *Piper betle*, and *Curcuma mangga* extracts in breast cancer cell lines. *Oxidants Antioxid Med Sci.* 2013;2:137-142.
- [22] Widowati W, Wijaya L, Wargasetia T, Bachtiar I, Yellianty Y, Laksmiawati D. Antioxidant, anticancer and apoptosis-inducing effects of Piper extracts in HeLa cells. *J Exp Integr Med.* 2013;3:225-230.
- [23] Laksmiawati DR, Widyastuti A, Karami N, Afifah E, Rihibiha DD, Nufus H, Widowati W. Anti-inflammatory effects of *Anredera cordifolia* and *Piper crocatum* extracts on lipopolysaccharide-stimulated macrophage cell line. *Bangladesh J Pharmacol.* 2017;12:35-40.
- [24] Adnyana I, Abuzaid A, Iskandar E, Kurniati N. Pancreatic lipase and α -amylase inhibitory potential of mangosteen (*Garcinia mangostana* Linn.) pericarp extract. *Int J Med Res Health Sci.* 2016;5:23-28.
- [25] Bera TK, Chatterjee K, Ghosh D. In vitro antioxidant properties of the hydro-methanol

- extract of the seeds of *Switenia mahagoni* (L.) Jacq. BGM. 2015;7:18-24.
- [26] Widowati W, Rani AP, Hamzah RA, Arumwardana S, Afifah E, Kusuma HSW, et al. Antioxidant and antiaging assays of *Hibiscus sabdariffa* extract and its compounds. Nat Prod Sci. 2017;23:192-200.
- [27] Widowati W, Darsono L, Suherman J, Yellianty Y, Maesaroh M. High Performance Liquid Chromatography (HPLC) analysis, antioxidant, antiaggregation of mangosteen peel extract (*Garcinia mangostana* L.). Int J Biosci Biochem Bioinform. 2014;4:458.
- [28] Rusmana D, Wahyudianingsih R, Elisabeth M, Balqis B, Maesaroh M, Widowati W. Antioxidant activity of *Phyllanthus niruri* extract, rutin and quercetin. Indones Biomed J. 2017;9:84-90.
- [29] Widowati W, Widyanto RM, Husin W, Ratnawati H, Laksmiawati DR, Setiawan B, et al. Green tea extract protects endothelial progenitor cells from oxidative insult through reduction of intracellular reactive oxygen species activity. Iran J Basic Med Sci. 2014;17:702-709.
- [30] Utami S, Sachrowardi QR, Damayanti NA, Wardhana A, Syarif I, Nafik S, et al. Antioxidants, anticollagenase and antielastase potentials of ethanolic extract of ripe sesoot (*Garcinia picrorrhiza* Miq.) fruit as antiaging. J Herbmed Pharmacol. 2018;7:88-93.
- [31] Tu PTB, Tawata S. Antioxidant, antiaging, and anti-melanogenic properties of the essential oil from two varieties of *Alpinia zerumbet*. Molecules. 2015;20:16723-16740.
- [32] Irshad M, Chaudhuri PS. Oxidant-antioxidant system: role and significance in human body. Indian J Exp Biol. 2002;40:1233-1239.
- [33] Blois MS. Antioxidant determination by the use of stable free radical. Nature. 1958;181:1199.
- [34] Thaakur SR. Free radical scavenging activity of *Jasminum sambac*. JGTPS. 2014;5:1658-1661.
- [35] Shekhar S, Prasad MP. Studies on antioxidant properties of *Jasminum* species by FRAP assay. Int J Pure App Biosci. 2015;3:52-57.
- [36] Chiu TM, CC Huang, TJ Lin, JY Fang, NL Wu, CF Hung. 2009. In vitro and in vivo anti-photoaging effects of an isoflavone extract from soybean cake. J Ethnopharmacol. 2126:108-113.
- [37] Kim JH, Cho YH, Lee BC, Pyo HB. Cosmetic composition comprising extracts of *Pterocarpus santalinus* capable of inhibiting activity. Korea Patent. 2007. <http://patent.ipexl.com/KR/1020050076940.html>.
- [38] Kim S, JH Chung. Berberine prevents UV-induced MMP-1 and reduction of type I procollagen expression in human dermal fibroblasts. Phytomed. 2008;15:749-753.
- [39] Maity N, NK Nema, MK Abedy, BK Sarkar, PK Mukherjee. 2011. Exploring *Tagetes erecta* Linn flower for the elastase, hyaluronidase and MMP-1 inhibitory activity. J Ethnopharmacol. 2011;137:1300-1305.
- [40] Moon HI, J Lee, PZ Ok, JH Chung. The effect of flavonol glycoside on the expressions of matrix metalloproteinase-1 in ultraviolet-irradiated cultured human skin fibroblasts. J Ethnopharmacol. 2005;101:176-179.
- [41] Satardekar KV, Deodhar MA. Anti-ageing ability of Terminalia species with special reference to hyaluronidase, elastase inhibition and collagen synthesis in vitro. Int J Pharmacogn Phytochem Res. 2010;2:30-34.
- [42] Aslam MN, Lansky EP, Varani J. Pomegranate as a cosmeceutical source: Pomegranate fractions promote proliferation and procollagen synthesis and inhibit matrix metallo proteinases-1 production in human skin cells. J Ethnopharmacol. 2006;103:311-318.
- [43] Raffetto JD, Khalil RA. Matrix metalloproteinases and their inhibitors in vascular remodeling and vascular disease. Biochem Pharmacol. 2008;75:346-349.
- [44] Kim Y, Uyama H, Kobayashi S. Inhibition effects of (+)-catechin-aldehyde polycondensates on proteinases causing proteolytic degradation of extracellular matrix. Biochem Biophys Res Commun. 2004;320:256-261.
- [45] Baylac S, Racine P. Inhibition of human leukocyte elastase by natural fragrant extracts

- of aromatic plants. *Int J Aromather.* 2004;14:179-182.
- [46] Melzig MF, Löser B, Ciesielski S. Inhibition of neutrophil elastase activity by phenolic compounds from plants. *Pharmazie.* 2001;56:967-970.
- [47] Siedle B, Ciesielski S, Murillo R, Löser B, Castro V, Klaas CA, et al. Sesquiterpene lactones as inhibitors of human neutrophil elastase. *Bioorg Med Chem.* 2002;10:2855-2861.
- [48] Piwowarski JP, Kiss AK, Kozłowska-Wojciechowska M. Anti-hyaluronidase and anti-elastase activity screening of tannin-rich plant materials used in traditional Polish medicine for external treatment of diseases with inflammatory background. *J Ethnopharm.* 2011;137:937-941.
- [49] Garg A, Garg S, Zaneveld LJD, Singla AK. Chemistry and pharmacology of the Citrus bioflavonoid hesperidin. *Phytother Res.* 2001;15:655-659.
- [50] Chen MC, Ye YY, Ji G, Liu JW. Hesperidin upregulates heme oxygenase-1 to attenuate hydrogen peroxide-induced cell damage in hepatic L02 cells. *J Agric Food Chem.* 2010;58:3330-3335.
- [51] Elavarasan J, Velusamy P, Ganesan T, Ramakrishnan SK, Rajasekaran D, Periandavan K. Hesperidin-mediated expression of Nrf2 and upregulation of antioxidant status in senescent rat heart. *J Pharm Pharmacol.* 2012;64:1472-1482.
- [52] Wilmsen PK, Spada DS, Salvador M. Antioxidant activity of the flavonoid hesperidin in chemical and biological systems. *J Agric Food Chem.* 2005;53:4757-4761.
- [53] Sun K, Xiang L, Ishihara S, Matsuura A, Sakagami Y, Qi J. Anti-aging effects of hesperidin on *Saccharomyces cerevisiae* via inhibition of reactive oxygen species and UTH1 gene expression. *Biosci Biotech Biochem.* 2012;76:640-645.
- [54] Li R, Cai L, Xie XF, Yang F, Li J. Hesperidin suppresses adjuvant arthritis in rats by inhibiting synoviocyte activity. *Phytother Res.* 2010;24:S71-S6.
- [55] Gülçin İ. Antioxidant activity of eugenol: a structure-activity relationship study. *J Med Food.* 2011;14:975-985.
- [56] Ogata M, Hoshi M, Urano S, Endo T. Antioxidant activity of eugenol and related monomeric and dimeric compounds. *Chem Pharm Bull.* 2000;48:1467-1469.

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau PEER REVIEW

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Antioxidant and Antiaging Activities of Jasminum Sambac Extract, and its Compounds

Jumlah Penulis : 7 Orang

Nama-nama Penulis : **Wahyu Widowati**, Widya Janeva B, Sri Nadya, Annisa Amalia, Seila Arumwardana, Hanna Sari W. Kusuma, Yukko Arinta

Status Penulis : ~~Penulis Pertama~~ / Penulis ke 1 / Penulis Korespondensi **)

Identitas Jurnal Ilmiah : a. Nama jurnal : Journal of Reports in Pharmaceutical Science
 b. Nomor ISSN : 2322-1232, 2322-5106
 c. Vol., No., Bulan, Tahun : Vol. 7., No. 3., July 2018
 d. Penerbit : Wolters Kluwer Medknow Publications
 e. DOI Artikel (jika ada) : -
 f. Alamat Web Jurnal : jrpsjournal.com/
 g. Terindeks di : Scopus Q3, SJR 0.155

Kategori Publikasi Jurnal Ilmiah: Jurnal Ilmiah Internasional / Internasional Bereputasi **)
 (beri tanda ✓ yang dipilih)

Jurnal Ilmiah Nasional Terakreditasi

Jurnal Ilmiah Nasional / Nasional terindeks ***)

HASIL PENILAIAN (Peer Review) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi	Nasional Terakreditasi	Nasional ***)	
		<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,7
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			4,6
c.	Kecukupan dan kemutakhiran data/ informasi dan metodologi (30%)	12			4,8
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			4,6
	Total	40			38,7

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur.....
 Penulisan memenuhi standar karya ilmiah akademis, metoda yang digunakan sesuai dan tepat digunakan pada penelitian ini
- b. Ruang lingkup & kedalaman pembahasan.....
 Penelitian tentang potensi ekstrak bunga melati dan senyawa-senyawanya (Hesperidin, Eugenol) sebagai antioksidan & antiaging
- c. Kecukupan & kemutakhiran data serta metodologi.....
 Ide dasar penelitian baik membahas perbandingan ekstrak dan senyawa sebagai antioksidan, metode DPPH, FRAP, ABTS, metode antiaging, ankihy alurodinase, antielastase dan antiokogenase
- d. Kelengkapan unsur dan kualitas penerbit.....
 Jurnal JRPTS terindeks Scopus Q3 SJR 0,16 Penerbit Wolters Kluwer Medknow Publications
- e. Indikasi plagiasi.....

tidak terdapat indikasi plagiarisme atau self plagiarisme

f. Kesesuaian bidang ilmu

Paper bidang bioteknia, biomedik, sesuai dengan bidang ilmu penulis

REVIEWER 1



(Prof. Dr. Chrismis Novalinda Ginting, M.Kes)

NIK : 0115127801

UNIVERSITAS PRIMA INDONESIA

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau *PEER REVIEW*

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Antioxidant and Antiaging Activities of Jasminum Sambac Extract, and its Compounds
 Jumlah Penulis : 7 Orang
 Nama-nama Penulis : **Wahyu Widowati**, Widya Janeva B, Sri Nadya, Annisa Amalia, Seila Arumwardana, Hanna Sari W. Kusuma, Yukko Arinta
 Status Penulis : ~~Penulis Pertama~~ / Penulis ke 1 / Penulis Korespondensi **)
 Identitas Jurnal Ilmiah : a. Nama jurnal : Journal of Reports in Pharmaceutical Science
 b. Nomor ISSN : 2322-1232, 2322-5106
 c. Vol., No., Bulan, Tahun : Vol. 7., No. 3., July 2018
 d. Penerbit : Wolters Kluwer Medknow Publications
 e. DOI Artikel (jika ada) : -
 f. Alamat Web Jurnal : jrpsjournal.com/
 g. Terindeks di : Scopus Q3, SJR 0.155

Kategori Publikasi Jurnal Ilmiah : ~~Jurnal Ilmiah Internasional~~ / Internasional Bereputasi **)
 (beri tanda \checkmark yang dipilih)
 Jurnal Ilmiah Nasional Terakreditasi
 Jurnal Ilmiah Nasional / Nasional terindeks ***)

HASIL PENILAIAN (*Peer Review*) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi <input checked="" type="checkbox"/>	Nasional Terakreditasi <input type="checkbox"/>	Nasional ***) <input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,8
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			11,5
c.	Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12			11,7
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			11,8
	Total	40			36,8

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur.....
Jurnal ini sudah cukup lengkap dan ada kesesuaian antara unsur dan isinya
- b. Ruang lingkup & kedalaman pembahasan.....
Ruang lingkup bahasan sudah memadai dan ada kedalaman pembahasannya
- c. Kecukupan & kemutakhiran data serta metodologi.....
Secara umum metodologi masih terbatas dan belum sepenuhnya
- d. Kelengkapan unsur dan kualitas penerbit.....
Kualitas penerbit sudah memenuhi reputasi yang baik
- e. Indikasi plagiasi.....

Belum ditemukan adanya unsur plagiasi

f. Kesesuaian bidang ilmu

Jurnal ini sesuai dengan bidang Ilmu yang diteliti

REVIEWER 2



(Prof. Dr. Ermi Girsang, M. Kes)

NIK : 0117057501

UNIVERSITAS PRIMA INDONESIA

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau *PEER REVIEW*

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : Antioxidant and Antiaging Activities of Jasminum Sambac Extract, and its Compounds
 Jumlah Penulis : 7 Orang
 Nama-nama Penulis : **Wahyu Widowati**, Widya Janeva B, Sri Nadya, Annisa Amalia, Seila Arumwardana, Hanna Sari W. Kusuma, Yukko Arinta
 Status Penulis : ~~Penulis Pertama~~ / Penulis ke 1 / Penulis Korespondensi **)
 Identitas Jurnal Ilmiah : a. Nama jurnal : Journal of Reports in Pharmaceutical Science
 b. Nomor ISSN : 2322-1232, 2322-5106
 c. Vol., No., Bulan, Tahun : Vol. 7., No. 3., July 2018
 d. Penerbit : Wolters Kluwer Medknow Publications
 e. DOI Artikel (jika ada) : -
 f. Alamat Web Jurnal : jrpsjournal.com/
 g. Terindeks di : Scopus Q3, SJR 0.155

Kategori Publikasi Jurnal Ilmiah: Jurnal Ilmiah Internasional / Internasional Bereputasi **)
 (beri tanda \checkmark yang dipilih)
 Jurnal Ilmiah Nasional Terakreditasi
 Jurnal Ilmiah Nasional / Nasional terindeks ***)

HASIL PENILAIAN (*Peer Review*) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi <input checked="" type="checkbox"/>	Nasional Terakreditasi <input type="checkbox"/>	Nasional ***) <input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,75
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			11,55
c.	Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12			11,75
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			11,7
	Total	40			36,95

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur.....
 Penulisan memenuhi Standard Karya Ilmiah akademis, metoda yang digunakan sesuai dan tepat digunakan pada penelitian ini
 Jurnal ini sudah cukup lengkap dan ada kesesuaian antara unsur dan kinya
- b. Ruang lingkup & kedalaman pembahasan.....
 Penelitian tentang potensi ekstrak bunga melati dan senyawa-senyawanya (lisperidin, Eugenol) sebagai antioksidan & antiaging

Ruang lingkup bahasan sudah memadai dan ada kedalaman pembahasannya

c. Kecukupan & kemutakhiran data serta metodologi... Ide dasar penelitian baik membahas perbandingan ekstrak dan senyawa sebagai antioksidan metode DPPH, FRAP, ABTS metode antiaging, antihyaluronidase, antielastase dan antikolagenase

Secara umum metodologi masih terbatas dan belum sepenuhnya

d. Kelengkapan unsur dan kualitas penerbit

Jurnal JAPTS terindeks scopus Q3 SJR 0,16 Penerbit Wolters kluwer Medknow Publications

Kualitas penerbit sudah memenuhi reputasi yang baik

e. Indikasi plagiasi

Tidak terdapat indikasi plagiarisme atau self plagiarisme

Tidak ditemukan adanya unsur plagiasi

f. Kesesuaian bidang ilmu

Jurnal ini sesuai dengan bidang ilmu yang diteliti

Medan,
Reviewer 2

(Prof. Dr. Ermi Girsang, M.Kes)
NIK : 0117057501
UNIVERSITAS PRIMA INDONESIA

Medan,
Reviewer 1

(Prof. Dr. Chrismis Novalinda Ginting, M.Kes)
NIK : 0115127801
UNIVERSITAS PRIMA INDONESIA