



Microstructural Characterization of the *Garcinia mangostana* Fruit at Different Age Level

Marisca Evalina Gondokesumo^{1*}, Bambang Pardjianto², Sutiman Bambang Sumitro³, Wahyu Widowati⁴ and Arbi Dimiyati⁵

¹Faculty of Medicine, Brawijaya University, Malang, Indonesia; mariscaevalina@gmail.com

²Department of Plastic Surgery, Saiful Anwar General Hospital, Faculty of Medicine, Brawijaya University, Malang, Indonesia

³Department of Biology, Faculty of Sciences, Brawijaya University, Malang, Indonesia

⁴Faculty of Medicine, Maranatha University, Bandung, Indonesia

⁵Center for Science and Technology of Advanced Materials, PSTBM-BATAN, Indonesia

Abstract

Mangosteen (*Garcinia mangostana*) is one of the most popular fruits which has been widely used medicinally. The major constituents are mostly found in the pericarp particularly, xanthenes, which are tricyclic isoprenylated polyphenols. Xanthenes have been reported to for its anti-oxidant, anti-inflammatory, anti-bacterial, anti-proliferative, proapoptotic and anti-carcinogenic activities. In this study, the mangosteen pericarps in various age levels were investigated with electron microscopy and X-ray spectroscopy. The results revealed the mangosteen at age level 3 showed the highest concentration of xanthone while those younger and older mangosteens exhibited lower xanthone content. The mangosteen pericarp with highest xanthone concentration showed crystalline structure and contained small amount of silver and gold particles which was analyzed with EDX measurement and confirmed by the XRD spectrum.

Keywords: Mangosteen, Particle Size, SEM, Xanthone, X-ray Diffraction, XRD

1. Introduction

Plants and their fruits have been used since ancient time as supplement and medicine to treat diseases. Approximately two-third of world population uses plant based products as their main medicine¹. *Garcinia mangostana* L. (Clusiaceae), commonly known as mangosteen, have been used as a traditional medicine in Southeastern Asia for the treatment of diarrhea, dysentery, inflammation, ulcers and wound healing^{2,3}. Mangosteen is cultivated mainly in Indonesia, Malaysia, Philipina and Thailand⁴⁻⁶. Medicinal properties of *G. mangostana* is associated with its bioactive compounds, xanthone. Xanthone is a tricyclic isoprenylated polyphenol. Previous investigation shows







that xanthone is antioxidant^{5,7}, anti-bacterial, anti-proliferative, proapoptotic, anti-carcinogen⁸⁻¹⁰ and anti-inflammatory¹¹.

Xanthone can be found in every part of mangosteen, mainly in fruit skin. Mangosteen contains 17% of pericarp¹², which contains 62.05% of water, 0.63% of fat, 0.71% of protein, 1.17% of total sugar and 35.61% of carbohydrate¹³. Xanthone content in the mangosteen pericarp is about 108 to 124 mg per 100 g^{14,15}. The ripeness level significantly affects the xanthone content of mangosteen¹⁶. In this study, we have investigated the physical and chemical properties of mangosteen pericarps depending on the age level in accordance to Palapol, et al. (2009)¹⁷ (Table 1), using various characterization techniques such as PSA (Particle Size Analyzer), Scanning Electron

*Author for correspondence

Microscope (SEM), X-ray Diffraction (XRD), and X-ray Fluorescence (XRF). The study aims to get insight into the microstructure of mangosteen pericarp in accordance to their age level and xanthone content. Furthermore, it can give insight to maximize the yield of xanthenes.

Table 1. Age level classification of mangosteen

Age Level	Properties
 Level 1	Green yellowish, high mucus content. The inner part of fruit cannot be separated from the flesh.
 Level 2	Yellow reddish with distribution of red colored fleck. Lower mucus content. The inner part of the fruit still cannot be separated from flesh.
 Level 3	Red brownish. Mucus still exists. The inner part of fruit can be separated from the flesh.
 Level 4	The color becomes red purple. The mucus is lesser. The inner part of fruit can be separated from the flesh and the fruit can be consumed.
 Level 5	Purple reddish. Fruit starts to ripe. The mucus disappears, and the fruit can be easily separated.
 Level 6	Purple blackish. The fruit is ripened and can be eaten immediately.

2. Methods

2.1 Extraction

The extraction of mangosteen was performed following standard procedure. Mangosteen pericarp was obtained from Bandung, West Java. Mangosteen was air dried until the water content was about 5% prior to crushing in an electrical blender. 700 g of mangosteen pericarp mass was macerated using 96% ethanol (1 : 5)

for 5 days and stirred every 3 hours. The extract was dried by evaporation and freeze dried to obtain the powder. The powder was stored in desiccators until used.

2.2 Characterization

The mangosteen extract in various age levels were subjected to microstructure characterization using PSA, SEM, XRD and XRF. For SEM analysis, the mangosteen extract was poured homogeneously onto the carbon double tip on the SEM sample holder stub. For the elemental composition in SEM, an energy dispersive X-ray (EDX) detector attached on SEM (SEM JEOL JSEM2650LV) was used. For XRD analysis Imperian with Cu source was used. The XRD results were analyzed using Ritveld Refinement to derive information about the phase and crystals may occur.

3. Results

PSA measurement of the mangosteen extract in various age levels is presented in Figure 1 and Table 2. Particle of mangosteen powder at level 5 exhibited the smallest size, whereas those at level 3 exhibited the highest size.

Table 2. Particle size of mangosteen pericarp extract as measured by PSA

Sample	Size (nm)
1	81.8 ± 21.1
2	72.6 ± 20.4
3	1244.1 ± 341.0
4	1112.2 ± 256.5
5	57.0 ± 14.5
6	199.2 ± 58.9

Data are presented in Mean ± Standard Deviation

Figure 2 shows the SEM image of the microstructure of mangosteen pericarp extract at the age levels 1-6 which generally showed granular and plate like structure composing of large coagulates. Sample of age levels 3 and 6 have shown clearly smaller powder grains than those of the sample of age levels 1, 2, 4 and 5. It was distinguished by the contrast deficiency by the sample with fine grain size. It is almost impossible to measure single grain on the sample 1, 2, 4 and 5 because the sample forms large block of particle agglomerations.






Figure 1. Particle size distribution of, a. Sample 1; b. Sample 2; c. Sample 3; d. Sample 4; e. Sample 5; f. Sample 6.






Figure 2. SEM SE images of mangosteen pericarp samples at age level and their corresponding EDX spectrum, a. Sample 1; b. Sample 2; c. Sample 3; d. Sample 4; e. Sample 5; f. Sample 6.

The corresponding EDX spectrums are also represented in Figure 2 on the right side revealing the elemental composition of the samples. Mainly the samples contain C, O, K and Cl. Sample 1 revealed the elemental content Si, Cl and K about 0.33%, 0.22% and 0.72% respectively. Sample 2 was crystalline in nature containing Al 0.1%, whilst Si was not detected. Level of Cl and K were significantly reduced. Sample 3 showed globular form, softer than samples 1 and 2 which was about 5 μm , as shown in EDX results. Both Cl and K content was significantly reduced, amorphous and its silicone was reduced as well. Sample 4 was seen as

granules and also in combined form. Sample 4 has granule size of 20 μm , crystalline and its level of Si was reduced. The content of Cl was 0.10% and K was 0.28%. Sample 5 showed granules form and also combined form. Its granule were between 5-50 μm , crystalline and Si content was reduced. Sample 5 were mainly composed by C, O, Cl and K with 61,18%, 37,49%; 0,19% and 0,69% respectively. Sample 6 showed granules form and also combined form. Granule was 10 μm , crystalline and Si content was reduced. Main components of sample 5 were C, O, Cl, and K with 62,41%, 36,67%; 0,23% and 0,68%. respectively.




Figure 3. XRD analysis of mangosteen peel extracts, a. Sample 1; b. Sample 2; c. Sample 3; d. Sample 4; e. Sample 5; f. Sample 6.

Table 3. Elemental concentration of mangosteen peel extract analyzed with X-ray fluorescence

Elements	Samples					
	1	2	3	4	5	6
K	50.96	50.59	53.58	50.21	51.22	51.03
Cl	13.45	11.18	12.83	9.61	11.90	12.20
Ca	4.92	8.10	1.48	8.56	5.90	4.67
Px	1.78	1.68	1.67	1.50	1.39	1.75
Sx	1.01	1.25	2.06	1.25	1.01	2.20
Na	1.37	1.27	1.08	1.67	1.77	1.22
cd	1.50	0.77	1.89	0.91	1.18	1.11
Ag	1.33	2.01	3.20	1.31	1.52	–
Fe	0.924	0.836	0.821	1.05	1.986	–
Zn	0.879	0.386	0.257	0.482	0.712	–
Mo	0.712	–	0.854	0.678	–	0.575
Rb	0.75	0.873	0.88	0.985	1.18	0.523
Mg	0.410	0.425	0.290	0.392	0.315	0.331
Si	0.295	0.294	0.249	0.280	0.569	0.401
Zr	0.239	0.209	0.125	0.219	0.167	0.249
Ar	0.291	0.291 / 0.254	0.339	0.246	0.203	0.203
Mn	0.218	0.218	0.156	0.453	0.195	0.293
Ti	0.115	0.115	–	–	0.089	0.504
Ba	–	–	0.80	–	0.84	–
Au	–	–	0.74	–	–	–

The XRD patterns of mangosteen pericarp extracts are shown in Figure 3. As shown in XRD analysis, samples 1, 2 and 3 were amorphous. Although sample 2 was previously seen as crystalline, it was mainly amorphous based on XRD results. Lowest peak in samples 4, 5 and 6 indicates carbon crystal.

XRF is X-ray method similar to EDX, yet it measures smaller element above 11 atomic number. XRF was performed to confirm results obtained from EDX and XRD analysis. Results of XRF are shown in Table 3. Most of elements were in accordance to previous results analyzed with EDX and there were also several more elements found. Elements K, Cl, Ca, Px, Sx, Na, Cd, Rb, Mg, Si, Zr, Ar and Mn were detected in all samples. Meanwhile, Ag, Fe, Zn, Mo and Ti was found only in sample 1. Similarly, Ag, Fe, Zn and Ti were detected in sample 2; Ag, Fe, Zn, Mo, Ba and Au were detected in sample 3; Ag, Fe, Zn and Mo were detected in sample 4; Ag, Fe, Zn, Ti and Ba were detected in sample 5 and Mo and Ti were detected in sample 6.

4. Discussion

Dehydration is an efficient alternative for fruit storage in which reduction of water is associated with the decline of chemical and enzymatic reactions that can damage foods. Gentle drying techniques, such as osmodehydration or lyophilisation (freeze drying), have been used as new alternative for the commercialization of tropical fruits to reduce the difficulties for its handling and transport and to acquire added value solids products¹⁸. In this study, we evaluated microstructure of freeze dried mangosteen pericarp extract at various age levels. The structural studies of the fruit powders were done using PSA (Particle Size Analyzer), Scanning Electron Microscope (SEM), X-ray Diffraction (XRD) and X-ray Fluorescence (XRF).

In the present study, particle size of the mangosteen extract was clearly heterogeneous. Particle of mangosteen powder at level 5 exhibited the smallest size, whereas those at level 3 exhibited the highest size. Particle size distribution does not follow a typical ideal Gauss bell

form. Additionally, it seems that the levels of age were not in correlation with the particle size. The particle size may depend on the processing method. This can be also originated from the coagulation of mangosteen particles.

SEM technique is a powerful tool for determining and observing the caking phenomenon on the powder surface. The corresponding EDX spectrums revealing the elemental composition of the samples, exhibited component C, O, K and Cl in samples. XRF is X-ray method similar to EDX, yet it measures smaller element above 11 atomic numbers. XRF was performed to confirm results obtained from EDX and XRD analysis. Most of elements were in accordance to previous results analyzed with EDX and there were also several more elements found.

Crystallization is essential for the stability of powdered juice and can be investigated through XRD analysis. Amorphous state is indicated with diffused and large peaks in which the molecules are disorderly displayed producing disperse bands, whereas crystalline materials yield sharp and defined peaks since they are presented in a highly ordered state¹⁸. In this study, samples 1, 2 and 3 were amorphous, whilst samples 4, 5 and 6 indicates carbon crystal.

Xanthenes are flavonoid commonly found in mangosteen peel¹⁹. In the present study, mangosteen at age level 3 showed the highest concentration of xanthone while those younger and older mangosteen exhibited lower xanthone content. Xanthenes have been reported to exhibit various medicinal properties including anti-microbial, anti-malarial, anti-oxidant and anti-inflammatory. It has been reported that scavenging activity of mangosteen is higher when the fruit were younger and its activity decrease as the fruit age increase¹⁶. It is suggested to be associated with decreased synthesis of antioxidants during the fruit ripening process resulting in oxidative stress. Chemical constituents and antioxidant capacity of fruit are affected by fruit growth stage in which immature fruit contains the highest antioxidant capacity²⁰. Study done by Huang, et al. also found decreased antioxidant activity in the late phase of fruit enlargement which also explains correlation between increasing oxidative stress and metabolic changes with aging of citrus fruits²¹.

5. Conclusion

Mangosteen at age level 3 showed the highest concentration of xanthone while those younger and older mangosteen exhibited lower xanthone content. Mangosteen pericarp with highest xanthone concentration has crystalline structure and contains small amount of silver and gold particles. These findings provide information regarding characteristics of mangosteen peel that can be further used for the optimization of xanthone content.

6. Acknowledgment

This study was personally funded by authors. We would also like to thank Indonesian Institute of Life Science (LIPI) for its assistance and supervision.

7. Conflict of interest

Authors declare that there is no conflict of interest.

8. Reference

1. Suksamrarn S, Suwannapoch N, Ratananukul P, Aroonlerk N, Suksamrarn A. Xanthenes from the Green Fruit Hulls of *Garcinia mangostana*. J Nat Prod. 2002; 65(5):761–3. PMID: 12027762. Crossref.
2. Moongkarndi P, Kosem N, Kaslungka S, Luanratana O, Pongpan N, Neungton N. Anti-proliferation, anti-oxidation and induction of apoptosis by *Garcinia mangostana* (mangosteen) on SKBR3 human breast cancer cell line. J Ethnopharmacol. 2004 Jan 1; 90(1):161–6. PMID: 14698525. Crossref.
3. Farnsworth NR, Bunapraphatsara N. Thai medicinal plants recommended for primary health care system. Medicinal Plant Information Center; 1992.
4. Ramasamy S, Mazlan NA, Ramli NA, Rasidi WN, Manickam S. Bioactivity and stability studies of anthocyanin-containing extracts from *Garcinia mangostana* L. and *Etligeria elatior* Jack. Sains Malaysiana. 2016; 45(4):559–65.
5. Tjahjani S, Widowati W, Khiong K, Suhendra A, Tjokropranoto R. Antioxidant properties of *Garcinia mangostana* L (mangosteen) rind. Procedia Chemist. 2014; 13:198–203. Crossref.

6. Aminah LN, Leong ST, Wong YS, Ong SA, Kairulazam CK. Biodiesel production of *Garcinia mangostana* Linn seeds by two-phase solvent extraction and alkali-catalyzed transesterification. *Int J Chem Eng Appl*. 2013; 4(3):92–5. Crossref.
7. Zarena AS, Sankar KU. A study of antioxidant properties from *Garcinia mangostana* L. pericarp extract. *Acta Sci Pol Technol Aliment*. 2009; 8(1):23–34.
8. Seesom W, Jaratrungratawee A, Suksamrarn S, Mekseepralard C, Ratananukul P, Sukhumsirichart W. Antileptospiral activity of xanthenes from *Garcinia mangostana* and synergy of gamma-mangostin with penicillin G. *BMC Complement Altern Med*. 2013; 13(1):182. PMID: 23866810 PMCID: PMC3734031. Crossref.
9. Shan T, Ma Q, Guo K, Liu J, Li W, Wang F, Wu E. Xanthenes from mangosteen extracts as natural chemopreventive agents: Potential anticancer drugs. *Curr Mol Med*. 2011; 11(8):666–77. PMID: 21902651 PMCID: PMC3237908. Crossref.
10. Akao Y, Nakagawa Y, Nozawa Y. Anti-cancer effects of xanthenes from pericarps of mangosteen. *Int J Mol Sci*. 2008; 9(3):355–70. PMID: 19325754 PMCID: PMC2635669. Crossref.
11. Widowati W, Darsono L, Suherman J, Fauziah N, Maesaroh M, Erawijantari PP. Anti-inflammatory effect of mangosteen (*Garcinia mangostana* L.) peel extract and its compounds in LPS-induced RAW264. 7 cells. *Nat Prod Sci*. 2016; 22(3):147–53. Crossref.
12. Chaovanalikit A, Mingmuang A, Kitbunluewit T, Choldumrongkool N, Sondee J, Chupratum S. Anthocyanin and total phenolics content of mangosteen and effect of processing on the quality of mangosteen products. *Int Food Res J*. 2012; 19(3):1047–53.
13. Yunitasari L. *Gempur 41 Penyakit dengan Buah Manggis*. Yogyakarta: Pustaka Baru Press; 2011.
14. Yatman E. Kulit Buah Manggis Mengandung Xanton yang Berkhasiat Tinggi. *Widya*. 2012; 29:2–9.
15. Iswari, K., Harnel and Afdi E. *Kajian Teknologi Pengolahan Manggis Mendukung Agribisnis Manggis di Sumbar*. Research Report of BPTP West Sumatera; 2006.
16. Kurniawati A. Character, xanthone content and antioxidant properties of mangosteen fruit's hull (*Garcinia mangostana* L.) at Several Fruit Growth Stadia. *Indones J Agron*. 2017; 39(3):188–92.
17. Palapol Y, Ketsa S, Stevenson D, Cooney JM, Allan AC, Ferguson IB. Color development and quality of mangosteen (*Garcinia mangostana* L.) fruit during ripening and after harvest. *Postharvest Biol Technol*. 2009; 51(3):349–53. Crossref.
18. Athmaselvi KA, Kumar C, Balasubramanian M, Roy I. Thermal, structural and physical properties of freeze dried tropical fruit powder. *J Food Process*. 2014; 2014.
19. Zadernowski R, Czaplicki S, Naczka M. Phenolic acid profiles of mangosteen fruits (*Garcinia mangostana*). *Food Chem*. 2009; 112(3):685–9. Crossref.
20. Çelik H, Ozgen M, Serce S, Kaya C. Phytochemical accumulation and antioxidant capacity at four maturity stages of cranberry fruit. *Sci Hort*. 2008; 117(4):345–8. Crossref.
21. Huang R, Xia R, Hu L, Lu Y, Wang M. Antioxidant activity and oxygen-scavenging system in orange pulp during fruit ripening and maturation. *Sci Hort*. 2007; 113(2):166–72. Crossref.