

A Computational Study to Predict Wound Healing Agents from the Peel of the Mangosteen (*Garcinia mangostana* L.) Extract

Marisca Evalina Gondokesumo^{1,2*}, Sutiman Bambang Sumitro³,
Kusworini Handono⁴, Bambang Pardjianto⁵,
Wahyu Widowati⁶, Didik Huswo Utomo³

¹Biomedical Sciences Doctoral Study Program, Faculty of Medicine
Brawijaya University
Veteran Str., Malang 65145, East Java, Indonesia
E-mail: marisca@staff.ubaya.ac.id

²Faculty of Pharmacy
University of Surabaya
Raya Rungkut Str., Surabaya 60293, East Java, Indonesia

³Department of Biology, Faculty of Mathematics and Natural Sciences
Brawijaya University
Veteran Str., Malang 65145, East Java, Indonesia
E-mails: sutiman@ub.ac.id, dknatow@gmail.com

⁴Department of Clinical Pathology, Faculty of Medicine
Brawijaya University
Veteran Str., Malang 65145, East Java, Indonesia
E-mail: dr.kusworini@gmail.com

⁵Department of Plastic Surgery, Faculty of Medicine
Brawijaya University
Veteran Str., Malang 65145, East Java, Indonesia
E-mail: bambangplast@yahoo.com

⁶Faculty of Medicine
Maranatha Christian University
65 Prof. drg. Surya Sumantri, M.P.H. Str.
Bandung 40164, West Java, Indonesia
E-mail: wahyu_w60@yahoo.com

*Corresponding author

Received: August 18, 2018

Accepted: June 05, 2020

Published: September 30, 2020

Abstract: This study aimed to identify potentially active compounds from mangosteen peel extracts that heal skin burns and to evaluate their molecular mechanisms. There are about 120 compounds that have been identified in mangosteen peel by datamining, including 60 types of xanthones. The SMILE format of each compound was downloaded from the PubChem database. The compound data was then analyzed for potential anti-inflammatory, antioxidant, and antibacterial activities with PASS SERVER. To prediction mechanisms, firstly, target protein predictions were made with Swiss target prediction and HitPick. Target Protein data was then analyzed by interaction with STRING to determine its molecular mechanism. Further analysis of the shortest pathway was done with Cytoscape. The screening process of fifty compounds, which are predicted to have anti-inflammatory, antioxidant, and antibacterial activities by PASS SERVER. The four highest potency compounds were Smeaxanthone A, Garcinone E, γ -mangosteen, and Gartanin, which were types of xanthone. Based on the result of computational study these four compounds have

potential targets prediction related to interleukin 6, epidermal growth factor, and transforming growth factor beta 1, that are involved in the regulation of epithelial cell proliferation which plays a role in the healing process of skin burns.

Keywords: *Mangosteen peel, Skin burns, Wound healing, Xanthones, Molecular docking.*

Introduction

Burn injuries contributed to 180,000 deaths in 2004, while nearly 11 million people worldwide were severely burned and required medical treatment [32]. Wound healing involves 3 overlapping phases: inflammation, cell proliferation, and remodeling [8]. Antioxidants, anti-inflammatory agents, and antimicrobials all have roles in the wound healing process and prevent further aggravation of the wound condition [5].

Many factors slow the wound repair process, such as the presence of oxygen free radicals [33]. Reactive Oxygen Species (ROS), along with their derivatives, are abundant at the wound site [6]. Open wounds are prone to bacterial infection and provide an entry point for systemic infections [5]. When a cell is injured, inflammatory mediators such as cytokines, tumor necrosis factor (TNF), interleukin-1 (IL-1) from leukocytes, monocytes and macrophages are released as a response to pain. When this happens the production of cyclooxygenase (COX) 5-lipoxygenase (5-LO), prostaglandins (PGE) and Nitric Oxide (NO) are increased [3, 19, 44].

Generally, wound treatment of skin burns with early antioxidant therapy reinforces cellular antioxidant defense mechanisms, decreases free oxygen radicals, and promotes the healing process [4]. Free radical scavenging agents involve in the de-activation, reduction, and ROS removal, as well as promoting the wound-healing process. Consequently, topical applications of compounds with free radical scavenging properties can protect tissues from oxidative damage and also significantly improve wound healing [8, 38]. Topical antibacterial agents are also commonly used to facilitate wound healing in patients [38].

Over many years, herbal medicines have been widely utilized for natural medication of skin burn injuries [2, 16, 17, 37]. *Garcinia mangostana* L. (Clusiaceae), known as mangosteen, has been used as a traditional medicine in southeastern Asia for diarrhea, dysentery, inflammation, and ulcers, as well as for wound healing [1, 27, 30].

Mangosteen rind extract and purified constituents have been subjected to a vast array of biological tests relating to infectious diseases, cancer chemoprevention and chemotherapy, diabetes, and neurological conditions [4, 21, 27, 43]. Mangosteen extract and its constituent xanthones have been shown to have antioxidant [7, 39], anti-inflammatory [44], and antibacterial activities [20].

In our previous research, we have analyzed the main components peel of mangosteen extract containing Smeaxanthone A, Garcinone E, γ -mangosteen, and Gartanin using LC-MS/MS [13]. We used peel of mangosteen extract in the administration of topical gels to treat burns on the skin of mice that show good recovery results with the mechanism of increasing Epidermal Growth Factor (EGF) expression through immunohistochemistry and confirmed by western blot protein results [14].

The purpose of computational studies is to prove that there are major components contained in mangosteen peel extract, which has strong potential for skin repair in burning conditions.

Materials and methods

Collection of compounds in mangosteen rind extract

Active compounds were identified from datamining the Knapsack herbal database. The structure of each compound was further analyzed through the simplified molecular-input line-entry system (SMILES) in PubChem (<http://pubchem.ncbi.nlm.nih.gov>).

Prediction of biology activity

Biological activities of compounds were analyzed with the Prediction of Activity Spectra for Substances (PASS) server (<http://www.pharmaexpert.ru/passonline/>). Predictions were made using the structure-activity relationship (SAR) approach. Compounds with a similar structure and active cluster were considered to possess similar activities. For the PASS Server, the probability that a compound was active (Pa) was set at a threshold 0.7, such that $Pa > 0.7$ indicated a compound with a high probability for activity on experimentation [10]. The biological properties measured included anti-inflammatory, antioxidant, and antibacterial activities. The prediction was based on cluster analysis of potential compounds. A combination of compounds with anti-inflammatory, antioxidant, and antibacterial activities was expected to enhance wound healing.

Prediction of targeted protein

The potentially most active compounds were further analyzed with the servers HitPick (<http://mips.helmholtz-muenchen.de/hitpick/>) and Swiss Target Prediction (<http://www.swisstargetprediction.ch/>), using a targeted-focused compound library approach to make identifications. The principle was to predict structural aspects and functional activities of new compounds based on similarities to compounds in the database [25]. The biochemical pathways of targeted proteins were then analyzed to study the processes that could be targeted by drugs.

Pathway of compound-protein interaction

Potential interactions between targeted and others proteins were analyzed with STRING (<http://string-db.org>). The primary interaction unit in STRING was one of functional association [36]. This database was usually used to propose new pathways regarding drug interactions. Pathways were further analyzed with Cytoscape (<http://www.cytoscape.org>) to determine the most essential pathway as well as the type of interaction in the protein network.

Results and discussion

Several docking tools were combined to provide more reliable results for the understanding of compound-target relationships and the construction of herb-target networks. In the present study, all compounds from mangosteen peel extract were predicted to be involved in wound healing by interacting with single or multiple target proteins. This constraint gave a clear molecular mechanism for novel formula designs where active herbs were linked with different disease-related proteins. Disease intervention could be effectively achieved by simultaneously attacking multiple targets [31, 45].

Compound collecting with the Knapsack database identified about 120 compounds in mangosteen rind extract. Furthermore, SMILE analysis from PubChem identified 50 compounds. Active compounds that have combined activities including antioxidant, antiinflammatory and antibacterial, could be promising candidates in wound healing (Fig. 1).




Fig. 1 Prediction of antioxidant, antiinflammatory, and antibacterial activity of compounds in mangosteen rind extract by PASS SERVER

Fig. 2 shows Smeatxanthone A possessed the highest probability with regard to active antioxidant, antiinflammatory, and antibacterial activities ($P_a = 0.700$), followed by Garcinone E ($P_a = 0.694$), γ -mangosteen ($P_a = 0.693$) and Gartanin ($P_a = 0.692$). The four compounds are major components in mangosteen peel extract, which have been identified by their contents using LC-MS/MS [13], and mangosteen peel extract in gel preparations has activity to repair burns on mice skin in our previous research [14].




Fig. 2 Four compounds (xanthenes group) with most active prediction as antioxidant, antiinflammatory, and antibacterial by PASS SERVER

The potential mechanism of these compounds was identified based on prediction analysis. Results showed potential targets of these compounds were related to the expression of interleukin 6 (IL6), EGF, and transforming growth factor beta 1 (TGFB1) (Fig. 3, Table 1).




Fig. 3 Herbs-protein interaction

Table 1. Expression score based on interaction prediction (scale 0-1)

Protein	Expression score
TGFB1->FGF2	0.450
TGFB2->IL6	0.517

Results showed that the expression of IL6 might be affected by mangosteen extracts through TGFB2. Expression of IL6 was higher than fibroblast growth factor 2 (FGF2). The primary mechanisms affected in the body comprised 2 areas of regulation, epithelial cell proliferation (FGF2, TGFB1, IL6, TGFB2), and leukocyte homeostasis (TGFB1, IL6). These pathways were strongly associated with wound healing, epithelium cell proliferation and leukocytic elimination of antigens at the wound site. The regulation of these pathways was further analyzed to identify the potential modes of action of the compounds in mangosteen rind (Fig. 4).




Fig. 4 Most activated pathways by mangosteen rind extract

Results showed prostaglandin-endoperoxide synthase 2 (PTGS2) had a significant role in these pathways. This enzyme would cause IL6 to be activated, mostly, followed by EGF and TGFB1. TGFB1, EGF, and IL6 have crucial roles in wound healing. The TGFB family of proteins consists of multifunctional regulators that have been linked to wound repair, affecting all cell types involved in the process. TGFB1 has the broadest spectrum of action. It can induce chemotaxis of the inflammatory cells into the wound [12], as well as both angiogenesis and extracellular matrix formation; and it also initiates granulation tissue formation during and after inflammation [12, 26]. The primary structural component of granulation tissue is collagen, which can be induced directly by TGFB1 [11, 23, 42]. TGFB1 can stimulate wound contraction by inducing fibroblasts to differentiate to myofibroblasts [41, 42]. These contractile cells express proteins such as α -smooth muscle actin (SMA), stress fibers, vinculin-containing adhesion complexes and fibronectin fibrils for wound contraction [41].

IL6 was previously reported to precisely modulate the expression of TGFB in the skin and IL6KO fibroblasts [24]. IL6 is a pleiotropic cytokine involved in the growth and differentiation of numerous cell types, including those of dermal and epidermal origin [35]. Epidermal keratinocytes are a primary producer of IL6 within the skin, while macrophages, Langerhans cells and fibroblasts in the dermis represent other sources of the cytokine [28]. Increased levels of IL6 have been associated with some skin pathologies, such as psoriasis [15], scleroderma [22] and systemic lupus erythematosus [9]. Overexpression of IL6 in the skin of normal rats induces epidermal proliferation and inflammation [34], while overexpression in transgenic mice results in little more than a thickened stratum corneum [40].

The EGF family comprises multiple mediators of tissue-specific homeostasis of proliferation and differentiation. Human epidermal keratinocytes maintain high steady-state levels of extracellular-signal-regulated kinase (ERK) activity, mostly via their autocrine production of EGF receptor (EGFR) ligands [18]. The EGFR can be activated by transactivation from a variety of G-protein-coupled receptors, integrins, and cytokine receptors, so that it can act as a significant transducer in disparate cell functions, including changes in proliferation rate, cellular shape, attachment and motility, and regulation of proinflammatory activation [29].

Bioinformatics study results are also in line with the results of burn repair in mice by inducing EGF expression by western blot analysis and EGF immunohistochemistry in previous studies [14].

Conclusion

There are four main compounds contained in mangosteen peel extract, which in comparative studies show anti-inflammatory, antibacterial, and antioxidant activities that play a role in the healing process of burns through signaling the interleukin 6 pathway, epidermal growth factor, and transforming growth factors beta 1, which is involved in the regulation of epithelial cell proliferation.

Acknowledgements

This study was personally funded by authors.

References

1. Adiputro D. L., H. Khotimah, M. A. Widodo, R. Romdoni, D. Sargowo (2013). Cathecins in Ethanolic Extracts of *Garcinia mangostana* Fruit Pericarp and Anti-inflammatory Effect in Atherosclerotic Rats, *Journal of Experimental and Integrative Medicine*, 3(2), 137-140.
2. Amini M., M. Kherad, D. Mehrabani, N. Azarpira, M. R. Panjehshahin, N. Tanideh (2010). Effect of Plantago Major on Burn Wound Healing in Rat, *Journal of Applied Animal Research*, 37(1), 53-56.
3. Anilkumar M. (2010). Ethnomedicinal Plants as Anti-inflammatory and Analgesic Agents, *Ethnomedicine: A Source of Complementary Therapeutics*, 2010, 267-293.
4. Chomnawang M. T., S. Surassmo, V. S. Nukoolkarn, W. Gritsanapan (2007). Effect of *Garcinia mangostana* on Inflammation Caused by Propionibacterium Acnes, *Fitoterapia*, 78(6), 401-408.
5. Dickson R. A., T. C. Fleischer, E. Ekuadzi, A. Y. Mensah, K. Annan, E. Woode (2010). Antibacterial, Antioxidant and Anti-inflammatory Properties of *Margaritaria discoidea*, a Wound Healing Remedy from Ghana, *Pharmacognosy Journal*, 2(17), 32-39.
6. Dunnill C., T. Patton, J. Brennan, J. Barrett, M. Dryden, J. Cooke, D. Leaper, N. T. Georgopoulos (2017). Reactive Oxygen Species (ROS) and Wound Healing: The Functional Role of ROS and Emerging ROS-modulating Technologies for Augmentation of the Healing Process, *International Wound Journal*, 14(1), 89-96.
7. Evacuasiyany E., H. Ratnawati, L. K. Liana, W. Widowati, M. Maesaroh, T. Mozef, C. Risdian (2014). Cytotoxic and Antioxidant Activities of Catechins in Inhibiting the Malignancy of Breast Cancer, *Oxidants and Antioxidants in Medical Science*, 3(2), 141-146.
8. Ezzat S. M., M. A. Choucry, Z. A. Kandil (2016). Antibacterial, Antioxidant, and Topical Anti-inflammatory Activities of *Bergia ammannioides*: A Wound-healing Plant, *Pharmaceutical Biology*, 54(2), 215-224.
9. Fugger L., N. Morling, K. Bendtzen, L. Ryder (1989). IL-6 Gene Polymorphism in Rheumatoid Arthritis, Pauciarticular Juvenile Rheumatoid Arthritis, Systemic Lupus Erythematosus, and in Healthy Danes, *Journal of Immunogenetics*, 16(6), 461-465.
10. Gangwar M., M. K. Gautam, A. K. Sharma, Y. B. Tripathi, R. K. Goel, G. Nath (2014). Antioxidant Capacity and Radical Scavenging Effect of Polyphenol Rich *Mallotus philippinensis* Fruit Extract on Human Erythrocytes: An *in vitro* Study, *The Scientific World Journal*, 2014, 279451, doi: 10.1155/2014/279451.

11. Ghosh A. K., Y. Mori, E. Dowling, J. Varga (2007). Trichostatin A Blocks TGF-beta-induced Collagen Gene Expression in Skin Fibroblasts: Involvement of Sp1, Biochemical and Biophysical Research Communication, 354(2), 420-426.
12. Gold L. I., J. J. Sung, J. W. Siebert, M. T. Longaker (1997). Type I (RI) and Type II (RII) Receptors for Transforming Growth Factor-beta Isoforms Are Expressed Subsequent to Transforming Growth Factor-beta Ligands during Excisional Wound Repair, The American Journal of Pathology, 150(1), 209-222.
13. Gondokesumo M. E., B. Pardjianto, K. Handono, S. B. Sumitro, W. Widowati, O. Wismandanu, H. Goenawan, T. Hartady, A. M. Rosdianto, R. Lesmana, N. Wathoni, U. Supratman (2019). *Garcinia mangostana* Extract Enhances Skin Epithelialization in Rat Induced Burn Injury, Pakistan Veterinary Journal, 39(3), 365-370.
14. Gondokesumo M. E., B. Pardjianto, S. B. Sumitro, W. Widowati, K. Handono (2019). Xanthones Analysis and Antioxidant Activity Analysis (Applying ESR) of Six Different Maturity Levels of Mangosteen Rind Extract (*Garcinia mangostana* Linn.), Pharmacognosy Journal, 11(2), 369-373.
15. Grossman R. M., J. Krueger, D. Yourish, A. Granelli-Piperno, D. P. Murphy, L. T. May, T. S. Kupper, P. B. Sehgal, A. B. Gottlieb (1989). Interleukin 6 Is Expressed in High Levels in Psoriatic Skin and Stimulates Proliferation of Cultured Human Keratinocytes, Proceedings of the National Academy of Sciences of the United States of America, 86(16), 6367-6371.
16. Hazrati M., D. Mehrabani, A. Japoni, H. Montasery, N. Azarpira, A. R. Hamidian-Shirazi, N. Tanideh (2010). Effect of Honey on Healing of *Pseudomonas aeruginosa* Infected Burn Wounds in Rat, Journal of Applied Animal Research, 37(2), 161-165.
17. Hosseini S. V., H. Niknahad, N. Fakhar, A. Rezaianzadeh, D. Mehrabani (2011). The healing Effect of Mixture of Honey, Putty, Vitriol and Olive Oil in *Pseudomonas aeruginosa* Infected Burns in Experimental Rat Model, Asian Journal of Animal and Veterinary Advances, 6(6), 572-579.
18. Iordanov M. S., R. J. Choi, O. P. Ryabinina, T. H. Dinh, R. K. Bright, B. E. Magun (2002). The UV (Ribotoxic) Stress Response of Human Keratinocytes Involves the Unexpected Uncoupling of the Ras-extracellular Signal-regulated Kinase Signaling Cascade from the Activated Epidermal Growth Factor Receptor, Molecular and Cellular Biology, 22(15), 5380-5394.
19. Iwalewa E. O., L. J. McGaw, V. Naidoo, J. N. Eloff (2007). Inflammation: The Foundation of Diseases and Disorders. A Review of Phytomedicines of South African Origin Used to Treat Pain and Inflammatory Conditions, African Journal of Biotechnology, 6(25), 2868-2885.
20. Janardhanan S., J. Mahendra, A. S. S. Girija, L. Mahendra, V. Priyadharsini (2017). Antimicrobial Effects of *Garcinia mangostana* on Cariogenic Microorganisms, Journal of Clinical and Diagnostic Research, 11(1), ZC19-ZC22.
21. Jung H. A., B. N. Su, W. J. Keller, R. G. Mehta, A. D. Kinghorn (2006). Antioxidant Xanthones from the Pericarp of *Garcinia mangostana* (Mangosteen), Journal of Agricultural and Food Chemistry, 54(6), 2077-2082.
22. Koch A. E., L. B. Kronfeld-Harrington, Z. Szekanecz, M. M. Cho, G. K. Haines, L. A. Harlow, R. M. Strieter, S. L. Kunkel, M. C. Massa, W. G. Barr (1993). *In situ* Expression of Cytokines and Cellular Adhesion Molecules in the Skin of Patients with Systemic Sclerosis. Their Role in Early and Late Disease, Pathobiology, 61(5-6), 239-246.
23. Lijnen H. R., E. Maquoi, D. Demeulemeester, B. V. Hoef, D. Collen (2002). Modulation of Fibrinolytic and Gelatinolytic Activity during Adipose Tissue Development in a Mouse Model of Nutritionally Induced Obesity, Thrombosis and Haemostasis, 88, 345-353.

24. Lockett-Chastain L. R., R. M. Gallucci (2009). Interleukin (IL)-6 Modulates Transforming Growth Factor- β Expression in Skin and Dermal Fibroblasts from IL-6-deficient Mice, *British Journal of Dermatology*, 161(2), 237-248.
25. Malo N., J. A. Hanley, S. Cerquozzi, J. Pelletier, R. Nadon (2006). Statistical Practice in High-throughput Screening Data Analysis, *Nature Biotechnology*, 24(2), 167-75.
26. McCartney-Francis N. L., S. M. Wahl (1994). Transforming Growth Factor Beta: A Matter of Life and Death, *Journal of Leukocyte Biology*, 55(3), 401-409.
27. Moongkarndi P., N. Kosem, S. Kaslungka, O. Luanratana, N. Pongpan, N. Neungton (2004). Antiproliferation, Antioxidation and Induction of Apoptosis by *Garcinia mangostana* (mangosteen) on SKBR3 Human Breast Cancer Cell Line, *Journal of Ethnopharmacology*, 90(1), 161-166.
28. Paquet P., G. E. Pierard (1996). Interleukin-6 and the Skin, *International Archives of Allergy and Immunology*, 109(4), 308-317.
29. Pastore S., F. Mascia, V. Mariani, G. Girolomoni (2008). The Epidermal Growth Factor Receptor System in Skin Repair and Inflammation, *Journal of Investigative Dermatology*, 128(6), 1365-1374.
30. Pedraza-Chaverri J., N. Cárdenas-Rodríguez, M. Orozco-Ibarra, J. M. Pérez-Rojas (2008). Medicinal Properties of Mangosteen (*Garcinia mangostana*), *Food and Chemical Toxicology*, 46(10), 3227-3239.
31. Pei J., N. Yin, X. Ma, L. Lai (2014). Systems Biology Brings New Dimensions for Structure-based Drug Design, *Journal of American Chemistry Society*, 136(33), 11556-11565.
32. Pereira R. F., C. C. Barrias, P. L. Granja, P. J. Bartolo (2013). Advanced Biofabrication Strategies for Skin Regeneration and Repair, *Nanomedicine*, 8(4), 603-621.
33. Rubis B. A., D. Danikas, M. Neumeister, W. G. Williams, H. Suchy, S. M. Milner (2002). The Use of Split-thickness Dermal Grafts to Resurface Full Thickness Skin Defects, *Burns*, 28(8), 752-759.
34. Sawamura D., X. Meng, S. Ina, M. Sato, K. Tamai, K. Hanada, I. Hashimoto (1998). Induction of Keratinocyte Proliferation and Lymphocytic Infiltration by *in vivo* Introduction of the IL-6 Gene into Keratinocytes and Possibility of Keratinocyte Gene Therapy for Inflammatory Skin Diseases Using IL-6 Mutant Genes, *Journal of Immunology*, 161(10), 5633-5639.
35. Sehgal P. B. (1990). Interleukin-6: Molecular Pathophysiology, *Journal of Investigative Dermatology*, 94(6), S2-S6.
36. Szklarczyk D., A. Franceschini, S. Wyder, K. Forslund, D. Heller, J. Huerta-Cepas, M. Simonovic, A. Roth, A. Santos, K. P. Tsafou, M. Kuhn (2014). STRING v10: Protein-protein Interaction Networks, Integrated over the Tree of Life, *Nucleic Acids Research*, 43(D1), D447-D452.
37. Tanideh N., P. Rokhsari, D. Mehrabani, S. M Samani, F. S. Sarvestani, M. J. Ashraf, O. K. Hosseinabadi, S. Shamsian, N. Ahmadi (2014). The Healing Effect of Licorice on *Pseudomonas aeruginosa* Infected Burn Wounds in Experimental Rat Model, *World Journal of Plastic Surgery*, 3(2), 99-106.
38. Thiem B., O. Grosslinka (2003). Antimicrobial Activity of *Rubus chamaemorus* Leaves, *Fitoterapia*, 75(1), 93-95.
39. Tjahjani S., W. Widowati (2013). The Potency of Xanthones as Antioxidant and Antimalarial, and Their Synergism with Artemisinin *in vitro*, *Journal of Indonesian Medecal Association*, 63(3), 95-99 (in Indonesian).
40. Turksen K., T. Kupper, L. Degenstein, I. Williams, E. Fuchs (1992). Interleukin 6: Insights to Its Function in Skin by Overexpression in Transgenic Mice, *Proceedings of the National Academy of Sciences of the United States of America*, 89(11), 5068-5072.

41. Vaughan M. B., E. W. Howard, J. J. Tomasek (2000). Transforming Growth Factor-beta1 Promotes the Morphological and Functional Differentiation of the Myofibroblast, *Experimental Cell Research*, 257(1), 180-189.
42. Wang X. J., G. Han, P. Owens, Y. Siddiqui, A. G. Li (2006). Role of TGF Beta-mediated Inflammation in Cutaneous Wound Healing, *Journal of Investigative Dermatology Symposium Proceedings*, 11(1), 112-117.
43. Weecharangsan W., P. Opanasopit, M. Sukma, T. Ngawhirunpat, U. Sotanaphun, P. Siripong (2006). Antioxidative and Neuroprotective Activities of Extracts from the Fruit Hull of Mangosteen (*Garcinia mangostana* Linn.), *Medical Principles and Practice*, 15, 281-287.
44. Widowati W., L. Darsono, J. Suherman, N. Fauziah, M. Maesaroh, P. P. Erawijantari (2016). Anti-inflammatory Effect of Mangosteen (*Garcinia mangostana* L.) Peel Extract and Its Compounds in LPS-induced RAW264.7 Cells, *National Product Sciences*, 22(3), 147-153.
45. Yang K., H. Bai, Q. Ouyang, L. Lai, C. Tang (2008). Finding Multiple Target Optimal Intervention in Disease-related Molecular Network, *Molecular Systems Biology*, 4, 228, doi: 10.1038/msb.2008.60.

Marisca Evalina Gondokesumo, Ph.D.

E-mails: marisca@staff.ubaya.ac.id



Marisca Evalina Gondokesumo studied at Biomedical Sciences Doctoral Program Faculty of Medicine, Brawijaya University, Malang, East Java, Indonesia. She working as a lecturer at the Faculty of Pharmacy, University of Surabaya (UBAYA), Surabaya, East Java, Indonesia. Her latest publication was Microstructural Characterization of the *Garcinia mangostana* Fruit at Different Maturity Level; Xanthones Analysis and Antioxidant Activity Analysis (Applying ESR) of Six Different Maturity Levels of Mangosteen Rind Extract (*Garcinia mangostana* Linn.)

Prof. Sutiman Bambang Sumitro

E-mail: sutiman@ub.ac.id



Sutiman Bambang Sumitro is a Professor in Cell and Nano Biology at the Graduate School of Biomedical and Advance Biology. He currently works at the Department of Biology, Faculty of Mathematics and Natural Sciences, Brawijaya University, Malang, East Java, Indonesia. Prof. Sutiman researches Cell and Nano Biology. His current project is Research based on Herbs Exploration and Use of Animal Models: Nature Materials towards Supporting Evidence-based Medicine. Prof. Sutiman is also giving lectures and supervises doctoral students at the Faculty of Medicine Brawijaya University.

Prof. Kusworini HandonoE-mails: rinihandono.fk@ub.ac.id, dr.kusworini@gmail.com

Kusworini Handono is a Professor in Clinical Pathology and Immunology, Faculty of Medicine, Brawijaya University, Malang, East Java, Indonesia. She is a Head of the Research Centre of Autoimmune, Rheumatic, and Allergy. Her interest is in autoimmunity, especially systemic lupus erythematosus. Her research is focused on Vitamin D, herbal, and vaccine, which has led to the development of a trial kinoid vaccine. Prof. Kusworini is also giving lectures and supervises doctoral students at the Faculty of Medicine Brawijaya University, and serve patients at the Dr. Saiful Anwar General Hospital.

Prof. Bambang PardjiantoE-mail: bambangplast@yahoo.com

Bambang Pardjianto is a Professor on Plastic Surgery, Department of Plastic Reconstructive and Aesthetic Surgery, Dr. Saiful Anwar General Hospital, Malang, East Java, Indonesia. Prof. Bambang is a lecturer in the Plastic Surgery Laboratory and Biomedical Sciences Doctoral Program Faculty of Medicine, Brawijaya University, Malang, East Java, Indonesia. His research is focused on plastic surgery, plastic reconstructive and aesthetic surgery.

Dr. Ir. Wahyu Widowati, M.Sc.Email: wahyu_w60@yahoo.com

Wahyu Widowati is a lecturer and a researcher at the Faculty of Medicine, Maranatha University, Bandung, West Java, Indonesia. Her research focuses on the potential of secretome and its cells from Wharton's Jelly stem cells for cancer, osteoarthritis therapy. She currently became a President Director of Aretha Medika Utama, Biomolecular & Biomedical Research Center, Bandung, West Java, Indonesia, since 2013.

Didik Huswo Utomo, Ph.D. StudentE-mail: didik.huswo@gmail.com

Didik Huswo Utomo is a Bioinformatician who is interested in natural product research. He received his B.Sc. degree at the Biology Department of the Brawijaya University, and his M.Sc. degree at the Bandung Institute of Technology, Indonesia with concern in Bioinformatics. Currently, he is a Director of the Bioinformatics Service Company (INBIO) and a Ph.D. student at Nagoya University, Japan.



© 2020 by the authors. Licensee Institute of Biophysics and Biomedical Engineering, Bulgarian Academy of Sciences. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau PEER REVIEW

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : A Computational Study to Predict Wound Healing Agents from the Peel of the Mangosteen (*Garcinia mangostana* L.) Extract

Jumlah Penulis : 6

Nama-nama Penulis : Marisca Evalina Gondokesumo, Sutiman Bambang Sumitro, Kusworini Handono, Bambang Pardjianto, **Wahyu Widowati**, Didik Huswo Utomo

Status Penulis : ~~Penulis Pertama~~ / Penulis ke 5 / ~~Penulis Korespondensi **~~)

Identitas Jurnal Ilmiah : a. Nama jurnal : International Journal Bioautomation
 b. Nomor ISSN : 1314-2321, 1314-1902
 c. Vol., No., Bulan, Tahun : Vol. 24, No. 3, 2020
 d. Penerbit : Bulgarian Academy of Sciences
 e. DOI Artikel (jika ada) : 10.7546/ijba.2020.24.3.000607
 f. Alamat Web Jurnal : http://biomed.bas.bg/bioautomation/
 g. Terindeks di : Scopus Q3, SJR 0.178

Kategori Publikasi Jurnal Ilmiah: ~~Jurnal Ilmiah Internasional~~ / Internasional Bereputasi **) Jurnal Ilmiah Nasional Terakreditasi Jurnal Ilmiah Nasional / Nasional terindeks ***)

HASIL PENILAIAN (Peer Review) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi	Nasional Terakreditasi	Nasional ***)	
		<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,7
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			11,6
c.	Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12			11,8
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			11,7
	Total	40			38,8

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur... Paper ditulis dengan baik, unsur lengkap, hasil sesuai judul penelitian, runtun mengikuti kaidah penulisan karya ilmiah internasional bereputasi.
- b. Ruang lingkup & kedalaman pembahasan... Data penelitian berupa penelitian in silico mencari senyawa aktif pada ekstrak kulit manggis yang memiliki aktivitas wound healing.
- c. Kecukupan & kemutakhiran data serta metodologi... Sumber pustaka mutakhir, metode in silico menunjukkan bahwa senyawa aktif memiliki aktivitas antioksidan, anti-inflamatory, antibakteri yaitu (methxanthone, Garcinone-E, γ -mangostin dan bartanin
- d. Kelengkapan unsur dan kualitas penerbitan... Int J Bioautomation terindeks scopus Q3, SJR 0,18. Penerbit Institute of Biophysics and Biomedical Engineering at the Bulgarian Academy of Sciences
- e. Indikasi plagiasi

Similarity Index 15%. Tidak terdapat indikasi plagiarisme atau self-plagiarism

f. Kesesuaian bidang ilmu

Paper bidang biomedik, biokimia sesuai dengan bidang ilmu penulis.

REVIEWER 1



(Prof. Dr. Chrismis Novalinda Ginting, M.Kes)

NIK : 0115127801

UNIVERSITAS PRIMA INDONESIA

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau *PEER REVIEW*

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : A Computational Study to Predict Wound Healing Agents from the Peel of the Mangosteen (*Garcinia mangostana* L.) Extract

Jumlah Penulis : 6

Nama-nama Penulis : Marisca Evalina Gondokesumo, Sutiman Bambang Sumitro, Kusworini Handono, Bambang Pardjianto, **Wahyu Widowati**, Didik Huswo Utomo

Status Penulis : ~~Penulis Pertama~~ / Penulis ke 5 / ~~Penulis Korespondensi~~ (**)

Identitas Jurnal Ilmiah : a. Nama jurnal : International Journal Bioautomation
 b. Nomor ISSN : 1314-2321, 1314-1902
 c. Vol., No., Bulan, Tahun : Vol. 24, No. 3, 2020
 d. Penerbit : Bulgarian Academy of Sciences
 e. DOI Artikel (jika ada) : 10.7546/ijba.2020.24.3.000607
 f. Alamat Web Jurnal : <http://biomed.bas.bg/bioautomation/>
 g. Terindeks di : Scopus Q3, SJR 0.178

Kategori Publikasi Jurnal Ilmiah: Jurnal Ilmiah Internasional / Internasional Bereputasi (**)
 (beri tanda \checkmark yang dipilih) Jurnal Ilmiah Nasional Terakreditasi
 Jurnal Ilmiah Nasional / Nasional terindeks (***)

HASIL PENILAIAN (*Peer Review*) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi	Nasional Terakreditasi	Nasional (***)	
		<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,6
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			11,0
c.	Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12			11,7
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			11,5
	Total	40			36,8

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur.....
Kelengkapan unsur isi artikel sudah cukup lengkap dan terdapat kesesuaian antara unsur dan banyu
- b. Ruang lingkup & kedalaman pembahasan.....
Cakupan pembahasan sudah cukup mendalam dengan tambahan teori yang jelas dan seluas
- c. Kecukupan & kemutakhiran data serta metodologi.....
Kemutakhiran data sudah terpenuhi dan metode yang dipakai sudah sesuai dengan tujuan penelitian
- d. Kelengkapan unsur dan kualitas penerbit.....
Kualitas penerbit sudah memenuhi kriteria dan kualitas jurnal masuk kategori yang baik
- e. Indikasi plagiasi

Belum ditemukan adanya unsur Indikasi plagiasi

f. Kesesuaian bidang ilmu

Jurnal ini sudah sesuai dengan bidang ilmu yang diteliti

REVIEWER 2



(Prof. Dr. Ermi Girsang, M. Kes)

NIK : 0117057501

UNIVERSITAS PRIMA INDONESIA

LEMBAR HASIL PENILAIAN
SEJAWAT SEBIDANG atau *PEER REVIEW*

KARYA ILMIAH : JURNAL ILMIAH

Judul Karya Ilmiah (Artikel) : A Computational Study to Predict Wound Healing Agents from the Peel of the Mangosteen (*Garcinia mangostana* L.) Extract

Jumlah Penulis : 6

Nama-nama Penulis : Marisca Evalina Gondokesumo, Sutiman Bambang Sumitro, Kusworini Handono, Bambang Pardjianto, **Wahyu Widowati**, Didik Huswo Utomo

Status Penulis : ~~Penulis Pertama~~ / Penulis ke 5 / ~~Penulis Korespondensi~~ (**)

Identitas Jurnal Ilmiah : a. Nama jurnal : International Journal Bioautomation
b. Nomor ISSN : 1314-2321, 1314-1902
c. Vol., No., Bulan, Tahun : Vol. 24, No. 3, 2020
d. Penerbit : Bulgarian Academy of Sciences
e. DOI Artikel (jika ada) : 10.7546/ijba.2020.24.3.000607
f. Alamat Web Jurnal : <http://biomed.bas.bg/bioautomation/>
g. Terindeks di : Scopus Q3, SJR 0.178

Kategori Publikasi Jurnal Ilmiah : Jurnal Ilmiah Internasional / Internasional Bereputasi (**)
(beri tanda \checkmark yang dipilih)
 Jurnal Ilmiah Nasional Terakreditasi
 Jurnal Ilmiah Nasional / Nasional terindeks (***)

HASIL PENILAIAN (*Peer Review*) :

No	Komponen Yang Dinilai	Nilai Maksimal JURNAL ILMIAH			Nilai Akhir Yang Diperoleh *)
		Internasional / Bereputasi	Nasional Terakreditasi	Nasional (***)	
		<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	
a.	Kelengkapan unsur isi karya (10%)	4			3,65
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12			11,6
c.	Kecukupan dan kemuakhiran data/ informasi dan metodologi (30%)	12			11,8
d.	Kelengkapan unsur dan kualitas penerbitan (30%)	12			11,6
	Total	40			38,6

Catatan Penilaian ARTIKEL oleh Reviewer :

- a. Kelengkapan dan kesesuaian unsur.....
Paper ditulis dengan baik, unsur lengkap, hasil sesuai judul penelitian, runtun mengikuti bab dan penulisan karya ilmiah internasional bereputasi
-
kelengkapan unsur isi artikel sudah cukup lengkap dan terdapat kesesuaian antara unsur dan isinya
- b. Ruang lingkup & kedalaman pembahasan.....
Data penelitian berupa penelitian *in silico*, mencari senyawa aktif pada ekstrak kulit manggis yang memiliki aktivitas wound healing.

Cakupan pembahasan sudah cukup mendalam dengan tambahan teori yang jelas dan sesuai
c. Kecukupan & kemutakhiran data serta metodologi. Sumber pustaka mutakhir, metode *in silico* menunjukkan bahwa senyawa aktif memiliki aktivitas antioksidan, antiinflamatory, antibakteri yaitu *Alneathxanthone*, *Garcinone-E*, *8-triangolin* dan *Bartarin*.

Kemutakhiran data sudah terpenuhi dan metode yang dipakai sudah sesuai dg tujuan penelitian
d. Kelengkapan unsur dan kualitas penerbit

Int J Bioautomation terindeks scopus Q4 SJR 0,18. Penerbit Institute of Biophysics and Biomedical Engineering at the Bulgarian Academy of Sciences.

Kualitas penerbit sudah memenuhi standar dan levelitas jurnal masuk kategori yang baik.

e. Indikasi plagiasi

Similarity index 15%. Tidak terdapat indikasi plagiarisme atau self-plagiarisme

Belum ditemukan adanya indikasi plagiasi.

f. Kesesuaian bidang ilmu

Paper bidang biomedik, biokimia sesuai dengan bidang ilmu penulis.


Jurnal ini sudah sesuai dengan bidang ilmu yang diteliti.

Medan,
Reviewer 2



(Prof. Dr. Ermi Girsang, M.Kes)
NIK : 0117057501
UNIVERSITAS PRIMA INDONESIA

Medan,
Reviewer 1



(Prof. Dr. Chrismis Novalinda Ginting, M.Kes)
NIK : 0115127801
UNIVERSITAS PRIMA INDONESIA