

# HAYATI

p-ISSN: 1978-3019  
e-ISSN: 2086-4094

## Journal of Biosciences

[ABOUT](#) [CURRENT](#) [ARCHIVES](#) [EDITORS](#) [SUBMISSIONS](#) [REGISTER](#)

[CONTACT](#) [SEARCH](#)

[Home](#) / [Editorial Team](#)

## Editorial Team

### Chief Editor

[Dr. Iman Rusmana](#), Bogor Agricultural University, Indonesia

### Managing Editor

[Dr. Kanthi Arum Widayati](#), Bogor Agricultural University, Indonesia

### Editorial Board

[Prof. Koji Nakamura](#), Kanazawa University, Japan

[Prof. Yasumasa Bessho](#), Nara Institute of Science and Technology, Japan

[Prof. Wei Zhang](#), Flinders University, Australia

[Prof. Lisdar Idwan Sudirman](#), Bogor Agricultural University, Indonesia

[Prof. Michel Raymond](#), Institut des Sciences de l'Evolution, France

[Prof. Dr. dr. Sri Budiarti](#), Bogor Agricultural University, Indonesia

[Prof. Youji Nitta](#), Fukushima University, Japan

[Prof. Dave B. Nedwell](#), University of Essex, United Kingdom

[Prof. Aris Tri Wahyudi](#), Bogor Agricultural University, Indonesia

[Assoc. Prof. Dr. Yaya Rukayadi](#), Department of Food Science. Faculty of Food Science and Technology. Universiti Putra Malaysia

[Dr. Patrick O'Connor](#), The University of Adelaide, Australia

[Dr. Ardiansyah](#), Universitas Bakrie, Indonesia

[Dr. Rika Indri Astuti](#), Bogor Agricultural University, Indonesia

[Dr. Mashuri Waite](#), University of Hawaii, USA

[Dr. Atsushi Ido](#), Ehime University, Japan

[Dr. Sastia Prama Putri](#), Osaka University, Japan

[Dr. Bambang Retnoaji](#), Gadjah Mada University, Indonesia

[Dr. Malcolm Wegener](#), The University of Queensland, Australia

[Dr. Miftahudin](#), Bogor Agricultural University, Indonesia

[Dr. Dodi Safari](#), Eijkman Institute, Indonesia

[Dr. Emmanuel Paradis](#), Institut des Sciences de l'Evolution de Montpellier, France

[Dr. Erin Phelps Riley](#), San Diego State University, USA

[Dr. Rebecca Johnson](#), Columbia University, USA

[Dr. Takuya Imamura](#), Kyushu University, Japan

[Prof. Dr. Dyah Perwitasari](#), Bogor Agricultural University, Indonesia

[Dr. Tsukasa Sanosaka](#), Keio University, Japan

[Dr. Ocky Karna Radjasa](#), Diponegoro University, Indonesia

[Dr. Tukirin Partomihardjo](#), Indonesian Research Center for Biology (LIPI), Indonesia

[Dr. Yamato Tsuji](#), Kyoto University, Japan

[Dr. Darjono](#), Indonesian Research Center for Biology (LIPI), Indonesia

[Prof. Dr. Hamim](#), Bogor Agricultural University, Indonesia


#### Technical Editor

[Dr. Puji Rianti](#), Bogor Agricultural University, Indonesia

[Mafrikhul Muttaqin, MSi](#), Bogor Agricultural University, Indonesia

#### Online Manager

[Jepri Agung Priyanto, MSi](#), Bogor Agricultural University, Indonesia



#### JOURNAL INFORMATION

[About The Journal](#)

[Editorial Team](#)

[Focus and Scope](#)

[Publication Ethics](#)

[Privacy Statement](#)

[Author Guidelines](#)

[Register New Account](#)

[Submit or Check Status](#)

[Contact](#)



**INDEXING**



**PUBLISHER**



**GOOGLE SCHOLAR CITATIONS**

HAYATI J.Biosci : GS citation

	All	Since 2017
Citations	7785	6035
h-index	40	34
i10-index	249	193

GS citation >

**VISITOR STATISTICS**

00548895

[View Hayati J. Biosci. Stats](#)



### HJB Most Read Papers

[Photosynthesis and Transpiration Rates of Rice Cultivated Under the System of Rice Intensification and the Effects on Growth and Yield](#)

510

10 November 2016

Nurul Hidayati, . Triadiati...

[Clinical and Hematological Parameters of Captive Timor Deer \(\*Rusa timorensis\*\) under Physical Restraint](#)

388

28 June 2022

Virgilius Martin Kelake Ked...

[Antibacterial and Antioxidant Activities of Fungal Endophytes Isolated from Medicinal Plants in Simeulue Island, Aceh](#)

339

28 June 2022

Ahmad Fathoni, Muhammad Ily...

[The Development of In Vitro Culture Sterilization Method of Gametophyte Explant \*Lopholejeunea\* sp.](#)

277

1 April 2021

Anna Widyastuti, Afiatry ...

[Propolis Components and Biological Activities from Stingless Bees Collected on South Sulawesi, Indonesia](#)

272

1 January 2020

Ryo Miyata, Muhamad Sahla...



This journal is published under the terms of the <https://creativecommons.org/licenses/by-nc/4.0/>



**IPB University**  
— Bogor Indonesia —

**HAYATI Journal of Biosciences (p-ISSN: 1978-3019; e-ISSN: 2086-4094)**

Department of Biology, Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, Dramaga Campus, Bogor 16680, Indonesia

Phone/Fax: +62-251-8421258; E-mail: [hayati.jbiosci@apps.ipb.ac.id](mailto:hayati.jbiosci@apps.ipb.ac.id); URL : <http://journal.ipb.ac.id/index.php/hayati>

Platform &  
workflow by



# HAYATI

p-ISSN: 1978-3019  
e-ISSN: 2086-4094


## Journal of Biosciences

[ABOUT](#) [CURRENT](#) [ARCHIVES](#) [EDITORS](#) [SUBMISSIONS](#) [REGISTER](#)

[CONTACT](#) [SEARCH](#)

[Home](#) / [Archives](#) / Vol. 27 No. 3 (2020): July 2020

### Vol. 27 No. 3 (2020): July 2020



**Published:** 2020-07-01

#### Articles

##### [Ecological Diversity of Microbial Consortium Feces of Beef Cattle and Lignite Coal](#)

Gina Chynthia Kamarudin Puteri, Roni Ridwan, Ellin Harlia  
180



##### [Apoptotic Potential of Secretome from Interleukin-Induced Natural Killer Cells toward Breast Cancer Cell Line by Transwell Assay](#)

Wahyu Widowati, Diana Krisanti Jasaputra, [Teresa Liliana Wargasetia](#), The Fransiska Eltania, Alya Mardhotillah Azizah, Mawar Subangkit, I Nyoman Ehrich Lister, Chrismis Novalinda Ginting, Ermi Girsang, Ahmad Faried  
186



##### [Impact of Methanotrophic and N2O-reducing Bacterial Inoculation on CH4 and N2O Emissions, Paddy Growth and Bacterial Community Structure in Paddy Field](#)

Yuli Siti Fatma, Iman Rusmana, Aris Tri Wahyudi, Hamim Hamim  
197



##### [Antagonistic Activity of Fungal Endophytes Isolated from Garcinia atroviridis against Colletotrichum gloeosporioides](#)

Nur Afeeqah Binti Mohamed Zanudin, Nor'Aishah Hasan, Patahayah Binti Mansor  
209



##### [Screening of Endophytic Bacteria as Biocontrol Agents Against Bacteria Leaf Blight \(Xanthomonas oryzae\)](#)

Rashidah Abd Halim, Nor'Aishah Hasan, Kogeethavani Ramachandran  
215



### Mammalian Diversity of Gunung Ledang, Johor, Peninsular Malaysia

Faiznur Ain Ahmad Bakri, Masatoshi Yasuda, Maryati Mohamed, Adlii Ikram Sharuddin, Muhazam Syah Hambar  
221



### In silico Vaccine Design against Dengue Virus Type 2 Envelope Glycoprotein

Muhammad Adnan, Matin Nuhamunada, Lisna Hidayati, Nastiti Wijayanti  
228



### Identification of Follower Status Based on Male Proximity Score in Crested Macaque

Andre Pasetha, Lisa Michelle Danish, Dyah Perwitasari-Farajallah, Muhammad Agil, Antje Engelhardt  
241




### Anatomical Properties and Near Infrared Spectra Characteristics of Four Shorea Species from Indonesia

Danang Sudarwoko Adi, Sung-Wook Hwang, Dwi Aji Pramiasari, Yusup Amin, Hairi Cipta, Ratih Damayanti, Wahyu Dwianto, Junji Sugiyama  
247



### Genetic Relationship between Tongka Langit Bananas (*Musa troglodytarum* L.) from Galunggung and Maluku, Indonesia, Based on ITS2

Fenny Martha Dwivany, Giasintha Stefani, Agus Sutanto, Husna Nugrahapraja, Ketut Wikantika, Adriana Hiariej, Topik Hidayat, I Nyoman Rai, Nisrina Sukriandi  
258



## JOURNAL INFORMATION

[About The Journal](#)

[Editorial Team](#)

[Focus and Scope](#)

[Publication Ethics](#)

[Privacy Statement](#)

[Author Guidelines](#)

[Register New Account](#)

[Submit or Check Status](#)

[Contact](#)



**INDEXING**



**PUBLISHER**



**GOOGLE SCHOLAR CITATIONS**

HAYATI J.Biosci : GS citation

	All	Since 2017
Citations	7785	6035
h-index	40	34
i10-index	249	193

GS citation >



## VISITOR STATISTICS

00548847

[View Hayati J. Biosci. Stats](#)



### HJB Most Read Papers

[Photosynthesis and Transpiration Rates of Rice Cultivated Under the System of Rice Intensification and the Effects on Growth and Yield](#)

510

10 November 2016

Nurul Hidayati, . Triadiati...

[Clinical and Hematological Parameters of Captive Timor Deer \(\*Rusa timorensis\*\) under Physical Restraint](#)

388

28 June 2022

Virgilius Martin Kelake Ked...

[Antibacterial and Antioxidant Activities of Fungal Endophytes Isolated from Medicinal Plants in Simeulue Island, Aceh](#)

339

28 June 2022

Ahmad Fathoni, Muhammad Ily...

[The Development of In Vitro Culture Sterilization Method of Gametophyte Explant \*Lopholejeunea\* sp.](#)

277

1 April 2021

Anna Widyastuti, Afiatry ...

[Propolis Components and Biological Activities from Stingless Bees Collected on South Sulawesi, Indonesia](#)

272

1 January 2020

Ryo Miyata, Muhamad Sahla...



This journal is published under the terms of the <https://creativecommons.org/licenses/by-nc/4.0/>



**IPB University**  
— Bogor Indonesia —

**HAYATI Journal of Biosciences (p-ISSN: 1978-3019; e-ISSN: 2086-4094)**

Department of Biology, Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, Dramaga Campus, Bogor 16680, Indonesia

Phone/Fax: +62-251-8421258; E-mail: [hayati.jbiosci@apps.ip.ac.id](mailto:hayati.jbiosci@apps.ip.ac.id); URL : <http://journal.ipb.ac.id/index.php/hayati>



## Apoptotic Potential of Secretome from Interleukin-Induced Natural Killer Cells toward Breast Cancer Cell Line by Transwell Assay

Wahyu Widowati<sup>1\*</sup>, Diana Krisanti Jasaputra<sup>1</sup>, Teresa Liliana Wargasetia<sup>1</sup>, The Fransiska Eltania<sup>1</sup>, Alya Mardhotillah Azizah<sup>2</sup>, Mawar Subangkit<sup>3</sup>, I Nyoman Ehrich Lister<sup>4</sup>, Chrismis Novalinda Ginting<sup>4</sup>, Ermi Girsang<sup>4</sup>, Ahmad Faried<sup>5</sup>

<sup>1</sup>Faculty of Medicine, Maranatha Christian University, Bandung, Indonesia

<sup>2</sup>Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung, Indonesia

<sup>3</sup>Laboratory of Veterinary Pathology, Faculty of Veterinary Medicine, IPB University, Bogor, Indonesia

<sup>4</sup>Faculty of Medicine, Universitas Prima Indonesia, Medan, Indonesia

<sup>5</sup>Department of Neurosurgery and Stem Cell Working Group, Faculty of Medicine, Padjadjaran University, Bandung, Indonesia

### ARTICLE INFO

#### Article history:

Received November 15, 2019

Received in revised form June 15, 2020

Accepted June 30, 2020

#### KEYWORDS:

MCF7,  
NK cells,  
apoptosis,  
interleukin,  
NKG2D

### ABSTRACT

Breast cancer (BC) is the number one cause of deaths from cancer in women. Metastasis in BC is caused by immunosurveillance deficiency, including impairment of Natural Killer (NK) cell maturation, low NK activity, and decreasing cytotoxicity. This study was performed to improve activating receptors and cytotoxicity of NK cells using interleukin 15 (IL15) against BC cells. Human recombinant IL15 was used to induce NK cells. To evaluate the potential of IL15 in inducing NK cells, we measured the activating and inhibiting receptors (NKG2D, NKG2A), apoptotic potency of NK cells on BC cells (MCF7) using transwell assay. The IL15 inducer on the NK cell were measured NKG2D, NKG2A gene expression with quantitative polymerase chain reaction (qPCR), (GzmB) secretion using ELISA, apoptotic gene expression of MCF7 using qPCR. IL15 increased NKG2D expression 4.01-9.13%, but IL15 could not affect toward NKG2A expression on NK cells. IL15-activated NK cells, inhibited BC cells proliferation, induced apoptotic BC cells 25.89-32.19%, induced apoptotic genes of BC cells bax, p53. IL15 increase NK activating receptor (NKG2D), inhibit BC cells proliferation, induce apoptotic percentage and induce apoptotic gene expression.

## 1. Introduction

Breast cancer (BC) is one of the main causes of death in the world. In 2012, it caused around 8.2 million deaths (Ferlay *et al.* 2015). BC is the most frequently diagnosed cancer and the leading cause of cancer death among females, accounting for 23% of the total cancer cases and 14% of the cancer deaths (Tirona *et al.* 2010). Metastasis in BC is caused by deficient immunosurveillance, including impairment of Natural Killer (NK) cell maturation, low NK cell counts in peripheral blood mononuclear cells (PBMCs), significantly lower NK activity in patients with BC than in healthy individuals (Dewan *et al.* 2009), decreased cytotoxic function (Levy *et al.* 2011; Hwang *et al.* 2012), NK abnormalities (Levy *et al.* 2011), poor tumor infiltrate (Albertsson *et al.*

2003; Esendagli *et al.* 2008), low NK cell numbers in tumors due to their inefficient homing into malignant tissues (Levy *et al.* 2011), and defective expression of activating receptors such as NKG2D, overexpression of inhibitory receptors NKG2A, CD158a, CD158b (Levy *et al.* 2011).

Immunotherapy using NK cells can be used to obtain the large and sufficient numbers of functional NK cells necessary for clinical therapy. The number, purity and state of NK cell proliferation and activation are key factors in immunotherapy (Cheng *et al.* 2013). NK cells are known as necessary effectors in suppressing cancer proliferation (Kelly *et al.* 2003; Smyth *et al.* 2005). Once NK cells recognize target cells such as cancer cells, they form an immunological synapse, and the secretory granules fuse with the presynaptic membrane and release perforin (Prf) and granzyme (Gzm) into the synaptic cleft. Released Prf provides transmembrane pores on the target cell

\* Corresponding Author

E-mail Address: wahyu\_w60@yahoo.com

and enables Gzm to diffuse into the cell. Gzm then initiates the apoptosis of the target cells, and the NK cells detach from the dying cells and can interact with other target cells to accomplish serial killing (Harada *et al.* 2017). Therefore, the focus of recent cancer therapy has been to promote and develop NK cells as drugs (Hwang *et al.* 2012), using NK effectors such as cytokines (Levy *et al.* 2011). This study was conducted to evaluate the effect of inducing Interleukin 15 (IL15) on NK cells toward the following: i) improvement of NK cell activating receptors (NKG2D), ii) inducement cytotoxic towards BC cells, iii) apoptotic inducer of NK cells, iv) apoptotic genes expression of BC cells, vii) morphology and density of MCF7 cells by transwell of NK and BC cells.

## 2. Materials and Methods

### 2.1. Culturing Breast Cancer and NK Cells

Breast cancer (MCF7) cells (ATCC® HTB-22™) obtained from Aretha Medika Utama, Biomolecular and Biomedical Research Center, Bandung, Indonesia were cultured in RPMI 1640 (Gibco, 11875093) supplemented with 10% FBS (Gibco, 26140079) and 1% antibiotic/antimycotic. Meanwhile, NK92 cells (ATCC® CRL-2407™) from Aretha Medika Utama, Biomolecular and Biomedical Research Center, Bandung, Indonesia were cultured in MEM-without nucleosides (Gibco, 12561056) supplemented with 12.5% FBS, 12.5% Horse Serum (Gibco, 16050122), Sodium bicarbonate (Merck, 106329), Myo-inositol (Sigma Aldrich, 17508), Folic Acid (Sigma Aldrich, F8758), IL-2 (BioLegend, 589106), 2-Mercaptoethanol (Gibco, 1628448), and 1% antibiotic/antimycotic (Gibco, 1772653). Cell cultures were maintained in T-75 flasks, at 37°C, 5% CO<sub>2</sub>. Culture mediums were changed every three days and sub-cultured at 80% confluence (Widowati *et al.* 2018; Widowati *et al.* 2019).

### 2.2. Determination of NKG2A and NKG2D Gene Expression

A suspension of NK92 cells (1 x 10<sup>6</sup> cells) were seeded onto T-25 Flask (TPP, 90026) and incubated for 24 hours at 37°C, 5% CO<sub>2</sub>. Afterwards, the cells were induced by human recombinant IL15 (BioLegend, 715902) to reach final concentration at 5 ng/ml and 10 ng/ml and incubated for 24 hours at 37°C, 5% CO<sub>2</sub>. The RNA isolation of NK92 was performed according to the protocol of RNA Isolation Kit (Bio-Rad, 732-6820). The concentration and purity of RNA of each sample was determined at 260/280 nm. The cDNA synthesis was performed using cDNA synthesis kit (Bio-rad, 170-8841) (Widowati *et al.* 2018; Afifah *et al.* 2019; Widowati *et al.* 2019). Primer sequences were used can be seen at Table 1.

### 2.3. Cytotoxic Assay

MCF7 and NK92 cells were seeded at a ratio 1:1 onto transwell 24 well-plate (Corning, 3396). MCF7 were seeded on the first day allowed to attach to the bottom chamber. Next day, NK92 induced by IL15 (0, 5, 10 ng/ml) were seeded to the upper chamber. The plate was incubated for 24 hours at 37°C, 5% CO<sub>2</sub>. Hereinafter, the upper chamber and the medium were discarded. The cells were washed one time using PBS 1x. After that, 500 µl Presto Blue 1x (Invitrogen, A13262) were added to each well and incubated for 1 hour at 37°C, 5% CO<sub>2</sub>. The solution was transferred to 96-well plate and the absorbance was read at 570 and 600 nm using spectrophotometer (Multiskan GO, ThermoScientific) (Widowati *et al.* 2018; Widowati *et al.* 2019).

### 2.4. Apoptosis Analysis Assay by Flow Cytometry

MCF7 and NK92 cells were seeded at a ratio 1:1 onto 6 well/plate (Corning, 3412). On the first day, MCF7 were seeded at the bottom chamber and

Table 1. Activator, inhibitor receptor genes of NK Cells, RT PCR protocols

Gene symbols	Primer sequence (5' to 3')	Product size (bp)	Annealing (°C)	Cycle	References
	Upper strand:sense				
	Lower strand:antisense				
NKG2A	5'-CCAGAGAAGCTCATTGTTGG-3' 5'-CCAATCCATGAGGATGGT-3'	168	51	40x	Gen Bank:AF461812.1
NKG2D	5'-CTGGGAGATGAGTGAATTCATA-3' 5'-GACTTCACCTTAAGTAAATC-3'	417	51	40x	Gen Bank:AF461811.1
GAPDH	5'-GGGCTGCTTTTAACTCTGGT-3' 5'-TGGCAGTTTTTCTAGACGG-3'	702	51	40x	Sadeghi <i>et al.</i> 2015

plates were incubated for 24 hours at 37°C, 5% CO<sub>2</sub>. On the next day, NK92 cells were added to the upper chamber. The NK92 cells were induced by IL15 (5 ng/ml and 10 ng/ml). The uninduced NK92 cells and MCF7 without transwell were also done as a control. The plate was then incubated for 24 hours at 37°C, 5% CO<sub>2</sub>. Subsequently, MCF7 at the bottom chamber were collected and washed two times by FACS buffer (2% FBS in PBS 1x). Pellets were resuspended in 100 µl FACS buffer and 5 µl Annexin V (BioLegend, Part 79998), 10 µl PI (BioLegend, Part 79997) were added. The cells were incubated for 30 minutes in 4°C dark room. Following the 30 minutes incubation, samples were added 400 µl annexin V-binding buffer (BioLegend, 640194) and immediately analyzed by flow cytometry (MACSQuant™ Analyzer 10, Miltenyi Biotec) (Widowati *et al.* 2018; Widowati *et al.* 2019).

### 2.5. Quantification of Granzyme-B Level

The quantitative determination of GzmB in the transwell-treatment, conditioned medium from NK cells from upper chamber and conditioned medium from MCF7 was performed using ELISA Kit Human Granzyme-B (Human ELISA KIT, ElabSci E-EL-H1617) followed the manufacturer protocol. Absorbance was read at 450 nm using spectrophotometer (Safta *et al.* 2015; Widowati *et al.* 2018, 2020).

### 2.6. Determination of Apoptosis Gene Expression

MCF7 cells had been transwelled with induced NK92 cells (IL15 with various concentration 0, 5, 10 ng/ml) for 24 h. Afterward, microscopic analysis was conducted under inverted phase-contrast microscope. Subsequently the MCF7 cells were collected to isolate the RNA. The RNA isolation of MCF7 was performed based on the protocol of RNA Isolation Kit (Bio-Rad, 732-6820). The concentration and purity of RNA of each sample was determined at 260/280 nm. The cDNA synthesis was performed using cDNA synthesis

kit (Bio-rad, 170-8841) (Widowati *et al.* 2018; Afifah *et al.* 2019; Widowati *et al.* 2019). Primer sequences can be seen at Table 2.

### 2.7. Statistical Analysis

All the data are presented as the mean ± standard deviation. The data were analyzed using one-way analysis of variance (ANOVA) followed by Tukey's post-hoc test for multiple comparisons with p-values less than 0.05 were considered significant.

## 3. Results

### 3.1. Effect of IL15 toward NK Cell Characteristics

To determine the effect of human recombinant IL15 toward NK receptors, we evaluated the NK receptors including gene expression of NKG2D and NKG2A (Figure 1). The data showed that IL15 (5, 10 ng/ml) significantly up-regulated NKG2D, meanwhile IL15 did not significantly down-regulate NKG2A.

Based on the data (Figure 1) shows that IL15 induced up-regulation of NKG2D gene expression, the highest NKG2D expression was IL15 (5 ng/ml) higher concentration of IL15 (10 ng/ml) lowered NKG2D expression. IL15 both concentrations (5, 10 ng/ml) didn't affect NKG2A expression of NK cells.

### 3.2. Cytotoxic Activity of IL15-Induced NK toward BC Cells

NK cells are involved in the elimination of tumor cells. To determine the effect of human recombinant IL15 on NK cells toward cytotoxic activity of BC cells. We determine the effect of IL15-induced NK toward MCF7 cells by transwell method. NK cells treated with IL15 at concentrations of 0, 5, 10 ng/ml (IL15-induced NK cells) against BC cell proliferation, we evaluated the inhibition of MCF7 proliferation (Figure 2).

Figure 2 shows that NK cells uninduced with IL15 had cytotoxic activity against MCF7 cells approximately 35.20% inhibitory growth cells. Both concentrations 5,

Table 2. Apoptotic primer for MCF7 cells, RT PCR protocols

Gene symbols	Primer sequence (5' to 3')	Product size (bp)	Annealing (°C)	Cycle	References
	Upper strand:sense				
	Lower strand:antisense				
p53	5'-AGAGTCTATAGCCCCACCCC-3' 5'-GCTCGACGCTAGGATCTGAC-3'	97	58	40x	Behbahani 2014; Widowati <i>et al.</i> 2019
bax	5'-TGCTTCAGGGTTTCATCCAG-3' 5'-GGCGGAATCATCCTCTG-3'	169	58	40x	Wang <i>et al.</i> 2009; Widowati <i>et al.</i> 2019
bcl2	5'-GGTCATGTGTGGAGAGCG-3' 5'-GGTGCCGGTTCAGGTACTCA-3'	89	58	40x	Guan <i>et al.</i> 2018; Widowati <i>et al.</i> 2019
β-Actin	5'-TCTGGCACACACCTTCTACAATG-3' 5'-AGCACAGCTGGATAGCAACG-3'	166	85	40x	Han <i>et al.</i> 2018




Figure 1. Effect IL15 inducer on NK cells toward NKG2A and NKG2D gene expression. The data are presented as histogram among treatments, this research was conducted in triplicate for each treatment. The asterisks symbol (\*) presents significant difference between (inducing IL15 5 ng/ml, inducing IL15 10 ng/ml) and non inducing of NK cells based on Tukey HSD post hoc test ( $p < 0.05$ ). (a) The IL15 didn't affect NKG2A gene expression. (b) The NKG2D expression of higher concentration (10 ng/ml IL15) was lower than 5 ng/ml IL15




Figure 2. Effect of IL15 inducer on NK cells toward proliferation inhibition of MCF7 cells. The data are presented as histogram among treatments, this research was conducted in triplicate for each treatment. The asterisks symbol (\*) presents significant difference between (inducing IL15 5 ng/ml, inducing IL15 10 ng/ml) and non inducing of NK cells based on Tukey HSD post hoc test ( $p < 0.05$ ). Figure 2 shows that IL15 activated NK cells to inhibit MCF7 cell proliferation and increased anticancer activity NK cells against proliferation of MCF7 cells. The higher concentration (10 ng/ml) of IL15 was not significant difference compared to lower concentration of IL15 (5 ng/ml) against MCF7 cells proliferation

10 ng/ml of IL15 induced NK cells to inhibit MCF7 cells proliferation about 65.60-70.29%.

**3.3. Effect of IL15-Induced NK Cells toward Apoptosis of MCF7 Cells**

NK cells require effectors to activate NK cells; thus, this study was conducted to evaluate the effect of IL15 toward apoptosis of MCF7 cells by transwell method. To determine the effect of human recombinant IL15 on NK cells toward apoptotic percentage of MCF7 cells, we evaluated apoptotic percentage of MCF7 cells (Figures 3). We used IL15 at levels of 0 (uninduced), 5 and 10 ng/ml. The data (Figure 3) show that IL15 increased apoptotic percentage of MCF7, the highest apoptotic potency was NK induced by 5 ng/ml, higher concentration of IL15 (10 ng/ml) lower apoptotic percentage than NK induced by 5 ng/ml.


Based on the apoptotic percentage of MCF7 cells were treated by IL15-induced NK cells using transwell method (Figure 3), shows that NK cells uninduced with IL15 was very low apoptotic potency (3.98%), IL15 activated NK cells and increased apoptotic potency against MCF7 cells, the best IL15 concentration was 5 ng/ml, higher concentration of IL15 significantly lowered apoptotic percentage of MCF7 cells.

**3.4. Effect of IL15-Induced NK Cells toward Granzyme Level in Transwell of MCF7 and NK Cells**

We measured NK cells were activated using IL15 (0, 5, 10 ng/ml). The cell ratios (MCF7: NK cells=1:1) were assessed for the ability to secrete GzmB. The effects


of IL15-induced NK (IL15-NK) transwelled with MCF7 cells, which made NK released cytoplasmic granules (GzmB) can be seen in Figure 4.

The GzmB level secreted by NK cells which co-cultured with MCF7 cells using transwell method (Figure 5). NK cells uninduced by IL15 secreted GzmB in low level (77.06 pg/ml), inducing IL15 (5, 10 ng/ml) increased secretion of GzmB, the highest GzmB level was secreted by NK cells which induced by IL15 (5 ng/



Transwell of MCF7 and IL15-induced NK cells

Figure 4. Effect of IL15 inducer on NK cells toward GzmB level of transwell MCF7 and NK cells. The data are presented as histogram among treatment, this research was conducted in triplicate for each treatment. The asterisks symbol (\*) present significant differences among concentrations of inducer IL15 (0, 5, 10 ng/ml) compared to control (MCF7 non treatment) toward GzmB level



Transwell of MCF7 and IL15-induced NK cells

Figure 3. Effect IL15 inducer on NK cells toward apoptotic percentage of MCF7 cells. The data are presented as histogram among treatment, this research was conducted in triplicate for each treatment. The asterisks symbol (\*) present significant differences among concentrations of IL15 inducer (0, 5, 10 ng/ml) compared to control (MCF7 non treatment) toward apoptotic percentage of MCF7



ml), meanwhile higher IL15 concentration (10 ng/ml) lowered GzmB level.

### 3.5. Effect IL15-Induced NK Cells toward Apoptotic and Antiapoptotic Genes Expression of MCF7

This study was the continued-research to elucidate the apoptosis mechanism of IL15-induced NK cells toward MCF7 cells, in order to determine the apoptotic inducing activity of IL15-induced NK cells toward MCF cells by transwell assay, the expression of apoptotic genes was determined by

qPCR. We measured the expression of proapoptotic genes, specifically p53, b-cell cll/lymphoma 2 (bcl-2), bcl2-associated x protein (bax). The p53, bax and bcl2 genes expression of MCF7 can be seen at Figure 5.

NK cells induced by IL15 significantly increased the apoptotic genes both low and high concentrations (5, 10 ng/ml) toward p53, bax genes expression (Figure 5a and 5b) but IL15-induced NK didn't affect bcl2 expression (Figure 5c). The highest p53 expression was MCF7 treated with 5 ng/ml IL15-induced NK cells, the higher concentration




Figure 5. Effect IL15 inducer on NK cells toward p53, bax and bcl2 gene expression of MCF7 cells. The data are presented as histogram among treatments, this research was conducted in triplicate for each treatment. The asterisks symbol (\*) presents significant difference between (inducing IL15 5 ng/ml, inducing IL15 10 ng/ml) compred control (MCF7 non treatment) based on Tukey HSD post hoc test ( $p < 0.05$ ). Figure 5 shows that IL15 (5, 10 ng/ml) up-regulated gene expression of p53 (Figure 5a), bax ( Figure 5b), but didnt affected bcl2 gene expression (Figure 5c)



of IL15 (10 ng/ml) was significantly lower p53 expression (Figure 5a). The bax gene expression of MCF7 cells treated with 10 ng/ml IL15-induced NK cells was unsignificantly difference compared to 5 ng/ml IL15-induced NK cells.

### 3.6. Effect IL15-Induced NK Cells against MCF7 Cells Morphology

This study was performed by investigating the effect of IL15-induced NK cells on viability, density, morphology of MCF7 cells (Figure 6). The




Figure 6. Morphological analysis using phase-contrast microscope at  $\times 100$  magnification (A1, A2, A3, A4) and  $\times 40$  magnification (B1, B2, B3, B4). Different concentration of IL15 as inducer for NK cells decreased the survival rate of MCF7 cells

result demonstrates that NK cells induced dose-dependent reduction in viability of MCF7 cells.

Based on the morphological MCF7 cells treated with NK cells using transwell assay shows that NK cells lower the viability and density BC cells. The highest viability and density cells was MCF7 (control), the lowest viability, density cells was MCF7 treated by IL15 (5 ng/ml) as NK cells inducer.

#### 4. Discussion

NK cell abnormalities in cancer patients such as imbalance of immune status, inclined to immunosuppression, decreased in NK cell numbers, decreased cytotoxicity, declined activating receptor (NKG2D, NKG2C, Nkp30, Nkp46) and inclined inhibitory receptor (NKG2A, CD158, CD158a). Recently has been developed the immunotherapy using activated-NK cells (Mamessier *et al.* 2011). IL2, IL12, IL15, and IL18, applied systemically and for *ex vivo* activation and expansion of NK cells, have improved NK cell antitumor activity by increasing the expression of NK cell activating receptors and by inducing cytotoxic effector molecules (Konjevic *et al.* 2016). Cytokines such as interleukins (IL2) can be used to increase the anticancer potency of NK cells (Widowati *et al.* 2019). Various types of stimulation have been reported to enable NK cells to achieve their full effector potential, such as IL15, IL12, IL18, IL21 (Lucas *et al.* 2007; Guia *et al.* 2008; Chaix *et al.* 2008; Brehm *et al.* 2011). Based on the result (Figure 1), IL15 induced activating receptor NKG2D, this result was validated with previous research that IL2 and IL15 induce the expression of KIRs and activating receptors (NKG2D and Nkp44) on NK cell surface (Rham *et al.* 2007; Boieri *et al.* 2017). This result was in line with previous research that metastatic melanoma (MM) patients exhibited decreased CD161 and NKG2D (Konjević *et al.* 2009). NK cells treated with IL15 up-regulated the expression of NK receptors, including Nkp30, Nkp46, NKG2C, and NKG2D (Szczepanski *et al.* 2010). IL15 didn't decline the inhibitory receptor (NKG2A), this result was not consistent with previous research that IL15+IL12 was significantly decline the NKG2A of HIV-infected individuals (Parasa *et al.* 2012).

NK cells have been used in clinical studies in order to treat various malignancies (Harada *et al.* 2017), NK cells abnormalities in cancer patients such as low NK cells counts in PBMC, decreased cytotoxicity (Levy *et al.* 2011). NK cells inhibited MCF7 cells proliferation, IL15 activated NK cells to inhibit MCF7 cells proliferation, this result was consistent with previous research that NK cells control tumor growth and metastasis

diffusion *in vivo* (Zamai *et al.* 2007). The low NK cell numbers in tumors due to their inefficient homing into malignant tissues (Levy *et al.* 2011). Decreasing NK cell numbers are observed in peripheral blood (PB) of cancer patients; therefore, NK cells decrease in tumor infiltrate (Levy *et al.* 2011). The activity and numbers of NK cells need to be enhanced for better efficacy (Mandal and Viswanathan 2002). NK cell infiltration in solid tumors was associated with a better prognosis (Villegas *et al.* 2002). IL-15 is essential for NK cell survival, differentiation, and proliferation (Anton *et al.* 2015). Improving NK cells cytotoxicity used ILs (IL15, IL18), the result showed that ILs increased TNF $\alpha$ , IFN $\gamma$  secretion by NK cells (Widowati *et al.* 2020)

IL15 induces optimal production of IFN- $\gamma$  from NK cells, subsequently induce apoptosis of the NK cells toward cancer cells (Ross and Caligiuri 1997; Widowati *et al.* 2020). The human MHC class I-negative of small cell lung cancer cell line (N592) genetically engineered to secrete IL-15, N592/IL-15, showed a reduced tumor growth rate (Orengo *et al.* 2003). IL15 and IL12 induces optimal production of IFN- $\gamma$  from NK cells (Ross and Caligiuri 1997). IFN $\gamma$  inhibited BC proliferation (MCF7) with median inhibitory 0.34  $\mu$ g/ml (Widowati *et al.* 2016). IFN $\gamma$  against MCF7 cells exhibited that the cytokines decreased the cell viability in a dose dependent manner (Widowati *et al.* 2016).

NK secrete GzmB, IL15-activated NK cells up-regulated GzmB secretion. This result was validated previous research that activated NK cells released higher levels of IFN- $\gamma$ , TNF- $\alpha$ , Prf1, and GzmB compared to non-induced NK cells. IL2, IL15, and IL18 increased the secretion of IFN- $\gamma$ , TNF- $\alpha$ , Prf1, and GzmB by co-culture cells (Widowati *et al.* 2020). IL15 in NK cell controls as well survival of mature NK cells in the periphery (Marçais *et al.* 2013; Widowati *et al.* 2020), mediated by up-regulation of anti-apoptotic bcl2 family members and down regulation of apoptotic (Marçais *et al.* 2013). The production of TNF- $\alpha$ , IFN- $\gamma$ , Prf1, GzmB increased when the ratio of NK cells and hWJMSCs was high (Widowati *et al.* 2019). NK cells activity were controlled by cytokine and ILs (IL2, IL12, IL15, IL18) (Domaica *et al.* 2012). NK cells eliminate malignantly transformed cells principally by releasing the contents of cytotoxic granules into the immune synapse formed with their target cell (Lieberman 2003). The granule mediators of target cell lysis are serine proteases, known as Gzm, which induce programmed cell death (Safta *et al.* 2015; Harada *et al.* 2017). Cytotoxicity of NK cells is executed mainly through the granule exocytosis pathway by releasing Prf1 and GzmB into the immunological synapse after

the conjugate formation with targets (Lieberman 2003). Human GzmB preferentially induces target cell apoptosis, induces a rapid accumulation of the tumor-suppressor protein p53 within target cells (Safta *et al.* 2015). Human GzmB-induced p53 accumulates on target cell mitochondria where it interacts with the prosurvival protein bcl2. This interaction allows the release of the proapoptotic protein bax from its inhibitory interaction with bcl2 (Safta *et al.* 2015). Several proteins that are involved in GzmB-induced apoptosis, including casp-9 and -3, bim, bid, bak, bax, and xiap. GzmB induced apoptosis cancer cells by involving induction of p53 tumor suppressor gene (Meslin *et al.* 2007).

IL15 inhibit MCF7 proliferation, induced apoptotic BC cells, this result also supported by MCF7 morphology, BC cells appear low density and viability. This result was supported by previous research that ILs (IL1, IL2, IL15) and CD28, serve as co-stimulatory factors, enhancing IFN- $\gamma$  production by NK cells, as

well as NK cell proliferation and cytotoxicity (Hunter *et al.* 1997; Cui *et al.* 2016). ILs (IL12, IL18) stimulate NK cell lines, increasing the secretion of IFN- $\gamma$  (Wang *et al.* 2012). IL12 and IL18 are critical regulators that activate NK cells via the production of cytokines and direct lysis of target cells (Vivier *et al.* 2011). IL15 activate, increase NK proliferation (Widowati *et al.* 2020). Activated NK cells induce IFN- $\gamma$ , TNF- $\alpha$ , Prf1, GzmB secretion and inhibit BC cells proliferation on co-cultured of MCF7 and NK cells (Widowati *et al.* 2020).

The proposed mechanism of our research which IL15 activated NK cells including increased NK cells number, up regulated activating receptor (NKG2D) but IL15 didn't influence inhibitor receptor (NKG2A), improved Prf, Gzm secretion. IL15-activated NK induced apoptosis of BC cells through increased p53, bax gene expression and inhibited BC cells proliferation. For the detail mechanism can be seen at Figure 7.




Figure 7. Proposed mechanism of activated NK cells to inhibit and kill breast cancer cells. NK cells abnormalities in breast cancer patients. The imbalance of immune status, which deficient immunosurveillance including low NK cell number in PBMC, poor tumor infiltrate to cancer cells, decreased cytotoxicity, defective expression of activating receptor (NKG2D). IL15 activated NK cells to kill target cancer cells through: cytoplasmic granules release (Prf, GzmB), death receptor-induced apoptosis (TNF $\alpha$ ), effector molecules production (IFN $\gamma$ ) and released antibody-dependent cellular cytotoxicity (ADCC). p53, bax induced apoptotic BC cells



## 5. Conclusion

IL15 improve, activate NK cells resulted in activating receptors (NKG2D), increasing GzmB secretion and cytotoxic activity on BC cells, inducing apoptotic genes bax, p53 expression and induce apoptotic cells on transwell assay.

## Acknowledgements

This study was supported by the Grants-in-Aid Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) 2019-2020, from Ministry of Research, Technology and Higher Education of the Republic of Indonesia. The authors like to thank to Hanna Sari Widya Kusuma, Aditya Rinaldy, Cintani Dewi Wahyuni, Kamila Yashfa Gunawan, Muhammad Aldi Maulana, Tri Wahyuni from Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung, West Java, Indonesia for their technical assistants.

## References

- Afifah E *et al.* 2019. Induction of matrix metalloproteinases in chondrocytes by interleukin IL-1 $\beta$  as an osteoarthritis model. *J Math Fundam Sci* 51:103-111.
- Albertsson PA *et al.* 2003. NK cells and the tumour microenvironment: implications for NK-cell function and anti-tumour activity. *Trends Immunol* 24:603-609.
- Anton OM *et al.* 2015. NK cell proliferation induced by IL-15 transpresentation is negatively regulated by inhibitory receptors. *J Immunol* 195:4810-4821.
- Behbahani M. 2014. Evaluation of *in vitro* anticancer activity of *Ocimum basilicum*, *Alhagi maurorum*, *Calendula officinalis* and their parasite *Cuscuta campestris*. *PLoS One* 9:e116049.
- Brehm C *et al.* 2011. IL-2 stimulated but not unstimulated NK cells induce selective disappearance of peripheral blood cells: Concomitant results to a phase I/II study. *PLoS One* 6:e2735.
- Boieri M *et al.* 2017. IL-12, IL-15, and IL-18 pre-activated NK cells target resistant T cell acute lymphoblastic leukemia and delay leukemia development *in vivo*. *Oncoimmunol* 6:e1274478.
- Chaix J *et al.* 2008. Priming of natural killer cells by Interleukin-18. *J Immunol* 181:1627-1631.
- Cheng M *et al.* 2013. NK cell-based immunotherapy for malignant disease. *Cellular Mol Immunol* 10:230-252.
- Cui R *et al.* 2016. Human mesenchymal stromal/stem cells acquire immunostimulatory capacity upon cross-talk with natural killer cells and might improve the NK cell function of immunocompromised patients. *Stem Cell Res Ther* 7:1-13.
- Dewan MZ *et al.* 2009. Natural killer activity of peripheral blood mononuclear cells in breast cancer patients. *Biomed Pharmacother* 63:703-706.
- Domaica CI *et al.* 2012. Human natural killer cell maturation defect supports *in vivo* CD56 bright to CD56 dim lineage development. *PLoS One* 7:e51677.
- Esendagli G *et al.* 2008. Malignant and non-malignant lung tissue areas are differentially populated by natural killer cells and regulatory T cells in non-small cell lung cancer. *Lung Cancer* 59:32-40.
- Ferlay J *et al.* 2015. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 136:359-386.
- Guan F *et al.* 2018. Induction of apoptosis by Bigelovii A through inhibition of NF- $\kappa$ B activity. *Mol Med Rep* 18:1600-1608.
- Guia S *et al.* 2008. A role for interleukin-12/23 in the maturation of human natural killer and CD56+ T cells *in vivo*. *Blood* 111:5008-5016.
- Han H *et al.* 2018. Inhibition of cell proliferation and migration through nucleobase-modified polyamidoamine-mediated p53 delivery. *Int J Nanomed* 13:1297-1311.
- Harada Y *et al.* 2017. Clinical applications of natural killer cells. Chapter V. DOI:10.5772/intechopen.68991
- Hwang YJ *et al.* 2012. A study on the immunomodulation effect of *Isodon japonicus* extract via splenocyte function and NK anti-tumor activity. *Int J Mol Sci* 13:4880-4888.
- Hunter CA *et al.* 1997. Type I interferons enhance production of IFN-gamma by NK cells. *Immunol Lett* 59:1-5.
- Kelly JM *et al.* 2003. Induction of tumor-specific T cell memory by NK cell-mediated tumor rejection. *Nat Immunol* 3:83-90.
- Konjevic G *et al.* 2009. Biomarkers of suppressed natural killer (NK) cell function in metastatic melanoma: decreased NKG2D and increased CD158a receptors on CD3-CD16+ NK cells. *Biomarkers* 4:258-270.
- Konjevic G *et al.* 2016. Natural killer cell receptors: alterations and therapeutic targeting in malignancies. *Immunol Res* 64:25-35.
- Levy EM *et al.* 2011. Natural killer cells in human cancer: from biological functions to clinical applications. *J Biomed Biotechnol* 2011:1-11.
- Lieberman J. 2003. The ABCs of granule-mediated cytotoxicity: new weapons in the arsenal. *Nat Rev Immunol* 3:361-370.
- Lucas M *et al.* 2007. Natural killer cell-mediated control of infections requires production of interleukin 15 by Type I IFN-triggered dendritic cells. *Immunity* 26:503-517.
- Mandal A, Viswanathan C. 2002. Natural killer cells: in health and disease. *Hematol Oncol Stem Cell Ther* 8:23-28.
- Mamessier E *et al.* 2011. Human breast cancer cells enhance self tolerance by promoting evasion from NK Cell antitumor immunity. *J Clin Invest* 121:3609-3622.
- Marcais A *et al.* 2013. Regulation of mouse NK cell development and function by cytokines. *Front Immunol* 4:1-14.
- Meslin F *et al.* 2007. Granzyme B-induced cell death involves induction of p53 tumor suppressor gene and its activation in tumor target cells. *J Biol Chem* 282:32991-32999.
- Orengo AM *et al.* 2003. Tumor cells engineered with il-12 and il-15 genes induce protective antibody responses in nude mice. *J Immunol* 171: 569-575.
- Parasa VRR *et al.* 2012. Effect of recombinant cytokines on the expression of natural killer cell receptors from patients with tb or/and hiv infection. *PLoS One* 7:e37448.
- Zamai L *et al.* 2007. NK cells and cancer. *J Immunol* 178:4011-4016.
- Rham de C *et al.* 2007. The proinflammatory cytokines IL-2, IL-15 and IL-21 modulate the repertoire of mature human natural killer cell receptors. *Arthritis Res Ther* 9: R125.
- Ross ME, Caligiuri MA. 1997. Cytokine-induced apoptosis of human natural killer cells identifies a novel mechanism to regulate the innate immune response. *Blood* 9: 910-918.
- Sadeghi F *et al.* 2015. The effect of estrogen on the expression of cartilage-specific genes in the chondrogenesis process of adipose-derived stem cells. *Adv Biomed Res* 4:1-16.

- Safta TB *et al.* 2015. Granzyme B-activated p53 interacts with Bcl-2 to promote cytotoxic lymphocyte-mediated apoptosis. *J Immunol* 194:418-428.
- Szczepanski MJ *et al.* 2010. Interleukin-15 enhances natural killer cell cytotoxicity in patients with acute myeloid leukemia by upregulating the activating NK cell receptors. *Cancer Immunol Immunother* 59:73-79.
- Smyth MJ *et al.* 2005. Activation of NK cell cytotoxicity. *Mol Immunol* 42:501-510.
- Tirona MT *et al.* 2010. Prevention of breast cancer (part 1): epidemiology, risk factors, and risk assessment tools. *Cancer Invest* 28:743-750.
- Villegas FR *et al.* 2002. Prognostic significance of tumor infiltrating natural killer cells subset CD57 in patients with squamous cell lung cancer. *Lung Cancer* 35:23-28.
- Vivier E *et al.* 2011. Innate or adaptive immunity? The example of natural killer cells. *Sci* 31:44-49.
- Wang J *et al.* 2009. Curcumin induces apoptosis through the mitochondria-mediated apoptotic pathway in HT-29 cells. *J Zhejiang Univ Sci B* 10:93-102.
- Wang R *et al.* 2012. Natural killer cell-produced IFN- $\gamma$  and TNF- $\alpha$  induce target cell cytotoxicity through up-regulation of ICAM-1. *J Leukoc Biol* 91:299-309.
- Widowati W *et al.* 2016. Selective cytotoxic potential of IFN- $\gamma$  and TNF- $\alpha$  on breast cancer cell lines (T47d and MCF-7). *Asian J Cell Biol* 11:1-12.
- Widowati W *et al.* 2018. Direct and indirect effect of TNF $\alpha$  and IFN $\gamma$  toward apoptosis in breast cancer cells. *Mol Cell Biomed Sci* 2:60-69.
- Widowati W *et al.* 2019. Effects of conditioned medium of co-culture IL-2 induced nk cells and human wharton's jelly mesenchymal stem cells (hwjmscs) on apoptotic gene expression in a breast cancer cell line (MCF-7). *J Math Fund Sci* 51:205-224.
- Widowati W *et al.* 2020. Effect of interleukins (IL2, IL15, IL18) on receptors activation and cytotoxic activity of natural killer cells in breast cancer cells. *African Health Sci* 20:1-12.