



Research article

Hepatotoxicity prevention in Acetaminophen-induced HepG2 cells by red betel (*Piper crocatum* Ruiz and Pav) extract from Indonesia via antioxidant, anti-inflammatory, and anti-necroticChrismis Novalinda Ginting^{a,*}, I Nyoman Ehrich Lister^a, Ermi Girsang^a, Wahyu Widowati^b, Dewani Tediana Yusepany^c, Alya Mardhotillah Azizah^c, Hanna Sari Widya Kusuma^c^a Faculty of Medicine, Universitas Prima Indonesia, Jl. Belanga No. 1, Medan 20118, North Sumatera, Indonesia^b Faculty of Medicine, Maranatha Christian University, Jl. Prof. Drg. Surya Sumantri No. 65, Bandung 40163, West Java, Indonesia^c Aretha Medika Utama, Biomolecular Research Center, Jl. Babakan Jeruk II No.9, Bandung 40163, West Java, Indonesia

ARTICLE INFO

Keywords:

Red betel leaves extract
Acetaminophen
HepG2 cells
Hepatoprotective
Biochemistry
Immunology
Inflammation
Natural product
Pharmaceutical science

ABSTRACT

Acetaminophen (APAP) is a widely used analgesic, but it may cause liver injury (hepatotoxicity) via oxidative stress that induced by N-acetyl-p-benzoquinone imine (NAPQI) in long term usage or overdose. Multiple inflammatory mediators were also found to contribute for this effect. Many medicinal plants was known for its antioxidant and anti-inflammatory activities and one of them is Red betel (*Piper crocatum* Ruiz and Pav) from Indonesia. In this study, the red betel leaves extract (RBLE) protective effect against APAP-induced HepG2 cells was determined. APAP-induced HepG2 as hepatotoxicity cell model was treated with RBLE at 25 and 100 µg/mL. Protective effects of RBLE toward hepatotoxicity were evaluated by several parameters: tumor necrosis factor- α (TNF- α) concentration, reactive oxygen species (ROS) level, live cells percentage, apoptotic cells percentage, necrotic cells percentage, death cells percentage, CYP2E1 and GPX gene expression. The RBLE treatments (both 25 and 100 µg/mL) increased CYP2E1 and GPX gene expression also live cells percentage, while decreased ROS level, TNF- α concentration, also the percentage of death and necrotic cells. Red Betel leaves ethanol extract has hepatoprotective effect via anti-inflammatory, anti-necrotic, and antioxidant potency in liver injury model.

1. Introduction

For many problem of drugs use, liver injury was continues to be a problem. It was represents a major challenge in designing potential therapies (Noh et al., 2015). Acetaminophen (paracetamol, APAP) is considered as first line analgesics. However, excessive use of APAP leads to liver injury even liver failure in human (Ganey et al., 2007; Ni et al., 2012). In small percentage, the cytochrome P450 2E1 (CYP2E1) enzymes was oxidizing the APAP and form N-acetyl-p-benzoquinone imine (NAPQI), a highly reactive intermediate, which is detoxified by covalent binding with glutathione (GSH). However, in APAP poisoning, it will generates excess NAPQI which evokes the GSH depletion that binds to macromolecules triggering oxidative stress, mitochondrial dysfunction, and ultimately resulting in hepatocellular death (Salminen et al., 2012; Uzi et al., 2013; Yuan et al., 2016). Although the mechanisms underlying hepatotoxicity that induced by APAP still unclear, some evidences was indicate that inflammation mediators such as tumor necrosis factor- α

(TNF- α) also oxidative stress was contribute to the APAP-induced acute liver damage pathology process (Uzkeser et al., 2012; Dragomir et al., 2012).

One of betel in Indonesia namely red betel (*Piper crocatum* Ruiz and Pav) has medicinal function and used as medicine since its introduce as medicinal plants producer in Blunyahrejo (Rinanda and Alga, 2012). It can be used to treat diabetes, gout, hepatitis, hypertension, and eye inflammation (Anugrahwati et al., 2016). In previous study, red betel leaves were found to have some secondary metabolite content like flavonoids, alkaloids, tannins, saponins, triterpenoids steroids, quinones, polyphenolics, and essential oil groups (Arambewela et al., 2005; Wulandari et al., 2018). In addition, red betel contains phenolic compounds in the form of hydrochavicol, cavitbetol acetate and eugenol (Swapna et al., 2012; Dervis et al., 2017). In previous studies, red betel leaves extract (RBLE) was shown have anti-inflammatory properties (Misra et al., 2009); antioxidant activity (Lister et al., 2019a); and also

* Corresponding author.

E-mail address: chrismis@unprimdn.ac.id (C.N. Ginting).

Table 1. RT-PCR details of β -Actin, CYP2E1, and GPX gene.

Gene Symbols	Primer Sequences (5' to 3') Upper strand: Sense Lower strand: Antisense	Annealing (°C)	Cycle	References
β -Actin	5'-TCTGGCACCACCTTCTACAATG-3' 5'-AGCACAGCCTGGATAGCAACG-3'	63	40	Widowati et al. (2019b) Afifah et al. (2019)
CYP2E1	5'-GTTCTTTGCGGGGACAGAGA-3' 5'-GAGGGTGATGAACCGCTGAA-3'	59	40	Kim et al., 2003
GPX	5'-CCAAGCTCATCACCTGGTCT-3' 5'-TCGATGCAATGGTCTGAA-3'	59	40	Ugusman et al. (2011)

have anticancer activity especially cervical cancer (Widowati et al., 2013) and breast cancer (Zulharini et al., 2018).

In this study, RBLE potential to suppress liver injury in APAP-induced HepG2 cells was conducted. The parameters that observed in this study was Reactive Oxygen Species (ROS) level; TNF- α level; Cytochrome P450 Family 2 Subfamily E Member 1 (CYP2E1) and Glutathione Peroxidase (GPX) gene expression; apoptotic, necrotic cells, and death cells percentage.

2. Materials and methods

2.1. Preparation of red betel leaves extract

The red betel (*P. crocatum* Ruiz and Pav) leaves that used in this study was obtained from Pabuaran Cilendek Timur, Indonesia and has been identified by Herbarium Bogoriense, Botanical Field Research Center for Biology-Indonesian Institute of Science, Indonesia. RBLE preparation was done by using maceration method. RBLE was obtained from our previous research and stored at -20 °C (Lister et al., 2019a, 2019b).

2.2. HepG2 cells culture and APAP-Induced HepG2

The cells that used in this study is human hepatocellular carcinoma (HepG2) cell line (ATCC, HB-8065TM) from Aretha Medika Utama Biomolecular and Biomedical Research Center, Bandung, Indonesia. It was grown in complete medium with composition: Modified Eagle Medium (MEM) (Biowest, L0416-500), fetal bovine serum (FBS) (Biowest, S1810) as much as 10% (v/v), antibiotic-antimycotic (Gibco, 15240062) as much as 1% (v/v), also nanomycopolitine (Biowest, LX16) addition as much as 1% of (v/v). Acetaminophen (Sigma Aldrich, A7085) with concentration at 40 mM was used to induce the hepatotoxicity. When the cells were confluent, it was rinsed using PBS and detached using trypsin-EDTA (Gibco, 25200072) with incubation at 37 °C. In 6 well plates, the cells was seeded (5×10^5 cells/well) and then incubated at the same temperature with 5% of CO₂ for 24 h. The cells was induced by RBLE and incubated again for 24 h. According to the treatment, it was divided into 5 groups: I) Normal Cells; II) DMSO1%; III) APAP 40 mM; IV) APAP 40 mM + RBLE 25 μ g/mL; V) APAP 40 Mm + RBLE 100 μ g/mL. Then it was centrifuged at 1600 rpm for 10 min. The supernatant was collected as sample for the Elisa assay (Luo et al., 2016; Aouache et al., 2018; Lister et al., 2019b).

2.3. Total protein assay

Bovine Serum Albumin (BSA) (Sigma Aldrich, A9576) was used as standard in this method. Briefly standard solutions as much as 20 μ L also same volume for the samples was mixed with Quick Start Dye Reagen 1X (Biorad, 5000205) as much as 200 μ L into each well in 96 well plate. The mixture then incubated at room temperature for around 5 min. The wavelength at 595 nm was used to determine the mixture absorbance by using microplate reader (MultiskanTM GO Micro plate Spectrophotometer, Thermo Scientific, 51119300) at 595 nm. The result from this assay was used for normalization of TNF- α data calculation (Pluemsamran et al., 2012; Widowati et al., 2019a).

2.4. TNF- α assay

This assay was measured using ELISA assay (BioLegend, 421701) and done according to the manufacturer's kit manual. Based on the manual, wavelength at 450 nm was used to determine the absorbance using microplate reader (Widowati et al., 2019a).

2.5. Apoptotic activity assay

The assay was conducted using methods that reported by Widowati et al. (2019b). Treated and control HepG2 cells were washed using PBS 1x and harvested using trypsin-EDTA for apoptotic assay. The pellet was washed using Annexin Binding Buffer 1X (Miltenyi Biotec, 130-092-820) 500 μ L and stained with Annexin V-FITC (BioLegend, 79998) and Propidium Iodide (BioLegend, 79997). Cells were incubated at 37 °C for 30 min in the dark. Cells were later suspended in Annexin Binding Buffer 1x. The HepG2 cells apoptotic percentage were analyzed using MACSquant Analyzer 10 (Miltenyi Biotec).

2.6. Reactive oxygen species (ROS) assay

HepG2 cells were digested with trypsin-EDTA after cultured around 7 days and 2.5×10^4 cells/0.5 mL were incubated for 45 min in 20 μ M DCF-DA at 37 °C and incubated again for 4 h in RBLE. Based on Prastuti et al. (2019) and Girsang et al. (2019), the 2', 7'-dichlorofluorescein diacetate (DCFDA)-Cellular Reactive Oxygen Species Detection Assay Kit (Abcam, ab113851) was used to measured intracellular ROS with modifications.

Table 2. RNA concentration and purity.

No.	Sample	Concentration (ng/ μ L)	Purity (Absorbance 260/280)
1.	Control cells	92.90	2.3212
2.	Positive control	90.10	2.0904
3.	Positive control + RBLE 25 μ g/mL	36.20	1.9676
4.	Positive control + RBLE 100 μ g/mL	40.00	2.0366




Figure 1. RBLE treatment effect toward TNF-α concentration in APAP-induced HepG2 cells as hepatotoxicity model. (A) TNF-α concentration (pg/mL) on hepatotoxicity model. (B) TNF-α concentration (pg/mg protein) on hepatotoxicity model. *Data was included as mean ± standard deviation. I) Normal cells as negative control; II) Normal cells + DMSO 1% as vehicle control; III) APAP-induced cells (Positive control); IV) Positive control + RBLE 25 µg/mL; V) Positive control + RBLE 100 µg/mL. Significance among treatments toward TNF-α concentration was presented as different letters (a,b) based on Tukey HSD post hoc test (P < 0.05).

2.7. The expression of GPX and CYP2E1 gene assay

Cells that has been harvested was processed for RNA isolation that will be used for futher assay. It was done by using the Aurum™ Total RNA mini Kit (Bio-Rad, 732-6820). RT-qPCR (Clever, GTC96S) was used to analyze the gene expression include the β-actin gene that constitutively expressed (Afifah et al., 2019; Widowati et al., 2019b). Table 1 was shown the primer sequence and Table 2 was shown RNA concentration and purity.

2.8. Statistical analysis

All data were obtained after doing it in triplicate. When the data has normal distribution, it was analyzed using ANOVA and Post Hoc Test using Tukey HSD with p < 0.05 while data didn't has normally distributed were analyzed with Kruskal Wallis and Post Hoc Test Mann Whitney using SPSS software (version 20.0). The data were presented as mean ± standard deviation.




Figure 2. RBLE effect toward apoptotic, necrotic, dead cells in hepatotoxicity model. (A) Live cells on hepatotoxicity model. (B) Early apoptotic on hepatotoxicity model. (C) Late apoptotic on hepatotoxicity model. (D) Necrotic on hepatotoxicity model. *Data was included as mean ± standard deviation. I) Normal cells as negative control; II) Normal cells + DMSO 1% as vehicle control; III) APAP-induced cells (Positive control); IV) Positive control + RBLE 25 µg/mL; V) Positive control + RBLE 100 µg/mL. There are significant different between all groups based on Kruskal-Wallis Test (P < 0.05) and Mann-Whitney Test (P < 0.05). It was marked as single star (*) marks for the statistical difference between positive control and negative control while the hashtag (#) mark for statistical difference between treatment and positive control.




Figure 3. RBLE effect toward ROS level in hepatotoxicity model. *Data was included as mean \pm standard deviation. I) Normal cells (Negative control); II) Normal cells + DMSO 1%; III) APAP-induced cells (Positive control); IV) Positive control + RBLE 25 μ g/mL; V) Positive control + RBLE 100 μ g/mL. Based on Kruskal-Wallis Test ($P < 0.05$), there are significant different among groups. It was marked as single star (*) for statistical difference between positive control and negative control also hashtag (#) for statistical difference between treatment and positive control.

3. Result

3.1. RBLE effect towards TNF- α concentration in APAP-induced HepG2 cells

APAP was increased the TNF- α concentration in HepG2 cells. When RBLE treatment was added, it was found can decrease the TNF- α concentration (Figure 1). Based on the result, RBLE has potential to suppress the TNF- α production in HepG2 cells that induced by APAP.

3.2. Effect of RBLE towards apoptotic, necrotic, and cell death in APAP-induced HepG2 cells

APAP decreased live cell percentage compare to normal HepG2 cells (Figure 2A). RBLE treatment decreased the percentage of apoptotic and necrotic significantly in APAP-induced HepG2 cells (Figure 2B–D). RBLE treatment can increase the live cells percentage also reduce the percentage of necrotic and dead cells in HepG2 cells that induced by APAP.




Figure 4. RBLE effect toward the expression of CYP2E1 gene in hepatotoxicity model. *Data was included as mean \pm standard deviation. I) Normal cells as negative control; II) Normal cells + DMSO 1%; III) APAP-induced cells (Positive control); IV) Positive control + RBLE 25 μ g/mL; V) Positive control + RBLE 100 μ g/mL. Based on ANOVA ($P < 0.05$) and Games-Howell ($P < 0.05$), there are significant different among all groups. It was marked as single star (*) marks for the difference between positive control and negative control also hashtag (#) mark for the difference between treatment and positive control.




Figure 5. RBLE effect toward the expression of GPX gene in hepatotoxicity model. *Data was included as mean \pm standard deviation. I) Normal cells (Negative control); II) Normal cells + DMSO 1%; III) APAP-induced cells (Positive control); IV) Positive control + RBLE 25 μ g/mL; V) Positive control + RBLE 100 μ g/mL. Based on Kruskal-Wallis Test ($P < 0.05$) and Mann-Whitney Test ($P < 0.05$), there are significant different between group. It was marked as single star (*) marks as difference between positive control and negative control while hashtag (#) mark as difference between treatment and positive control.

3.3. RBLE effect towards ROS level in liver injury model

ROS level increased significantly after APAP induction and reduced significantly when injured HepG2 cells were treated with RBLE (Figure 3). RBLE had potential to decrease ROS level in liver injury model.

3.4. RBLE effect on CYP2E1 and GPX gene expression in liver injury model

CYP2E1 gene expression decreased significantly in APAP-induced HepG2 cells. RBLE treatments increased the CYP2E1 gene expression significantly compare to the APAP-induced HepG2 cells group (Figure 4). GPX gene expression decreased in APAP-induced HepG2 cells. RBLE treatments could increase the GPX gene expression significantly (Figure 5). RBLE treatments had ability to increase the CYP2E1 and GPX gene expression.

4. Discussions

Betel leaves had been known to contain many active compounds, mainly hydrochavicol, cavibetol acetate and eugenol (Begam et al., 2018). Based on previous study, it had been demonstrated that red betel leaves extract, along with its active constituents: eugenol and hydroxychavicol, can scavenging H_2O_2 and DPPH also reducing FRAP and ABTS radicals that indicated their antioxidant activity (Lister et al., 2019a). Eugenol also had been reported could decrease the ALT and AST activities and LDH level in liver injury model that induced by APAP (Lister et al., 2019b).

The presence of APAP toxic metabolite NAPQI caused Kupffer cells activation that leads to TNF- α release (Legert et al., 2015). TNF- α , one of inflammatory cytokine, involved in oxidative stress injury (Barman et al., 2016; Jaeschke et al., 2012). It was mediated death receptor pathway apoptosis by activating caspase 3 that act as a central effector to cleave various cellular substrates and trigger cell apoptosis eventually (Nagase et al., 2002; Truong et al., 2016). While apoptosis and necrosis frequently coexist in liver pathological conditions and the cell death balance may be dictated by the particular insult (Antoine et al., 2010).

RBLE treatment was found can decrease the TNF- α level in liver injury model based on the study result. One of active compound in RBLE, eugenol, had been studied have effect on reduction of inflammatory cells infiltration and generation of cytokines from Kupffer cells include ability to suppress TNF- α level in liver injury model (Yogalakshmi et al., 2010).




Figure 6. Proposed RLBE hepatoprotective mechanism in liver injury model. *CYP2E1 act to transform APAP to NAPQI. It induce GSH depletion then induce production of ROS. The excessive of ROS decrease the GPX gene expression leads to increase cell death. NAPQI activate the kupper cell and leads to TNF- α production that induce the JNK signaling pathways that also increase ROS level and leads to upregulate cell necrosis; increase inflammation; and induce cell death. While JNK induced the Bcl-2 down-regulation and Bax up-regulation, resulting in activation of caspase 9 and caspase 3 that leads to apoptosis cells. The RLBE treatments could inhibit the excessive ROS and TNF- α . It also could lowering the necrosis and apoptosis that leads to lowering inflammation. RBLE treatments decrease the cell death and increase the survival hepatic cells.

Phenolic compound had anti-inflammatory effect as another study from Yuan et al. (2016) also stated that a phenolic compound ferulic acid could decrease the TNF- α level in mice induced with APAP.

Figure 2 shows that the APAP induction increased the apoptotic, necrotic, and death cells percentage, while RBLE treatments had successfully reduce death cells and maintain live cells at higher level. This data was in line with previous research that less apoptotic cells were seen in ferulic acid treatment in injury liver model (Yuan et al., 2016).

In APAP-induced hepatotoxicity model, oxidative stress played an important role and it was characterized by ROS accumulation (Nagi et al., 2010; Du et al., 2016). NAPQI, a reactive metabolite formed from APAP, could react rapidly with GSH and aggravating oxidative stress in conjunction with mitochondrial dysfunction that induced hepatocellular damage (Smith et al., 2016; Kang et al., 2017). The enzymatic antioxidant defense system known can detoxified ROS. Previous study exhibited that RBLE had antioxidant potential (Lister et al., 2019a). Based on the result, RBLE proved to suppress the ROS level in liver injury model, this result was in line with Parikh et al. (2015) that found phenolic compounds in *Brassica juncea* hydromethanolic extract such as quercetin and catechin could reduce the ROS level in APAP-induced HepG2 cells. Thus, the RBLE hepatoprotective mechanism might result from diminishing generation of ROS.

In the metabolism of endogenous and exogenous compounds wide variety, CYP2E1 has important functions that relevant to chemical toxicity and carcinogenesis in liver (Gonzalez, 2007). ROS was one of compounds that generated by CYP2E1 that increase mitochondrial membrane permeability and lipid peroxidation, which induce apoptosis via pro-apoptotic factors release and activate caspase 3 (Lee and Wei, 2007). Based on the result, RBLE treatments decreased CYP2E1 gene expression in APAP-induced HepG2 cells, probably by its high phenolic compounds. This result was in line with previous research that in APAP-induced hepatotoxicity, the ferulic acid could inhibit the up-regulation of CYP2E1 expression (Yuan et al., 2016).

The enzymatic antioxidant defense system primary part against oxidative stress is GPX that directly eliminating ROS (Truong et al., 2016). When free radicals formed rapidly, GPX functions will become inefficient and leads to hepatocytes damage (Roh et al., 2018). GPX level can be used as indicator of the oxidative stress response (Wang et al., 2016). Based on the result, it was shown that APAP could decrease the GPX expression, however RBLE treatments could counter this effect. It was indicated that RBLE can protects cells/livers from APAP-inducer through an antioxidant defense system enhancement. Truong et al. (2016) also stated that a phenolic compound, mainly quercitrin could restore GPX expression and attenuates APAP-induced liver damage.

Based on this study, RBLE was shown have antioxidant, anti-necrotic, and anti-inflammatory activities. It mechanism as hepatoprotective agent in live injury was shown in Figure 6 that proposed by us based on the study result and literature review.

5. Conclusion

Red betel leaves extract treatments could reduce TNF- α level, reduce cell's apoptosis and increase live cells percentage, reduce intracellular ROS, reduce CYP2E1 and increase GPX level in HepG2 cells. This marked the hepatoprotective potential of RBLE through antioxidant, anti-necrotic, and anti-inflammatory activities. Further research on *in vivo* model is needed to confirm current result.

Declarations

Author contribution statement

C.N. Ginting, I.N.E. Lister and E. Girsang: Conceived and designed the experiments.

W. Widowati: Conceived and designed the experiments; Analyzed and interpreted the data.

D.T. Yusepany: Performed the experiments; Contributed reagents, materials, analysis tools or data.

A.M. Azizah: Analyzed and interpreted the data; Wrote the paper.

H.S.W. Kusuma: Contributed reagents, materials, analysis tools or data; Wrote the paper.

Funding statement

This work was funded by Universitas Prima Indonesia, Medan, Indonesia.

Declaration of interests statement

The authors declare no conflict of interest.

Additional information

No additional information is available for this paper.

Acknowledgements

We extend our gratitude to Jenifer Kiem Aviani, Rr. Anisa Siwanti Handayani, Kamila Yashfa Gunawan, Dwi Surya Artie, and Seila Arumwardana from Aretha Medika Utama, Biomolecular and Biomedical Research Center for their technical support.

References

- Afifah, E., Mozef, T., Sandra, F., Arumwardana, S., Rihibiha, D.D., Nufus, H., Rizal, R., Amalia, A., Bachtiar, I., Murti, H., Widowati, W., 2019. Induction of matrix metalloproteinases in chondrocytes by Interleukin IL-1 β as an osteoarthritis model. *J. Math. Fund. Sci.* 51 (2), 103–111.
- Antoine, D.J., Williams, D.P., Kipar, A., Laverty, H., Park, B.K., 2010. Diet restriction inhibits apoptosis and HMGB1 oxidation and promotes inflammatory cell recruitment during acetaminophen hepatotoxicity. *Mol. Med.* 16 (11–12), 479–490.
- Anugrahwati, M., Purwaningsih, T., Manggalarini, J.A., Alnavis, N.B., Wulandari, D.N., Pranowo, H.D., 2016. Extraction of ethanolic extract of red betel leaves and its cytotoxicity test on HeLa cells. *Proc. Eng.* 148, 1402–1407.
- Aouache, R., Biquard, L., Vaiman, D., Miralles, F., 2018. Oxidative stress in preeclampsia and placental diseases. *Int. J. Mol. Sci.* 19 (5), 1496.
- Arambewela, L.S.R., Arawwawala, L.D.A.M., Ratnasooriya, W.D., 2005. Antidiabetic activities of aqueous and ethanolic extracts of *Piper betle* leaves in rats. *J. Ethnopharmacol.* 102 (2), 239–245.
- Barman, P.K., Mukherjee, R., Prusty, B.K., Suklabaidya, S., Senapati, S., Ravindran, B., 2016. Chitohexaose protects against acetaminophen-induced hepatotoxicity in mice. *Cell Death Dis.* 7 (5), e2224.
- Begam, K.M.F., Ravichandran, P., Manimekalai, V., 2018. Phytochemical analysis of some selected varieties of *Piper betle* L. *Int. J. Curr. Pharmaceut. Res.* 10 (2), 89–93.
- Dervis, E., Yurt Kilcar, A., Medine, E.L., Tekin, V., Cetkin, B., Uygur, E., Muftuler, F.Z.B., 2017. In vitro incorporation of radioiodinated eugenol on adenocarcinoma cell lines (Caco2, MCF7, and PC3). *Cancer Biother. Radiopharmaceutic.* 32 (3), 75–81.
- Dragomir, A.C., Sun, R., Mishin, V., Hall, L.B., Laskin, J.D., Laskin, D.L., 2012. Role of galectin-3 in acetaminophen-induced hepatotoxicity and inflammatory mediator production. *Toxicol. Sci.* 127 (2), 609–619.
- Du, K., Ramachandran, A., Jaeschke, H., 2016. Oxidative stress during acetaminophen hepatotoxicity: sources, pathophysiological role and therapeutic potential. *Redox Biol.* 10, 148–156.
- Ganey, P.E., Luyendyk, J.P., Newport, S.W., Eagle, T.M., Maddox, J.F., Mackman, N., Roth, R.A., 2007. Role of the coagulation system in acetaminophen-induced hepatotoxicity in mice. *Hepatology* 46 (4), 1177–1186.
- Girsang, E., Lister, I.N.E., Ginting, C.N., Nasution, S.L., Suhartina, S., Munshy, U.Z., Rizal, R., Widowati, W., 2019. Antioxidant and anti-inflammatory activity of chlorogenic acid on lead-induced fibroblast cells. *J. Phys. Conf.* 1374, 012006.
- Gonzalez, F.J., 2007. The 2006 Bernard B. Brodie Award Lecture. Cyp2e1. *Drug Metabol. Dispos.* 35 (1), 1–8.
- Jaeschke, H., Williams, C.D., Ramachandran, A., Bajt, M.L., 2012. Acetaminophen hepatotoxicity and repair: the role of sterile inflammation and innate immunity. *Liver Int.* 32 (1), 8–20.
- Kang, E.S., Lee, J., Homma, T., Kurahashi, T., Kobayashi, S., Nabeshima, A., Yamada, S., Seo, H.G., Miyata, S., Sato, H., Fujii, J., 2017. xCT deficiency aggravates acetaminophen-induced hepatotoxicity under inhibition of the transsulfuration pathway. *Free Radic. Res.* 51 (1), 80–90.
- Kim, S.S., Oh, O.J., Min, H.Y., Park, E.J., Kim, Y., Park, H.J., Han, Y.N., Lee, S.K., 2003. Eugenol suppresses cyclooxygenase-2 expression in lipopolysaccharide-stimulated mouse macrophage RAW264.7 cells. *Life Sci.* 73 (3), 337–348.
- Lee, H.C., Wei, Y.H., 2007. Oxidative stress, mitochondrial DNA mutation, and apoptosis in aging. *Exp. Biol. Med.* 232 (5), 592–606.
- Legert, K.G., Tsilingaridis, G., Remberger, M., Ringdén, O., Heimdahl, A., Yucel-Lindberg, T., Dahllöf, G., 2015. The relationship between oral mucositis and levels of pro-inflammatory cytokines in serum and in gingival crevicular fluid in allogeneic stem cell recipients. *Supp. Care Cancer* 23 (6), 1749–1757.
- Lister, I.N.E., Ginting, C.N., Girsang, E., Armansyah, A., Marpaung, H.H., Sinaga, A.P.F., Handayani, R.A.S., Rizal, R., 2019a. Antioxidant properties of red betel (*Piper crocatum*) leaf extract and its compounds. *J. Nat. Remedies* 19 (4), 198–205.
- Lister, I.N.E., Ginting, C.N., Girsang, E., Amansyah, A., Chiuman, L., Yanti, N.L.W.E., Rizal, R., Widowati, W., 2019b. Hepatoprotective effect of eugenol on acetaminophen-induced hepatotoxicity in HepG2 cells. *J. Phys. Conf.* 1374 (1), 012009.
- Luo, R., Wang, Y., Xu, P., Cao, G., Zhao, Y., Shao, X., Li, Y.X., Chang, C., Peng, C., Wang, Y.L., 2016. Hypoxia-inducible miR-210 contributes to preeclampsia via targeting thrombospondin type I domain containing 7A. *Sci. Rep.* 6, 19588.
- Misra, P., Kumar, A., Khare, P., Gupta, S., Kumar, N., Dube, A., 2009. Pro-apoptotic effect of the landrace bangla mahoba of *Piper betle* on *Leishmania donovani* may be due to the high content of eugenol. *J. Med. Microbiol.* 58 (8), 1058–1066.
- Nagase, M., Shiota, T., Tsushima, A., Alam, M.M., Fukuoka, S., Yoshizawa, T., Sakato, N., 2002. Molecular mechanism of satratoxin-induced apoptosis in HL-60 cells: activation of caspase-8 and caspase-9 is involved in activation of caspase-3. *Immunol. Lett.* 84 (1), 23–27.
- Nagi, M.N., Almakki, H.A., Sayed-Ahmed, M.M., Al-Bekairi, A.M., 2010. Thymoquinone supplementation reverses acetaminophen-induced oxidative stress, nitric oxide production and energy decline in mice liver. *Food Chem. Toxicol.* 48 (8–9), 2361–2365.
- Ni, H.M., Bockus, A., Boggess, N., Jaeschke, H., Ding, W.X., 2012. Activation of autophagy protects against acetaminophen-induced hepatotoxicity. *Hepatology* 55 (1), 222–232.
- Noh, J.R., Kim, Y.H., Hwang, J.H., Choi, D.H., Kim, K.S., Oh, W.K., Lee, C.H., 2015. Sulforaphane protects against acetaminophen-induced hepatotoxicity. *Food Chem. Toxicol.* 80, 193–200.
- Parikh, H., Pandita, N., Khanna, A., 2015. Phytoextract of Indian mustard seeds acts by suppressing the generation of ROS against acetaminophen-induced hepatotoxicity in HepG2 cells. *Pharm. Biol.* 53 (7), 975–984.
- Pluemsamran, T., Onkokoong, T., Panich, U., 2012. Caffeic acid and ferulic acid inhibit UVA-induced matrix metalloproteinase-1 through regulation of antioxidant defense system in keratinocyte HaCaT cells. *Photobiol.* 88 (4), 961–968.
- Prahastuti, S., Hidayat, M., Hasiana, S.T., Widowati, W., Amalia, A., Qodariah, R.L., Rizal, R., Kusuma, H.S.W., Khoiriyah, Z., 2019. Ethanolic extract of jati belanda (*Guazuma ulmifolia* L.) as therapy for chronic kidney disease in vitro model. *J. Rep. Pharm. Sci.* 8 (2), 229–235.
- Rinanda, T., Alga, D.M., 2012. Antibacterial activity of red betel (*Piper crocatum*) leaf methanolic extracts against methicillin resistant *Staphylococcus aureus*. *Proceedings of The 2nd Annual Int. Conference, Syiah Kuala University 2012 & The 8th IMT-GT Uninet Biosciences Conference-Life Sci.* 2 (1), 270–275.
- Roh, T., De, U., Lim, S.K., Kim, M.K., Choi, S.M., Lim, D.S., Yoon, S., Kacew, S., Kim, H.S., Lee, B.M., 2018. Detoxifying effect of pyridoxine on acetaminophen-induced hepatotoxicity via suppressing oxidative stress injury. *Food Chem. Toxicol.* 114, 11–22.
- Salminen, W.F., Yang, X., Shi, Q., Greenhaw, J., Davis, K., Ali, A.A., 2012. Green tea extract can potentiate acetaminophen-induced hepatotoxicity in mice. *Food Chem. Toxicol.* 50 (5), 1439–1446.
- Smith, A.K., Petersen, B.K., Ropella, G.E., Kennedy, R.C., Kaplowitz, N., Ookhtens, M., Hunt, C.A., 2016. Competing mechanistic hypotheses of acetaminophen-induced hepatotoxicity challenged by virtual experiments. *PLoS Comput. Biol.* 12 (12), 1005253.
- Swapna, N.L., Ammani, K., Saripalli, H.P., 2012. Antioxidant activity of makkathotapapada leaves of piper betel L. *Cv. Kapoori. Free Radic. Antioxidants* 2 (4), 68–72.
- Truong, V.L., Ko, S.Y., Jun, M., Jeong, W.S., 2016. Quercitrin from *Toona sinensis* (Juss.) M. Roem. attenuates acetaminophen-induced acute liver toxicity in HepG2 cells and mice through induction of antioxidant machinery and inhibition of inflammation. *Nutrition* 8 (7), 431.
- Ugusman, A., Zakaria, Z., Hui, C.K., Nordin, N.A.M.M., 2011. *Piper sarmentosum* inhibits ICAM-1 and Nox4 gene expression in oxidative stress-induced human umbilical vein endothelial cells. *BMC Complement. Altern. Med.* 11 (1), 1–8.
- Uzi, D., Barda, L., Scaiewicz, V., Mills, M., Mueller, T., Gonzalez-Rodriguez, A., Valverde, A.M., Iwawaki, T., Nahmias, Y., Xavier, R., Chung, R.T., 2013. CHOP is a critical regulator of acetaminophen-induced hepatotoxicity. *J. Hepatol.* 59 (3), 495–503.
- Uzkeser, M., Karakus, E., Albayrak, A., Kiki, I., Bayir, Y., Cadirci, E., Unal, D., Halici, Z., Karadeniz, A., 2012. Protective effect of *Panax ginseng* against N-acetyl-p-aminophenol-induced hepatotoxicity in rats. *African J. Pharm. Pharmacol.* 6 (36), 2634–2642.
- Wang, W., Guan, C., Sun, X., Zhao, Z., Li, J., Fu, X., Qiu, Y., Huang, M., Jin, J., Huang, Z., 2016. Tanshinone IIA protects against acetaminophen-induced hepatotoxicity via activating the Nrf2 pathway. *Phytomedicine* 23 (6), 589–596.
- Widowati, W., Wijaya, L., Wargasetia, T.L., Bachtiar, I., Yellianty, Y., Laksmiawati, D.R., 2013. Antioxidant, anticancer and apoptosis-inducing effects of piper extracts in HeLa cells. *J. Exp. Integr. Med.* 3 (3), 225–230.
- Widowati, W., Prahastuti, S., Ekayanti, N.L.W., Munshy, U.Z., Kusuma, H.S.W., Wibowo, S.H.B., Amalia, A., Widodo, W.S., Rizal, R., 2019a. Anti-inflammation assay of black soybean extract and its compounds on lipopolysaccharide-induced RAW 264.7 cell. *J. Phys. Conf.* 1374, 012052.
- Widowati, W., Jasaputra, D.K., Onggowidjaja, D., Sumitro, S.B., Widodo, M.A., Afifah, E., Rihibiha, D.D., Rizal, R., Amalia, A., Kusuma, H.S.W., Murti, H., Bachtiar, I., 2019b.

- Effects of conditioned medium of co-culture IL-2 induced NK cells and human wharton's jelly mesenchymal stem cells (hWJMSCs) on apoptotic gene expression in a breast cancer cell line (MCF-7). *J. Math. Fund. Sci.* 51 (3), 205–224.
- Wulandari, N., Meiftasari, A., Fadliyah, H., Jenie, R.I., 2018. Red betel leaves methanolic extract (*Piper crocatum* Ruiz & Pav.) increases cytotoxic effect of doxorubicin on WiDr colon cancer cells through apoptosis induction. *Indones. J. Cancer Chemoprevention.* 9 (1), 1–8.
- Yogalakshmi, B., Viswanathan, P., Anuradha, C.V., 2010. Investigation of antioxidant, anti-inflammatory and DNA-protective properties of eugenol in thioacetamide-induced liver injury in rats. *Toxicology* 268 (3), 204–212.
- Yuan, J., Ge, K., Mu, J., Rong, J., Zhang, L., Wang, B., Wan, J., Xia, G., 2016. Ferulic acid attenuated acetaminophen-induced hepatotoxicity through down-regulating the cytochrome P 2E1 and inhibiting toll-like receptor 4 signaling-mediated inflammation in mice. *American J. Transl. Res.* 8 (10), 4205.
- Zulharini, M., Sutejo, I.R., Fadliyah, H., Jenie, R.I., 2018. Methanolic extract of red betel leaves (*Piper crocatum* Ruiz & Pav) perform cytotoxic effect and antimigration activity toward metastatic breast cancer. *Indones. J. Cancer Chemoprevention.* 8 (3), 94–100.

articleHepatotoxicity
prevention in Acetaminophen-
induced HepG2 cells by
redbetel (Piper crocatumRuiz
and Pav) extract from Indonesia
via antioxidant, anti-
inflammatory, and anti-necrotic

by Wahyu Widowati

Submission date: 28-Oct-2021 04:17AM (UTC+0700)

Submission ID: 1685937903

File name: 1-s2.0-S2405844020324634-main.pdf (862.34K)

Word count: 5232

Character count: 28682



Contents lists available at ScienceDirect

Heliyon

journal homepage: www.cell.com/heliyon

Research article

Hepatotoxicity prevention in Acetaminophen-induced HepG2 cells by red betel (*Piper crocatum* Ruiz and Pav) extract from Indonesia via antioxidant, anti-inflammatory, and anti-necroticChrismis Novalinda Ginting^{a,*}, I Nyoman Ehrich Lister^a, Ermi Girsang^a, Wahyu Widowati^b, Dewani Tediana Yusepany^c, Alya Mardhotillah Azizah^c, Hanna Sari Widya Kusuma^c^a Faculty of Medicine, Universitas Prima Indonesia, Jl. Belanga No. 1, Medan 20118, North Sumatera, Indonesia^b Faculty of Medicine, Maranatha Christian University, Jl. Prof. Drg. Surya Sumantri No. 65, Bandung 40163, West Java, Indonesia^c Aretha Medika Utama, Biomolecular Research Center, Jl. Babakan Jeruk II No.9, Bandung 40163, West Java, Indonesia

ARTICLE INFO

Keywords:

Red betel leaves extract
Acetaminophen
HepG2 cells
Hepatoprotective
Biochemistry
Immunology
Inflammation
Natural product
Pharmaceutical science

ABSTRACT

Acetaminophen (APAP) is a widely used analgesic, but it may cause liver injury (hepatotoxicity) via oxidative stress that induced by *N*-acetyl-*p*-benzoquinone imine (NAPQI) in long term usage or overdose. Multiple inflammatory mediators were also found to contribute for this effect. Many medicinal plants was known for its antioxidant and anti-inflammatory activities and one of them is Red betel (*Piper crocatum* Ruiz and Pav) from Indonesia. In this study, the red betel leaves extract (RBLE) protective effect against APAP-induced HepG2 cells was determined. APAP-induced HepG2 as hepatotoxicity cell model was treated with RBLE at 25 and 100 µg/mL. Protective effects of RBLE toward hepatotoxicity were evaluated by several parameters: tumor necrosis factor-α (TNF-α) concentration, reactive oxygen species (ROS) level, live cells percentage, apoptotic cells percentage, necrotic cells percentage, death cells percentage, CYP2E1 and GPX gene expression. The RBLE treatments (both 25 and 100 µg/mL) increased CYP2E1 and GPX gene expression also live cells percentage, while decreased ROS level, TNF-α concentration, also the percentage of death and necrotic cells. Red Betel leaves ethanol extract has hepatoprotective effect via anti-inflammatory, anti-necrotic, and antioxidant potency in liver injury model.

1. Introduction

For many problem of drugs use, liver injury was continues to be a problem. It was represents a major challenge in designing potential therapies (Noh et al., 2015). Acetaminophen (paracetamol, APAP) is considered as first line analgesics. However, excessive use of APAP leads to liver injury even liver failure in human (Ganey et al., 2007; Ni et al., 2012). In small percentage, the cytochrome P450 2E1 (CYP2E1) enzymes was oxidizing the APAP and form *N*-acetyl-*p*-benzoquinone imine (NAPQI), a highly reactive intermediate, which is detoxified by covalent binding with glutathione (GSH). However, in APAP poisoning, it will generates excess NAPQI which evokes the GSH depletion that binds to macromolecules triggering oxidative stress, mitochondrial dysfunction, and ultimately resulting in hepatocellular death (Salminen et al., 2012; Uzi et al., 2013; Yuan et al., 2016). Although the mechanisms underlying hepatotoxicity that induced by APAP still unclear, some evidences was indicate that inflammation mediators such as tumor necrosis factor-α

(TNF-α) also oxidative stress was contribute to the APAP-induced acute liver damage pathology process (Uzkeser et al., 2012; Dragomir et al., 2012).

One of betel in Indonesia namely red betel (*Piper crocatum* Ruiz and Pav) has medicinal function and used as medicine since its introduce as medicinal plants producer in Blunyahrejo (Rinanda and Alga, 2012). It can be used to treat diabetes, gout, hepatitis, hypertension, and eye inflammation (Anugrahwati et al., 2016). In previous study, red betel leaves were found to have some secondary metabolite content like flavonoids, alkaloids, tannins, saponins, triterpenoids steroids, quinones, polyphenolics, and essential oil groups (Arambewela et al., 2005; Wulandari et al., 2018). In addition, red betel contains phenolic compounds in the form of hydrochavicol, cavibetol acetate and eugenol (Swapna et al., 2012; Dervis et al., 2017). In previous studies, red betel leaves extract (RBLE) was shown have anti-inflammatory properties (Misra et al., 2009); antioxidant activity (Lister et al., 2019a); and also

5

* Corresponding author.

E-mail address: chrismis@unprimdn.ac.id (C.N. Ginting).<https://doi.org/10.1016/j.heliyon.2020.e05620>

Received 26 August 2020; Received in revised form 20 October 2020; Accepted 24 November 2020

2405-8440/© 2020 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Table 1. RT-PCR details of β -Actin, CYP2E1, and GPX gene.

Gene Symbols	Primer Sequences (5' to 3') Upper strand: Sense Lower strand: Antisense	Annealing (°C)	Cycle	References
β -Actin	5'-TCTGGACCACACCTTCTACAATG-3' 5'-AGCACAGCCTGGATAGCAACG-3'	63	40	Widowati et al. (2019b) Affiah et al. (2019)
CYP2E1	5'-GTTCITTTGCGGGGACAGAGA-3' 5'-GAGGGTGATGAACCGCTGAA-3'	59	40	Kim et al., 2003
GPX	5'-CCAAGCTCATCACCTGGTCT-3' 5'-TCGATGTCATGGTCTGGAA-3'	59	40	Ugusman et al. (2011)

have anticancer activity especially cervical cancer (Widowati et al., 2013) and breast cancer (Zulharini et al., 2018).

In this study, RBLE potential to suppress liver injury in APAP-induced HepG2 cells was conducted. The parameters that observed in this study was Reactive Oxygen Species (ROS) level; TNF- α level; Cytochrome P450 Family 2 Subfamily E Member 1 (CYP2E1) and Glutathione Peroxidase (GPX) gene expression; apoptotic, necrotic cells, and death cells percentage.

2. Materials and methods

2.1. Preparation of red betel leaves extract

The red betel (*P. crocatum* Ruiz and Pav) leaves that used in this study was obtained from Pabuaran Cilendek Timur, Indonesia and has been identified by Herbarium Bogoriense, Botanical Field Research Center for Biology-Indonesian Institute of Science, Indonesia. RBLE preparation was done by using maceration method. RBLE was obtained from our previous research and stored at -20 °C (Lister et al., 2019a, 2019b).

2.2. HepG2 cells culture and APAP-Induced HepG2

The cells that used in this study is human hepatocellular carcinoma (HepG2) cell line (ATCC, HB-8065TM) from Aretha Medika Utama Biomolecular and Biomedical Research Center, Bandung, Indonesia. It was grown in complete medium with composition: Modified Eagle Medium (MEM) (Biowest, L0416-500), fetal bovine serum (FBS) (Biowest, S1810) as much as 10% (v/v), antibiotic-antimycotic (Gibco, 15240062) as much as 1% (v/v), also nanomycopolitine (Biowest, LX16) addition) as much as 1% of (v/v). Acetaminophen (Sigma Aldrich, A7085) with concentration at 40 mM was used to induce the hepatotoxicity. When the cells were confluent, it was rinsed using PBS and detached using trypsin-EDTA (Gibco, 25200072) with incubation at 37 °C. In 6 well plates, the cells was seeded (5×10^5 cells/well) and then incubated at the same temperature with 5% of CO₂ for 24 h. The cells was induced by RBLE and incubated again for 24 h. According to the treatment, it was divided into 5 groups: I) Normal Cells; II) DMSO1%; III) APAP 40 mM; IV) APAP 40 mM + RBLE 25 μ g/mL; V) APAP 40 Mm + RBLE 100 μ g/mL. Then it was centrifuged at 1600 rpm for 10 min. The supernatant was collected as sample for the Elisa assay (Luo et al., 2016; Aouache et al., 2018; Lister et al., 2019b).

2.3. Total protein assay

Bovine Serum Albumin (BSA) (Sigma Aldrich, A9576) was used as standard in this method. Briefly standard solutions as much as 20 μ L also same volume for the samples was mixed with Quick Start Dye Reagen 1X (Biorad, 5000205) as much as 200 μ L into each well in 96 well plate. The mixture then incubated at room temperature for around 5 min. The wavelength at 595 nm was used to determine the mixture absorbance by using microplate reader (MultiskanTM GO Micro plate Spectrophotometer, Thermo Scientific, 51119300) at 595 nm. The result from this assay was used for normalization of TNF- α data calculation (Pluemsamran et al., 2012; Widowati et al., 2019a).

2.4. TNF- α assay

This assay was measured using ELISA assay (BioLegend, 421701) and according to the manufacturer's kit manual. Based on the manual, wavelength at 450 nm was used to determine the absorbance using microplate reader (Widowati et al., 2019a).

2.5. Apoptotic activity assay

The assay was conducted using methods that reported by Widowati et al. (2019b). Treated and control HepG2 cells were washed using PBS 1x and harvested using trypsin-EDTA for apoptotic assay. The pellet was washed using Annexin Binding Buffer 1X (Miltenyi Biotec, 130-092-820) 500 μ L and stained with Annexin V-FITC (BioLegend, 79998) and Propidium Iodide (BioLegend, 79997). Cells were incubated at 37 °C for 30 min in the dark. Cells were later suspended in Annexin Binding Buffer 1x. The HepG2 cells apoptotic percentage were analyzed using MACSquant Analyzer 10 (Miltenyi Biotec).

2.6. Reactive oxygen species (ROS) assay

HepG2 cells were digested with trypsin-EDTA after cultured around 7 days and 2.5×10^4 cells/0.5 mL were incubated for 45 min in 20 μ M DCF-DA at 37 °C and incubated again for 4 h in RBLE. Based on Prastuti et al. (2019) and Girsang et al. (2019), the 2', 7'-dichlorofluorescein diacetate (DCFDA)-Cellular Reactive Oxygen Species Detection Assay Kit (Abcam, ab113851) was used to measured intracellular ROS with modifications.

Table 2. RNA concentration and purity.

No.	Sample	Concentration (ng/ μ L)	Purity (Absorbance 260/280)
1.	Control cells	92.90	2.3212
2.	Positive control	90.10	2.0904
3.	Positive control + RBLE 25 μ g/mL	36.20	1.9676
4.	Positive control + RBLE 100 μ g/mL	40.00	2.0366




Figure 1. RBLE treatment effect toward TNF- α concentration in APAP-induced HepG2 cells as hepatotoxicity model. (A) TNF- α concentration (pg/mL) on hepatotoxicity model. (B) TNF- α concentration (pg/mg protein) on hepatotoxicity model. *Data was included as mean \pm standard deviation. I) Normal cells as negative control; II) Normal cells + DMSO 1% as vehicle control; III) APAP-induced cells (Positive control); IV) Positive control + RBLE 25 μ g/mL; V) Positive control + RBLE 100 μ g/mL. Significance among treatments toward TNF- α concentration were analyzed with Kruskal Wallis and Post Hoc Test Mann Whitney using SPSS software (version 20.0). The data were presented as mean \pm standard deviation. Significance was presented as different letters (a,b) based on Tukey HSD post hoc test ($P < 0.05$).

2.7. The expression of GPX and CYP2E1 gene assay

Cells that has been harvested was processed for RNA isolation that will be used for futher assay. It was done by using the Aurum™ Total RNA mini Kit (Bio-Rad, 732-6820). RT-qPCR (Clever, GTC96S) was used to analyze the gene expression include the β -actin gene that constitutively expressed (Afifah et al., 2019; Widowati et al., 2019b). Table 1 was shown the primer sequence and Table 2 was shown RNA concentration and purity.

2.8. Statistical analysis

All data were obtained after doing it in triplicate. When the data has normal distribution, it was analyzed using ANOVA and Post Hoc Test using Tukey HSD with $p < 0.05$ while data didn't has normally distributed were analyzed with Kruskal Wallis and Post Hoc Test Mann Whitney using SPSS software (version 20.0). The data were presented as mean \pm standard deviation.




Figure 2. RBLE effect toward apoptotic, necrotic, dead cells in hepatotoxicity model. (A) Live cells on hepatotoxicity model. (B) Early apoptotic on hepatotoxicity model. (C) Late apoptotic on hepatotoxicity model. (D) Necrotic on hepatotoxicity model. *Data was included as mean \pm standard deviation. I) Normal cells as negative control; II) Normal cells + DMSO 1% as vehicle control; III) APAP-induced cells (Positive control); IV) Positive control + RBLE 25 μ g/mL; V) Positive control + RBLE 100 μ g/mL. There are significant different between all groups based on Kruskal-Wallis Test ($P < 0.05$) and Mann-Whitney Test ($P < 0.05$). It was marked as single star (*) marks for the statistical difference between positive control and negative control while the hashtag (#) mark for statistical difference between treatment and positive control.




Figure 3. RBLE effect toward ROS level in hepatotoxicity model. *Data was included as mean \pm standard deviation. I) Normal cells (Negative control); II) Normal cells + DMSO 1%; III) APAP-induced cells (Positive control); IV) Positive control + RBLE 25 μ g/mL; V) Positive control + RBLE 100 μ g/mL. Based on Kruskal-Wallis Test ($P < 0.05$), there are significant different among groups. It was marked as single star (*) for statistical difference between positive control and negative control also hashtag (#) for statistical difference between treatment and positive control.

3. Result

3.1. RBLE effect towards TNF- α concentration in APAP-induced HepG2 cells

APAP increased the TNF- α concentration in HepG2 cells. When RBLE treatment was added, it was found that RBLE can decrease the TNF- α concentration (Figure 1). Based on the result, RBLE has potential to suppress the TNF- α production in HepG2 cells that induced by APAP.

3.2. Effect of RBLE towards apoptotic, necrotic, and cell death in APAP-induced HepG2 cells

APAP decreased live cell percentage compared to normal HepG2 cells (Figure 2A). RBLE treatment decreased the percentage of apoptotic and necrotic significantly in APAP-induced HepG2 cells (Figure 2B–D). RBLE treatment can increase the live cells percentage also reduce the percentage of necrotic and dead cells in HepG2 cells that induced by APAP.




Figure 4. RBLE effect toward the expression of CYP2E1 gene in hepatotoxicity model. *Data was included as mean \pm standard deviation. I) Normal cells as negative control; II) Normal cells + DMSO 1%; III) APAP-induced cells (Positive control); IV) Positive control + RBLE 25 μ g/mL; V) Positive control + RBLE 100 μ g/mL. Based on ANOVA ($P < 0.05$) and Games-Howell ($P < 0.05$), there are significant different among all groups. It was marked as single star (*) marks for the difference between positive control and negative control also hashtag (#) mark for the difference between treatment and positive control.




Figure 5. RBLE effect toward the expression of GPX gene in hepatotoxicity model. *Data was included as mean \pm standard deviation. I) Normal cells (Negative control); II) Normal cells + DMSO 1%; III) APAP-induced cells (Positive control); IV) Positive control + RBLE 25 μ g/mL; V) Positive control + RBLE 100 μ g/mL. Based on Kruskal-Wallis Test ($P < 0.05$) and Mann-Whitney Test ($P < 0.05$), there are significant different between group. It was marked as single star (*) marks as difference between positive control and negative control while hashtag (#) mark as difference between treatment and positive control.

3.3. RBLE effect towards ROS level in liver injury model

ROS level increased significantly after APAP induction and reduced significantly when injured HepG2 cells were treated with RBLE (Figure 3). RBLE had potential to decrease ROS level in liver injury model.

3.4. RBLE effect on CYP2E1 and GPX gene expression in liver injury model

CYP2E1 gene expression decreased significantly in APAP-induced HepG2 cells. RBLE treatments increased the CYP2E1 gene expression significantly compared to the APAP-induced HepG2 cells group (Figure 4). GPX gene expression decreased in APAP-induced HepG2 cells. RBLE treatments could increase the GPX gene expression significantly (Figure 5). RBLE treatments had ability to increase the CYP2E1 and GPX gene expression.

4. Discussions

Betel leaves had been known to contain many active compounds, mainly hydrochavicol, cavibetol acetate and eugenol (Begam et al., 2018). Based on previous study, it had been demonstrated that red betel leaves extract, along with its active constituents: eugenol and hydroxychavicol, can scavenge H_2O_2 and DPPH also reducing FRAP and ABTS radicals that indicated their antioxidant activity (Lister et al., 2019a). Eugenol also had been reported could decrease the ALT and AST activities and LDH level in liver injury model that induced by APAP (Lister et al., 2019b).

The presence of APAP toxic metabolite NAPQI caused Kupffer cells activation that leads to TNF- α release (Legert et al., 2015). TNF- α , one of inflammatory cytokine, involved in oxidative stress injury (Barman et al., 2016; Jaeschke et al., 2012). It was mediated by death receptor pathway apoptosis by activating caspase 3 that act as a central effector to cleave various cellular substrates and trigger cell apoptosis eventually (Nagase et al., 2002; Truong et al., 2016). While apoptosis and necrosis frequently coexist in liver pathological conditions and the cell death balance may be dictated by the particular insult (Antoine et al., 2010).

RBLE treatment was found can decrease the TNF- α level in liver injury model based on the study result. One of active compound in RBLE, eugenol, had been studied have effect on reduction of inflammatory cells infiltration and generation of cytokines from Kupffer cells include ability to suppress TNF- α level in liver injury model (Yogalakshmi et al., 2010).




Figure 6. Proposed RBLE hepatoprotective mechanism in liver injury model. *CYP2E1 act to transform APAP to NAPQI. It induce GSH depletion then induce production of ROS. The excessive of ROS decrease the GPX gene expression leads to increase cell death. NAPQI activate the kupper cell and leads to TNF- α production that induce the JNK signaling pathways that also increase ROS level and leads to upregulate cell necrosis; increase inflammation; and induce cell death. While JNK induced the Bcl-2 down-regulation and Bax up-regulation, resulting in activation of caspase 9 and caspase 3 that leads to apoptosis cells. The RBLE treatments could inhibit the excessive ROS and TNF- α . It also could lowering the necrosis and apoptosis that leads to lowering inflammation. RBLE treatments decrease the cell death and increase the survival hepatic cells.

Phenolic compound had anti-inflammatory effect as another study from Yuan et al. (2016) also stated that a phenolic compound ferulic acid could decrease the TNF- α level in mice induced with APAP.

Figure 2 shows that the APAP induction increased the apoptotic, necrotic, and death cells percentage, while RBLE treatments had successfully reduce death cells and maintain live cells at higher level. This data was in line with previous research that less apoptotic cells were seen in ferulic acid treatment in injury liver model (Yuan et al., 2016).

In APAP-induced hepatotoxicity model, oxidative stress played an important role and it was characterized by ROS accumulation (Nagi et al., 2010; Du et al., 2016). NAPQI, a reactive metabolite formed from APAP, could react rapidly with GSH and aggravating oxidative stress in conjunction with mitochondrial dysfunction that induced hepatocellular damage (Smith et al., 2016; Kang et al., 2017). The enzymatic antioxidant defense system known can detoxified ROS. Previous study exhibited that RBLE had antioxidant potential (Lister et al., 2019a). Based on the result, RBLE proved to suppress the ROS level in liver injury model, this result was in line with Parikh et al. (2015) that found phenolic compounds in *Brassica juncea* hydromethanolic extract such as quercetin and catechin could reduce the ROS level in APAP-induced HepG2 cells. Thus, the RBLE hepatoprotective mechanism might result from diminishing generation of ROS.

In the metabolism of endogenous and exogenous compounds wide variety, CYP2E1 has important functions that relevant to chemical toxicity and carcinogenesis in liver (Gonzalez, 2007). ROS was one of compounds that generated by CYP2E1 that increase mitochondrial membrane permeability and lipid peroxidation, which induce apoptosis via pro-apoptotic factors release and activate caspase 3 (Lee and Wei, 2007). Based on the result, RBLE treatments decreased CYP2E1 gene expression in APAP-induced HepG2 cells, probably by its high phenolic compounds. This result was in line with previous research that in APAP-induced hepatotoxicity, the ferulic acid could inhibit the up-regulation of CYP2E1 expression (Yuan et al., 2016).

The enzymatic antioxidant defense system primary part against oxidative stress is GPX that directly eliminating ROS (Truong et al., 2016). When free radicals formed rapidly, GPX functions will become inefficient and leads to hepatocytes damage (Roh et al., 2018). GPX level can be used as indicator of the oxidative stress response (Wang et al., 2016). Based on the result, it was shown that APAP could decrease the GPX expression, however RBLE treatments could counter this effect. It was indicated that RBLE can protects cells/livers from APAP-inducer through an antioxidant defense system enhancement. Truong et al. (2016) also stated that a phenolic compound, mainly quercitrin could restore GPX expression and attenuates APAP-induced liver damage.

Based on this study, RBLE was shown have antioxidant, anti-necrotic, and anti-inflammatory activities. Its mechanism as hepatoprotective agent in live injury was shown in Figure 6 that proposed by us based on the study result and literature review.

5. Conclusion

Red betel leaves extract treatments could reduce TNF- α level, reduce cell's apoptosis and increase live cells percentage, reduce intracellular ROS, reduce CYP2E1 and increase GPX level in HepG2 cells. This marked the hepatoprotective potential of RBLE through antioxidant, anti-necrotic, and anti-inflammatory activities. Further research on *in vivo* model is needed to confirm current result.

Declarations

Author contribution statement

C.N. Ginting, I.N.E. Lister and E. Girsang: Conceived and designed the experiments.

W. Widawati: Conceived and designed the experiments; Analyzed and interpreted the data.

D.T. Yusepany: Performed the experiments; Contributed reagents, materials, analysis tools or data.

A.M. Azizah: Analyzed and interpreted the data; Wrote the paper.

H.S.W. Kusuma: Contributed reagents, materials, analysis tools or data; Wrote the paper.

Funding statement

This work was funded by Universitas Prima Indonesia, Medan, Indonesia.

7 Declaration of interests statement

The authors declare no conflict of interest.

Additional information

No additional information is available for this paper.

Acknowledgements

We extend our gratitude to Jenifer Kiem Aviani, Rr. Anisa Siwanti Handayan, Kamila Yashfa Gunawan, Dwi Surya Artie, and Seila Arumwardana from Aretha Medika Utama, Biomolecular and Biomedical Research Center for their technical support.

References

- Affiah, E., Moez, T., Sandra, F., Arumwardana, S., Rihibiha, D.D., Nufus, H., Rizal, R., Amalia, A., Bachtiar, I., Murti, H., Widowati, W., 2019. Induction of matrix metalloproteinases in chondrocytes by Interleukin IL-1 β as an osteoarthritis model. *J. Math. Fund. Sci.* 51 (2), 103–111.
- Antoine, D.J., Williams, D.P., Kipar, A., Laverty, H., Park, B.K., 2010. Diet restriction inhibits apoptosis and HMGB1 oxidation and promotes inflammatory cell recruitment during acetaminophen hepatotoxicity. *Mol. Med.* 16 (11–12), 479–490.
- Anugrahwati, M., Purwaningsih, T., Mangalarini, J.A., Alnavis, N.B., Wulandari, D.N., Pranowo, H.D., 2016. Extraction of ethanolic extract of red betel leaves and its cytotoxicity test on HeLa cells. *Proc. Eng.* 148, 1402–1407.
- Aouache, R., Biquard, L., Vaiman, D., Miralles, F., 2018. Oxidative stress in preeclampsia and placental diseases. *Int. J. Mol. Sci.* 19 (5), 1496.
- Arambewela, L.S.R., Arawawala, L.D.A.M., Ratnasooriya, W.D., 2005. Antidiabetic activities of aqueous and ethanolic extracts of *Piper betle* leaves in rats. *J. Ethnopharmacol.* 102 (2), 239–245.
- Barman, P.K., Mukherjee, R., Prusty, B.K., Suklabaidya, S., Senapati, S., Ravindran, B., 2016. Chitoheaxose protects against acetaminophen-induced hepatotoxicity in mice. *Cell Death Dis.* 7 (5), e2224.
- Begam, K.M.F., Ravichandran, P., Manimekalai, V., 2018. Phytochemical analysis of some selected varieties of *Piper betle* L. *Int. J. Curr. Pharmaceut. Res.* 10 (2), 89–93.
- Dervis, E., Yurt Kilar, A., Medine, E.I., Tekin, V., Cetkin, B., Uygur, E., Muftuler, F.Z.B., 2017. In vitro incorporation of radioiodinated eugenol on adenocarcinoma cell lines (Caco2, MCF7, and PC3). *Cancer Biother. Radiopharmaceutic.* 32 (3), 75–81.
- Dragomir, A.C., Sun, R., Mishin, V., Hall, L.B., Laskin, J.D., Laskin, D.L., 2012. Role of galectin-3 in acetaminophen-induced hepatotoxicity and inflammatory mediator production. *Toxicol. Sci.* 127 (2), 609–619.
- Du, K., Ramachandran, A., Jaeschke, H., 2016. Oxidative stress during acetaminophen hepatotoxicity: sources, pathophysiological role and therapeutic potential. *Redox Biol.* 10, 148–156.
- Ganey, P.E., Luyendyk, J.P., Newport, S.W., Eagle, T.M., Maddox, J.F., Mackman, N., Roth, R.A., 2007. Role of the coagulation system in acetaminophen-induced hepatotoxicity in mice. *Hepatology* 46 (4), 1177–1186.
- Girsang, E., Lister, I.N.E., Ginting, C.N., Nasution, S.L., Suhartina, S., Munshy, U.Z., Rizal, R., Widowati, W., 2019. Antioxidant and anti-inflammatory activity of chlorogenic acid on lead-induced fibroblast cells. *J. Phys. Conf.* 1374, 012006.
- Gonzalez, F.J., 2007. The 2006 Bernard B. Brodie Award Lecture. Cyp2e1. *Drug Metabol. Dispos.* 35 (1), 1–8.
- Jaeschke, H., Williams, C.D., Ramachandran, A., Bajt, M.L., 2012. Acetaminophen hepatotoxicity and repair: the role of sterile inflammation and innate immunity. *Liver Int.* 32 (1), 8–20.
- Kang, E.S., Lee, J., Homma, T., Kurahashi, T., Kobayashi, S., Nabeshima, A., Yamada, S., Seo, H.G., Miyata, S., Sato, H., Fujii, J., 2017. xCT deficiency aggravates acetaminophen-induced hepatotoxicity under inhibition of the transsulfuration pathway. *Free Radic. Res.* 51 (1), 80–90.
- Kim, S.S., Oh, O.J., Min, H.Y., Park, E.J., Kim, Y., Park, H.J., Han, Y.N., Lee, S.K., 2003. Eugenol suppresses cyclooxygenase-2 expression in lipopolysaccharide-stimulated mouse macrophage RAW264.7 cells. *Life Sci.* 73 (3), 337–348.
- Lee, H.C., Wei, Y.H., 2007. Oxidative stress, mitochondrial DNA mutation, and apoptosis in aging. *Exp. Biol. Med.* 232 (5), 592–606.
- Legert, K.G., Tsilingaridis, G., Remberger, M., Ringden, O., Heimdahl, A., Yucel-Lindberg, T., Dahllof, G., 2015. The relationship between oral mucositis and levels of pro-inflammatory cytokines in serum and in gingival crevicular fluid in allogeneic stem cell recipients. *Supp. Care Cancer* 23 (6), 1749–1757.
- Lister, I.N.E., Ginting, C.N., Girsang, E., Amansyah, A., Marpaung, H.H., Sinaga, A.P.F., Handayani, R.A.S., Rizal, R., 2019a. Antioxidant properties of red betel (*Piper crocatum*) leaf extract and its compounds. *J. Nat. Remedies* 19 (4), 198–205.
- Lister, I.N.E., Ginting, C.N., Girsang, E., Amansyah, A., Chiuman, L., Yanti, N.L.W.E., Rizal, R., Widowati, W., 2019b. Hepatoprotective effect of eugenol on acetaminophen-induced hepatotoxicity in HepG2 cells. *J. Phys. Conf.* 1374 (1), 012009.
- Luo, R., Wang, Y., Xu, P., Cao, G., Zhao, Y., Shao, X., Li, Y.X., Chang, C., Peng, C., Wang, Y.L., 2016. Hypoxia-inducible miR-210 contributes to preeclampsia via targeting thrombospondin type I domain containing 7A. *Sci. Rep.* 6, 19588.
- Misra, P., Kumar, A., Khare, P., Gupta, S., Kumar, N., Dube, A., 2009. Pro-apoptotic effect of the lardrace bangla mahoba of *Piper betle* on *Leishmania donovani* may be due to the high content of eugenol. *J. Med. Microbiol.* 58 (8), 1058–1066.
- Nagase, M., Shiota, T., Tsushima, A., Alam, M.M., Fukuoka, S., Yoshizawa, T., Sakato, N., 2002. Molecular mechanism of satratoxin-induced apoptosis in HL-60 cells: activation of caspase-8 and caspase-9 is involved in activation of caspase-3. *Immunol. Lett.* 84 (1), 23–27.
- Nagi, M.N., Almaki, H.A., Sayed-Ahmed, M.M., Al-Bekairi, A.M., 2010. Thymoquinone supplementation reverses acetaminophen-induced oxidative stress, nitric oxide production and energy decline in mice liver. *Food Chem. Toxicol.* 48 (8–9), 2361–2365.
- Ni, H.M., Bockus, A., Boggess, N., Jaeschke, H., Ding, W.X., 2012. Activation of autophagy protects against acetaminophen-induced hepatotoxicity. *Hepatology* 55 (1), 222–232.
- Noh, J.R., Kim, Y.H., Hwang, J.H., Choi, D.H., Kim, K.S., Oh, W.K., Lee, C.H., 2015. Sulforaphane protects against acetaminophen-induced hepatotoxicity. *Food Chem. Toxicol.* 80, 193–200.
- Parikh, H., Pandita, N., Khanna, A., 2015. Phytoextract of Indian mustard seeds acts by suppressing the generation of ROS against acetaminophen-induced hepatotoxicity in HepG2 cells. *Pharm. Biol.* 53 (7), 975–984.
- Pluemsamran, T., Onkkoosong, T., Panich, U., 2012. Caffeic acid and ferulic acid inhibit UVA-induced matrix metalloproteinase-1 through regulation of antioxidant defense system in keratinocyte HaCaT cells. *Photochem. Photobiol.* 88 (4), 961–968.
- Prahastuti, S., Hidayat, M., Hasiana, S.T., Widowati, W., Amalia, A., Qodariah, R.L., Rizal, R., Kusuma, H.S.W., Khoiriyah, Z., 2019. Ethanolic extract of jati belanda (*Guazuma ulmifolia* L.) as therapy for chronic kidney disease in vitro model. *J. Rep. Pharm Sci.* 8 (2), 229–235.
- Rinanda, T., Alga, D.M., 2012. Antibacterial activity of red betel (*Piper crocatum*) leaf methanolic extracts against methicillin resistant *Staphylococcus aureus*. *Proceedings of The 2nd Annual Int. Conference, Syiah Kuala University 2012 & The 8th IMT-GT Uninet Biosciences Conference-Life Sci. Eng.* 2 (1), 270–275.
- Roh, T., De, U., Lim, S.K., Kim, M.K., Choi, S.M., Lim, D.S., Yoon, S., Kacew, S., Kim, H.S., Lee, B.M., 2018. Detoxifying effect of pyridoxine on acetaminophen-induced hepatotoxicity via suppressing oxidative stress injury. *Food Chem. Toxicol.* 114, 11–22.
- Salminen, W.F., Yang, X., Shi, Q., Greenhaw, J., Davis, K., Ali, A.A., 2012. Green tea extract can potentiate acetaminophen-induced hepatotoxicity in mice. *Food Chem. Toxicol.* 50 (5), 1439–1446.
- Smith, A.K., Petersen, B.K., Ropella, G.E., Kennedy, R.C., Kaplowitz, N., Ookhtens, M., Hunt, C.A., 2016. Competing mechanistic hypotheses of acetaminophen-induced hepatotoxicity challenged by virtual experiments. *PLoS Comput. Biol.* 12 (12), 1005253.
- Swapna, N.L., Ammani, K., Saripalli, H.P., 2012. Antioxidant activity of mokkathatapapada leaves of piper betel L. Cv. Kapoori. *Free Radic. Antioxidants* 2 (4), 68–72.
- Truong, V.L., Ko, S.Y., Jun, M., Jeong, W.S., 2016. Quercitrin from *Toona sinensis* (Juss.) M. Roem. attenuates acetaminophen-induced acute liver toxicity in HepG2 cells and mice through induction of antioxidant machinery and inhibition of inflammation. *Nutrition* 8 (7), 431.
- Ugusman, A., Zakaria, Z., Hui, C.K., Nordin, N.A.M.M., 2011. *Piper sarmentosum* inhibits ICAM-1 and Nox4 gene expression in oxidative stress-induced human umbilical vein endothelial cells. *BMC Complement. Altern. Med.* 11 (1), 1–8.
- Uzi, D., Barda, L., Scaiewicz, V., Mills, M., Mueller, T., Gonzalez-Rodriguez, A., Valverde, A.M., Iwawaki, T., Nahmias, Y., Xavier, R., Chung, R.T., 2013. CHOP is a critical regulator of acetaminophen-induced hepatotoxicity. *J. Hepatol.* 59 (3), 495–503.
- Uzkeser, M., Karakas, E., Albayrak, A., Kiki, I., Bayir, Y., Cadirci, E., Unal, D., Halici, Z., Karadeniz, A., 2012. Protective effect of *Panax ginseng* against N-acetyl-p-aminophenol-induced hepatotoxicity in rats. *African J. Pharm. Pharmacol.* 6 (36), 2634–2642.
- Wang, W., Guan, C., Sun, X., Zhao, Z., Li, J., Fu, X., Qiu, Y., Huang, M., Jin, J., Huang, Z., 2016. Tanshinone IIA protects against acetaminophen-induced hepatotoxicity via activating the Nrf2 pathway. *Phytomedicine* 23 (6), 589–596.
- Widowati, W., Wijaya, L., Wargasetia, T.L., Bachtiar, I., Yellianty, Y., Laksmiawati, D.R., 2013. Antioxidant, anticancer and apoptosis-inducing effects of piper extracts in HeLa cells. *J. Exp. Integr. Med.* 3 (3), 225–230.
- Widowati, W., Prahastuti, S., Ekayanti, N.L.W., Munshy, U.Z., Kusuma, H.S.W., Wibowo, S.H.B., Amalia, A., Widodo, W.S., Rizal, R., 2019a. Anti-inflammation assay of black soybean extract and its compounds on lipopolysaccharide-induced RAW 264.7 cell. *J. Phys. Conf.* 1374, 012052.
- Widowati, W., Jasputra, D.K., Onggowidjaja, D., Sumitro, S.B., Widodo, M.A., Affiah, E., Rihibiha, D.D., Rizal, R., Amalia, A., Kusuma, H.S.W., Murti, H., Bachtiar, I., 2019b.

- Effects of conditioned medium of co-culture IL-2 induced NK cells and human wharton's jelly mesenchymal stem cells (hWJMSCs) on apoptotic gene expression in a breast cancer cell line (MCF-7). *J. Math. Fund. Sci.* 51 (3), 205–224.
- Wulandari, N., Meiftasari, A., Fadliyah, H., Jenie, R.L., 2018. Red betel leaves methanolic extract (*Piper crocatum* Ruiz & Pav.) increases cytotoxic effect of doxorubicin on WIDr colon cancer cells through apoptosis induction. *Indones. J. Cancer Chemoprevention.* 9 (1), 1–8.
- Yogalakshmi, B., Viswanathan, P., Anuradha, C.V., 2010. Investigation of antioxidant, anti-inflammatory and DNA-protective properties of eugenol in thioacetamide-induced liver injury in rats. *Toxicology* 268 (3), 204–212.
- Yuan, J., Ge, K., Mu, J., Rong, J., Zhang, L., Wang, B., Wan, J., Xia, G., 2016. Ferulic acid attenuated acetaminophen-induced hepatotoxicity through down-regulating the cytochrome P 2E1 and inhibiting toll-like receptor 4 signaling-mediated inflammation in mice. *American J. Transl. Res.* 8 (10), 4205.
- Zulharini, M., Sutejo, I.R., Fadliyah, H., Jenie, R.L., 2018. Methanolic extract of red betel leaves (*Piper crocatum* Ruiz & Pav) perform cytotoxic effect and antimigration activity toward metastatic breast cancer. *Indones. J. Cancer Chemoprevention.* 8 (3), 94–100.

articleHepatotoxicity prevention in Acetaminophen-induced HepG2 cells by redbetel (Piper crocatumRuiz and Pav) extract from Indonesia via antioxidant,anti-inflammatory, and anti-necrotic

ORIGINALITY REPORT

12%

SIMILARITY INDEX

11%

INTERNET SOURCES

9%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	repository.maranatha.edu Internet Source	4%
2	peerj.com Internet Source	2%
3	pubmed.ncbi.nlm.nih.gov Internet Source	1%
4	Submitted to University of Technology, Sydney Student Paper	1%
5	Submitted to Universitas Merdeka Malang Student Paper	1%
6	ri.conicet.gov.ar Internet Source	1%
7	eprints.whiterose.ac.uk Internet Source	1%
8	etd.aau.edu.et Internet Source	1%

9	www.hindawi.com Internet Source	1 %
10	Xing-Yue Xu, Zi Wang, Shen Ren, Jing Leng, Jun-nan Hu, Zhi Liu, Chen Chen, Wei Li. "Improved protective effects of American ginseng berry against acetaminophen-induced liver toxicity through TNF- α -mediated caspase-3/-8/-9 signaling pathways", Phytomedicine, 2018 Publication	1 %
11	coek.info Internet Source	1 %
12	res.mdpi.com Internet Source	1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On

BEALL'S LIST

OF POTENTIAL PREDATORY JOURNALS AND PUBLISHERS

[PUBLISHERS](#)[STANDALONE JOURNALS](#)[VANITY PRESS](#)[CONTACT](#)[OTHER](#)

Potential predatory scholarly open-access journals

Instructions: simply enter the journal's name or its URL in the search box above. If the journal has a publisher that was assessed to be predatory, then it is included on the [Publishers](#) list.

Original list

This is an archived version of the Beall's list – a list of potential predatory journals created by a librarian [Jeffrey Beall](#). We will only update links and add notes to this list.

Last updated January 9, 2017

Update

Here we include journals that were not originally on the Beall's list, but may be predatory.

Last updated February 5, 2021

Original description

This is a list of questionable, scholarly open-access standalone journals. For journals published by a publisher, please look for the publisher on the list of publishers, [here](#). This list is only for single, standalone journals. We recommend that scholars read the available reviews, assessments and descriptions provided here, and then decide for themselves whether they want to

[GO TO UPDATE](#)

BEALL'S LIST

OF POTENTIAL PREDATORY JOURNALS AND PUBLISHERS

[PUBLISHERS](#)[STANDALONE JOURNALS](#)[VANITY PRESS](#)[CONTACT](#)[OTHER](#)

Potential predatory scholarly open-access publishers

Instructions: first, find the journal's publisher – it is usually written at the bottom of the journal's webpage or in the "About" section. Then simply enter the publisher's name or its URL in the search box above. If the journal does not have a publisher use the [Standalone Journals](#) list.

All journals published by a predatory publisher are potentially predatory unless stated otherwise.

Original list

[GO TO UPDATE](#)

This is an archived version of the Beall's list – a list of potential predatory publishers created by a librarian [Jeffrey Beall](#). We will only update links and add notes to this list.

Last updated December 31, 2016

Update

Here we include publishers that were not originally on the Beall's list, but may be predatory.

Last updated March 7, 2021

Excluded – decide after reading

Useful pages

[List of journals falsely claiming to be indexed by DOAJ](#)

[DOAJ: Journals added and removed](#)

[Nonrecommended medical periodicals](#)

[Retraction Watch](#)

[Flaky Academic Journals Blog](#)

[List of scholarly publishing stings](#)

Conferences

[Questionable conferences \[archive\]](#)

[How to avoid predatory conferences](#)

[Flaky Academic Conferences Blog](#)