

Daftar Pustaka

1. Perkeni. *Konsensus Pengelolaan Dan Pencegahan Diabetes Melitus Tipe 2 Di Indonesia 2015*. Vol 1.; 2015. doi:10.1017/CBO9781107415324.004.
2. Of S, Care diabetes M. *STANDARDS OF MEDICAL CARE Standards of Medical Care in Diabetes D 2016*. Vol 39.; 2016.
3. WHO. *GLOBAL REPORT ON DIABETES*.; 2016.
4. Kementerian Kesehatan. *Riset Kesehatan Dasar*.; 2013.
5. Gohel MG, Chacko AN. Serum GGT activity and hsCRP level in patients with type 2 diabetes mellitus with good and poor glycemc control: An evidence linking oxidative stress, inflammation and glycemc control. *J Diabetes Metab Disord*. 2013;12(1):56. doi:10.1186/2251-6581-12-56.
6. Sudoyo A w, Bambang S hadi, Idris A. *Buku Ajar Ilmu Penyakit Dalam (Edisi 6)*. Jakarta: Interna Publishing; 2014.
7. PFÜTZNER A, FORST T. High-Sensitivity C-Reactive Protein as Cardiovascular Risk Marker in Patients with Diabetes Mellitus. *DIABETES Technol Ther*. 2006;8(1):28-37.
8. Siracuse JJ, Chaikof EL. Diabetes and Peripheral Vascular Disease. 2012:13-27. doi:10.1007/978-1-62703-158-5.
9. Hall J. *Guyton and Hall Textbook of Medical Physiology*. Elsevier; 2016.
10. Edition S. *DIABETES*. 2015th ed.; 2015. International Diabetes Federation.
11. DeFronzo R. From the Triumvirate to the Ominous Octet: A New Paradigm for the Treatment of Type 2 Diabetes Mellitus. 2009.
12. Kementerian Kesehatan RI. *Profil Kesehatan Indonesia 2015*.; 2016. doi:351.077 Ind.
13. Tipe DM, Indonesia DI. *KONSENSUS PENGELOLAAN DAN PENCEGAHAN DIABETES MELITUS TIPE 2 DI*.; 2011.
14. Eroschenko V. *Atlas Histologi Difiore (Edisi 11)*.; 2010.

15. Kumar V. *Robbins Basic Pathology 10th Edition*. Elsevier; 2013.
16. Parrinello CM, Lutsey PL, Ballantyne CM, Folsom AR, Pankow JS, Selvin E. Six-year change in high-sensitivity C-reactive protein and risk of diabetes, cardiovascular disease, and mortality. *Am Heart J*. 2015;170(2):380-389.e4. doi:10.1016/j.ahj.2015.04.017.
17. Barbalho SM, Toffano RJ, Bechara MD, et al. Association between Hyperglycemia , C-Reactive Protein and Other Risk Factors in Patients at Cardiovascular Risk. *J Diabetes Mellit*. 2016;(February):16-24. doi:http://dx.doi.org/10.4236/jdm.2016.61003.
18. Lal T, Moses A, Dorairaj VS, Muthukumar S, Banu K. HIGH SENSITIVITY C-REACTIVE PROTEIN in TYPE-2 DIABETES MELLITUS. *Int J Curr Res*. 2015;7(June 2010).
19. Amanullah S, Jarari A, Govindan M, Basha MI, Khatheeja S. Association of hs-CRP with Diabetic and Non-diabetic individuals. *JJBS Jordan J Biol Sci*. 2010;3(1):7-12.
20. Soinio M, Maeniemi J, Laakso M, Lehto S, Ronnema T. High-Sensitivity C-Reactive Protein and Coronary Heart Disease Mortality in Patients With Type 2 Diabetes: A 7-year follow-up study. *Diabetes Care*. 2006;29(2):329-333. doi:10.2337/diacare.29.02.06.dc05-1700.
21. Lima LM, Carvalho M das G, Soares AL, et al. High-sensitivity C-reactive protein in subjects with type 2 diabetes mellitus and/or high blood pressure. *Arq Bras Endocrinol Metabol*. 2007;51(6):956-960. <http://www.ncbi.nlm.nih.gov/pubmed/17934663>.