



UNIVERSITAS
KRISTEN
MARANATHA

Jl. Prof. drg. Surya Sumantri, M.P.H. No. 65
Bandung - 40164, Jawa Barat, Indonesia
Telp: +62 22-201 2186 / 200 3450, ext. 1905 / 1906
Email: fkg@dent.maranatha.edu
www.maranatha.edu

Fakultas Kedokteran Gigi

SURAT TUGAS
No. 075/FKG-UKM/I/2023

Yang bertanda tangan di bawah ini

Nama : Dr. Ignatius Setiawan, drg., M.M.
Jabatan : Dekan Fakultas Kedokteran Gigi UK.Maranatha

Dengan ini menugaskan kepada :

No	Nama Dosen	NIK
1.	Dr. Vinna Kurniawati Sugiaman, drg., M.Kes., PBO., CMC.	120005
2.	Silvia Naliani, drg., Sp.Pros., M.K.G., CMC.	120029

Melaksanakan Publikasi Jurnal "*Fungistatic effect of Moringa oleifera Lam. on the metabolism changes of Candida albicans*" di *Journal of Pharmacy & Pharmacognosy Research* pada Januari 2023.

Demikian agar tugas ini dilaksanakan dengan sebaik-baiknya.

Bandung, 31 Januari 2023
Dekan Fakultas Kedokteran Gigi
Universitas Kristen Maranatha


Dr. Ignatius Setiawan, drg., M.M.
NIK. 120010



JPPRes

Journal of Pharmacy
& Pharmacognosy
Research

CELEBRATING
10 YEARS
2013-2023




Volume 11, Issue 1

J Pharm Pharmacogn Res 11(1), (Jan-Feb) 2023




IFC (**Journal of Pharmacy & Pharmacognosy Research**).

1.- Original Article


Samiaa J. Abdulwahid-Kurdi, Muhsin J. Abdulwahid, Usman Magaji, Zeiad A. Aghwan, Rodziah Atan, Kasrin A. Hamadamin (2023) **Medicinal plants used in Soran district Kurdistan region of Iraq, an ethnobotanicals study.** | [Plantas medicinales utilizadas en el distrito de Soran, región del Kurdistan de Irak, un estudio etnobotánico]. J Pharm Pharmacogn Res 11(1): 1-32. https://doi.org/10.56499/jppres22.1484_11.1.1  jppres_pdf_fre [1.13 Mb]

2.- Original Article

Alberto Calle, Cecilia K. Curi-Borda, Cervando Gutiérrez, Lily Salcedo, Yonny Flores, Giovanna R. Almanza (2023) **Quantification and *in vitro* photo-protective studies of phenolic compounds from *Baccharis papillosa* Rusby.** | [Cuantificación y estudios de fotoprotección *in vitro* de compuestos fenólicos de *Baccharis papillosa* Rusby]. J Pharm Pharmacogn Res 11(1): 33-46. https://doi.org/10.56499/jppres22.1477_11.1.33  jppres_pdf_free [724 Kb]


3.- Original Article

Raghad R. Al-Yousef, Manal M. Abbas, Razan Obeidat, Manal A. Abbas (2023) **Rhamnetir decreases the expression of HMG-CoA reductase gene and increases LDL receptor in HepG2 cells.** | [Ramnetina disminuye la expresión del gen de la HMG-CoA reductasa y


aumenta los receptores de LDL en las células HepG2]. J Pharm Pharmacogn Res 11(1): 47-54. https://doi.org/10.56499/jppres22.1507_11.1.47  jppres_pdf_free [355 Kb]

4.- Original Article

Hendri Susilo, Mochammad Thaha, Budi Susetyo Pikir, Mochamad Yusuf Alsagaff, Satriyo Dwi Suryantoro, Ifan Ali Wafa, Nando Reza Pratama, David Setyo Budi, Bayu Satria Wiratama, Citrawati Dyah Kencono Wungu (2023) **The role of plasma angiotensin-converting enzyme and interleukin-6 levels on the prognosis of non-dialysis chronic kidney disease patients.** | [Papel de los niveles plasmáticos de la enzima convertidora de angiotensina e interleucina-6 en el pronóstico de los pacientes con enfermedad renal crónica no sometidos a diálisis]. J Pharm Pharmacogn Res 11(1): 55-62.


https://doi.org/10.56499/jppres22.1518_11.1.55  jppres_pdf_free [341 Kb]

5.- Original Article

Lizette Gil-del Valle, Rosario Gravier-Hernández, Mario M. Delgado-Guerra, Joniel A. Sánchez-Márquez, Olga E. López-Fernández, Miguel A. Acosta-Suárez, Teresa Rosell-Guerra, Rodolfo Suárez-Iznaga, Raiza Martínez-Casanueva, Zullyt Zamora-Rodriguez, Lid A. Fernández-García, Yusimit Bermudez-Alfonso, María C. Hernández-Gonzalez-Abreu, Gabino Garrido (2023) **Oxidative stress in diverse clinical conditions of SARS-CoV-2 Cuban hospitalized patients.** | [Estrés oxidativo en diferentes condiciones clínicas de pacientes cubanos hospitalizados con SARS-CoV-2]. J Pharm Pharmacogn Res 11(1): 63-75. https://doi.org/10.56499/jppres22.1535_11.1.63  jppres_pdf_free [474 Kb]


6.- Review

Eka Arum Cahyaning Putri, Misbakhul Munir, Hayuris Kinandita Setiawan, Lilik Herawati, Gadis Meinar Sari, Citrawati Dyah Kencono Wungu, Hendri Susilo, Henry Sutanto (2023) **Adverse cardiac events following mRNA COVID-19 vaccination: A systematic review and meta-analysis.** | [Eventos cardíacos adversos tras la vacunación con COVID-19 ARNm: Una revisión sistemática y metaanálisis]. J Pharm Pharmacogn Res 11(1): 76-100.


https://doi.org/10.56499/jppres22.1524_11.1.76  jppres_pdf_free [652 Kb]

7.- Original Article


John Jáuregui-Nongrados, Angel T. Alvarado, Miguel Mucha, Ana M. Muñoz, Haydee Chávez, Aura Molina-Cabrera, Pompeyo A. Cuba-García, Elizabeth J. Melgar-Merino, Mario Bolarte-Arteaga, Jaime A. Mori-Castro (2023) **Bioadsorption of silver ions by calcareous chitin, chitin and chitosan.** | [Bioadsorción de iones de plata por quitina calcárea, quitina y quitosano]. J Pharm Pharmacogn Res 11(1): 101-109.

https://doi.org/10.56499/jppres22.1529_11.1.101  jppres_pdf_free [572 Kb]


8.- Original Article

An Nisa Fithri, Yuyun Yueniwati, I Wayan Arsana, Husnul Khotimah, Wiwit Nurwidyaningtyc (2023) **Epithelial thinning in vaginal atrophy related to lowering of calcitonin gene-related protein, vascular endothelial growth factor, and nerve growth factor expressions in a menopausal rat model.** | [Adelgazamiento epitelial en la atrofia vaginal relacionado con la disminución de las expresiones de la proteína relacionada con el gen de la calcitonina, el factor de crecimiento endotelial vascular y el factor de crecimiento nervioso en un modelo de rata menopáusica]. J Pharm Pharmacogn Res 11(1): 110-116. https://doi.org/10.56499/jppres22.1512_11.1.110  jppres_pdf_free [511 Kb]


9.- Original Article

Hawa Purnama Celala Ary Cane, Musri Musman, Mustanir Yahya, Nurdin Saidi, Darusmar Darusman, Muhammad Nanda, Diva Rayyan Rizki, Kana Puspita (2023) **Phytochemical screening and antibacterial activity of ethnomedicinal plants from Gayo Lues Highland, Indonesia.** | [Cribado fitoquímico y actividad antibacteriana de plantas etnomedicinales del altiplano de Gayo Lues, Indonesia]. J Pharm Pharmacogn Res 11(1): 117-128. https://doi.org/10.56499/jppres22.1526_11.1.117  jppres_pdf_free [709 Kb]


10.- Original Article

Hanifa R. Putri, Rhesi Kristiana, I Wayan Mudianta, Edwin Setiawan, Aty Widyawaruyanti, Nitra Nuengchamng, Nungruthai Suphrom, Suciati Suciati (2023) **Metabolite profile and *in vitro* cholinesterase inhibitory activity of extract and fractions of *Aaptos suberitoides*.** | [Perfil de metabolitos y actividad inhibidora de la colinesterasa *in vitro* de extracto y fracciones de *Aaptos suberitoides*]. J Pharm Pharmacogn Res 11(1): 129-136. https://doi.org/10.56499/jppres22.1511_11.1.129  jppres_pdf_free [490 Kb]

11.- Original Article

Sara Tlemcani, Amal Lahkimi, Nouredine Eloutassi, Ahmed Bendaoud, Anouar Hmamou Hicham Bekkari (2023) **Ethnobotanical study of medicinal plants in the Fez-Meknes region of Morocco.** | [Estudio etnobotánico de plantas medicinales en la región de Fez-Meknes de Marruecos]. J Pharm Pharmacogn Res 11(1): 137-159. https://doi.org/10.56499/jppres22.1459_11.1.137  jppres_pdf_free [890 Kb]


12.- Expert Opinion

Jorge Duconge-Soler, Víctor Mangas Sanjuan, Gledys Reynaldo Fernández (2023) **Conceptos y aplicaciones de los parámetros farmacocinéticos: Una guía para el salón de clases.** | [Concepts and applications of pharmacokinetic parameters: A guide for the classroom]. J Pharm Pharmacogn Res 11(1): 160-178. https://doi.org/10.56499/jppres22.1472_11.1.160  jppres_pdf_free [962 Kb]

13.- Original Article

Basri A. Gani, Cut Soraya, Vinna Kurniawati Sugiaman, Fitri Yunita Batubara, Dharli Syafriz Silvia Naliani, Sri Rezeki, Subhani Jakfar, Muhammad Nazar, Kemala Hayati (2023)


Fungistatic effect of *Moringa oleifera* Lam. on the metabolism changes of *Candida albicans*. | [Efecto fungistático de *Moringa oleifera* Lam. sobre los cambios en el metabolismo de *Candida albicans*]. J Pharm Pharmacogn Res 11(1): 179-190.

https://doi.org/10.56499/jppres22.1533_11.1.179  jppres_pdf_free [853 Kb]

14.- Original Article

Sulistyo Mulyo Agustini, Edi Widjajanto, Muhaimin Rifa'i, Sofia Mubarika Haryana, Nurdian Diana Lyrawati, Usi Sukorini, Noviana Dwi Lestari (2023) **Anti-leukemic activity of *Cyperus rotundus* L. on human acute myeloid leukemia HL-60 cells *in vitro*.** | [Actividad antileucémica de *Cyperus rotundus* L. sobre células humanas de leucemia mieloide aguda HL-60 *in vitro*]. J Pharm Pharmacogn Res 11(1): 191-197.


[Actividad antileucémica de *Cyperus rotundus* L. sobre células humanas de leucemia mieloide aguda HL-60 *in vitro*]. J Pharm Pharmacogn Res 11(1): 191-197.

https://doi.org/10.56499/jppres22.1502_11.1.191  jppres_pdf_free [542 Kb]

15.- Original Article

Zahrotunisa, Silvia Surini (2023) **Advance in transdermal delivery of calcitonin using nanostructured lipid carrier-based emulgel.** | [Avance en la administración transdérmica de calcitonina mediante un emulgel nanoestructurado basado en un portador lipídico]. J Pharm Pharmacogn Res 11(1): 198-207.

[Avance en la administración transdérmica de calcitonina mediante un emulgel nanoestructurado basado en un portador lipídico]. J Pharm Pharmacogn Res 11(1): 198-207.

https://doi.org/10.56499/jppres22.1538_11.1.198  jppres_pdf_free [485 Kb]

© 2023 Journal of Pharmacy & Pharmacognosy Research (JPPRes)



JPPRes

Journal of Pharmacy
& Pharmacognosy
Research

[Home](#) [About](#) [Archive](#) [Authors](#) [Reviewers](#) [Events](#) [Blog](#)

 [Twitter](#)  [LinkedIn](#)  [Facebook](#)



Editorial Board



JPPRes

Journal of Pharmacy
& Pharmacognosy
Research

Gabino Garrido

Editor-in-Chief

[Scholar Google](#)

[Scopus](#)

[ResearchGate](#)

[ORCID](#)

GarVal Editorial Ltda.

[Email](#)

Marisela Valdés González

Executive Editor

[ORCID](#)

GarVal Editorial Ltda.



Editorial and Design Manager

[LinkedIn](#)[Behance](#)[Web](#)

AVAGAX Studio

Antofagasta, Chile

[Email](#)

Editorial Board Members



Antonio Vassallo

Dipartimento di Scienze. Università degli Studi della Basilicata, Potenza, Italia.

[Research Gate](#)[ORCID](#)[Scopus](#)

Bárbara B. Garrido Suárez

Laboratorio de Farmacología y Toxicología, Centro de Investigación y Desarrollo de Medicamentos, La Habana, Cuba.

**Carla Delporte**

Facultad de Ciencias Químicas y Farmacéuticas, Universidad de Chile, Santiago, Chile

[Research Gate](#)[ORCID](#)**Damaris Silveira**

Faculdade de Ciências da Saúde, Universidade de Brasília, Brazil

[Lattes](#)[ORCID](#)**Douglas S. de Almeida Chaves**

Dep. de Ciências Farmacêuticas (DCFAR), Instituto de Ciências Biológicas e da Saúde,
Universidade Federal Rural do Rio de Janeiro, Rodovia, RJ. Brazil

[ResearchGate](#)[ORCID](#)[Lattes](#)**Edgar Pastene**

Universidad de Concepción, Concepción, Chile



Etile Spegazzini

Universidad de Belgrano, Buenos Aires, Argentina

[ResearchGate](#)



Farid Chemat

Université Avignon et des Pays de Vaucluse, Avignon, France

[ResearchGate](#)



Fidel O. Castro

Departamento de Ciencias Pecuarias, Universidad de Concepción, Chillán, Chile

[ResearchGate](#)

[ORCID](#)



Guilherme N. L. do Nascimento

Universidade Federal de Tocantins. Brazil

**Jacqueline Sepúlveda**

Universidad de Concepción, Concepción, Chile

[ResearchGate](#)[ORCID](#)**Jelena Nadinic**

Universidad de Buenos Aires, Argentina.

[ResearchGate](#)[ORCID](#)**José H. Isaza-Martínez**

Universidad del Valle, Cali, Colombia

[ResearchGate](#)[ORCID](#)**Juan C. Sepúlveda-Arias**

Universidad Tecnológica de Pereira, Pereira, Colombia

**Madan Mohan Gupta**

School of Pharmacy, Faculty of Medical Sciences. The University of the West Indies,
Trinidad & Tobago, West Indies

[Google Scholar](#)[ORCID](#)**Mahomoodally M. Fawzi**

Department of Health Sciences, Faculty of Science, University of Mauritius. Mauritius.

[ORCID](#)[ResearchGate](#)**María Inés Isla**

Facultad de Ciencias Naturales e Instituto Miguel Lillo. Universidad Nacional de
Tucumán, Tucumán, Argentina.

[ResearchGate](#)[ORCID](#)**Marisol Fernández Alfonso**

Facultad de Farmacia, Universidad Complutense de Madrid, España

**Mayank Gangwar**

Department of Pharmacology, Institute of Medical Sciences, Banaras Hindu University,
Varanasi (UP) India

[Scholar Google](#)[ORCID](#)**Silvia Debenedetti**

Pharmacy Career. Universidad de Belgrano, Buenos Aires, Argentina

[ORCID](#)[ResearchGate](#)**Vikash Kumar Ruhil**

PDM College of Pharmacy, Bahadurgarh, Haryana, India

[Google Scholar](#)**Yasser Shahzad**

Department of Pharmacy. COMSATS University Islamabad, Pakistan



JPPRes

Journal of Pharmacy
& Pharmacognosy
Research



The reviewers will be recruited among researchers and clinicians with high international reputation in Pharmacy and Pharmacognosy.

Last Update: August 20th, 2023



JPPRes

Journal of Pharmacy
& Pharmacognosy
Research

[Home](#) [About](#) [Archive](#) [Authors](#) [Reviewers](#) [Events](#) [Blog](#)

[Twitter](#) [LinkedIn](#) [Facebook](#)

© 2013-2020 by the authors; licensee JPPRes, Antofagasta, Chile. This journal is an open-access journal distributed under the terms and conditions of the Creative Commons Attribution license-Non Commercial 4.0 international. The content on this site is intended for health professionals. If you are not a health professional, please talk to your doctor about any doubts or concerns regarding your health

Made with ♥ by **AVAGAX Studio**



Fungistatic effect of *Moringa oleifera* Lam. on the metabolism changes of *Candida albicans*

[Efecto fungistático de *Moringa oleifera* Lam. sobre los cambios en el metabolismo de *Candida albicans*]

Basri A. Gani^{1*}, Cut Soraya², Vinna Kurniawati Sugiaman³, Fitri Yunita Batubara⁴, Dharli Syafriza⁵, Silvia Naliani⁶, Sri Rezeki⁷, Subhani Jakfar⁸, Muhammad Nazar⁹, Kemala Hayati¹⁰

¹Department of Oral Biology, Denstistry Faculty, Universitas Syiah Kuala. Jl., Kopelma Darussalam, Kecamatan Syiah Kuala, Kota Banda Aceh, Indonesia.

²Department of Conservative Dentistry, Dentistry Faculty, Universitas Syiah Kuala. Jl. Kopelma Darussalam, Kecamatan Syiah Kuala, Kota Banda Aceh, Indonesia.

³Department of Oral Biology, Dentistry Faculty, Universitas Kristen Maranatha. Jl. Prof. Drg. M.P.H. No. 65, Bandung, Jawa Barat, Indonesia.

⁴Department of Conservative Dentistry, Dentistry Faculty, Universitas Sumatera Utara. Jl. Alumni No.2, Kampus USU, Medan, Sumatera Utara, Indonesia.

⁵Department of Pedodontics, Dentistry Faculty, Universitas Syiah Kuala. Jl., Kopelma Darussalam, Kecamatan Syiah Kuala, Kota Banda Aceh, Indonesia.

⁶Department of Prosthodontics, Dentistry Faculty, Universitas Kristen Maranatha. Jl. Prof. Drg. Surya Sumantri, M.P.H. No. 65, Bandung, Jawa Barat, Indonesia.

⁷Department of Oral Medicine, Dentistry Faculty, Universitas Syiah Kuala. Jl., Kopelma Darussalam, Kecamatan Syiah Kuala, Kota Banda Aceh, Indonesia.

⁸Department of Dental Material, Denstistry Faculty, Universitas Syiah Kuala. Jl., Kopelma Darussalam, Kecamatan Syiah Kuala, Kota Banda Aceh, Indonesia.

⁹Department of Chemistry Education, Faculty of Teacher and Training, Universitas Syiah Kuala. Jl. Tgk. Hasan Krueng Kalee, Kopelma Darussalam, Kecamatan Syiah Kuala, Kota Banda Aceh, Aceh, Indonesia.

¹⁰Department of Dental Radiology, Dentistry Faculty, Universitas Syiah Kuala. Jl. Kopelma Darussalam, Kecamatan Syiah Kuala, Kota Banda Aceh, Aceh, Indonesia.

*E-mail: basriunoe@unsyiah.ac.id

Abstract

Context: *Candida albicans* is a pathological agent that triggers oral candidiasis because *C. albicans* can adapt to changes to increase growth and adhesion through biofilm formation mechanisms. *Moringa oleifera* Lam. has been reported to have fungistatic properties and increases the metabolism change of *C. albicans*.

Aims: To evaluate the fungistatic effect of *M. oleifera* leaves ethanolic extract (MOLE) on the metabolism changes of *C. albicans* cells associated with growth and biofilm formation.

Methods: The assessment of metabolism changes (stress response and metabolic alterations) of *C. albicans* by the action of MOLE was performed by mean of FTIR, growth assessment by spectrophotometry, biofilm formation with 1% crystal violet, also read by spectrophotometry, and observation of biofilm mass with a microscope.

Results: MOLE showed substantial absorption values based on topological polar surface area (<140 Å²). Concentrations of 25% and 6.25% of MOLE increased the stress response (metabolism changes) of *C. albicans* (66-75%), meanwhile 50% and nystatin (100.000 IU/mL) were similar in inducing metabolism changes of *C. albicans*. All concentrations of *M. oleifera* could inhibit the growth of *C. albicans* at all incubation times (24, 48, and 72 h) with an Optical Density (OD) of 0.02-0.05 (<300 CFU/mL) and were able to degrade the biofilm formation of *C. albicans* on a scale substantial at 24 and 48 h (OD>0.4), and moderate scale at 72 h (OD 0.2-0.39).

Conclusions: The extract of *M. oleifera* has increased metabolism changes (stress response) of *C. albicans* cells, which correlate with the ability to inhibit growth and biofilm formation for 24, 48, and 72 h.

Keywords: biofilm; *Candida albicans*; fungistatic; growth; stress response.

Resumen

Contexto: *Candida albicans* es un agente patológico que desencadena la candidiasis oral porque *C. albicans* puede adaptarse a los cambios para aumentar el crecimiento y la adhesión a través de mecanismos de formación de biopelículas. Se ha informado que *Moringa oleifera* Lam. tiene propiedades fungistáticas y aumenta el cambio de metabolismo de *C. albicans*.

Objetivos: Evaluar el efecto fungistático del extracto etanólico de hojas de *M. oleifera* (MOLE) sobre los cambios en el metabolismo de las células de *C. albicans* asociados con el crecimiento y la formación de biopelículas.

Métodos: La evaluación de los cambios en el metabolismo (respuesta al estrés y alteraciones metabólicas) de *C. albicans* por acción de MOLE fue realizada mediante FTIR, la evaluación del crecimiento mediante espectrofotometría, la formación de biofilm con cristal violeta al 1%, también leído mediante espectrofotometría, y la observación de la masa del biofilm con un microscopio.

Resultados: MOLE presentó valores de absorción sustanciales basados en el área de superficie polar topológica (<140 Å²). Las concentraciones de 25% y 6,25% de MOLE aumentaron la respuesta de estrés (cambios en el metabolismo) de *C. albicans* (66-75%), mientras que el 50% y la nistatina (100.000 UI/mL) fueron similares en la inducción de cambios en el metabolismo de *C. albicans*. Todas las concentraciones de *M. oleifera* pudieron inhibir el crecimiento de *C. albicans* en todos los tiempos de incubación (24, 48 y 72 h) con una densidad óptica (DO) de 0,02,0,05 (<300 ufc/mL) y fueron capaces de degradar la formación de biopelículas de *C. albicans* en una escala sustancial a las 24 y 48 h (DO>0,4), y moderada a las 72 h (DO 0,2-0,39).

Conclusiones: El extracto de *M. oleifera* ha aumentado los cambios en el metabolismo (respuesta al estrés) de las células de *C. albicans*, que se correlacionan con la capacidad de inhibir el crecimiento y la formación de biopelículas durante 24, 48 y 72 h.

Palabras Clave: biopelícula; *Candida albicans*; crecimiento; fungistático; respuesta al estrés.

AUTHOR INFO

ARTICLE INFO

Received: November 8, 2022.

Accepted: January 1, 2023.

Available Online: January 30, 2023.

ORCID:

[0000-0002-4592-398X](https://orcid.org/0000-0002-4592-398X) (BAG)

[0000-0002-8025-5534](https://orcid.org/0000-0002-8025-5534) (CS)

[0000-0002-3688-6718](https://orcid.org/0000-0002-3688-6718) (VKS)

[0000-0002-4159-9748](https://orcid.org/0000-0002-4159-9748) (FYB)

[0000-0001-5556-1671](https://orcid.org/0000-0001-5556-1671) (DS)

[0000-0002-5846-1200](https://orcid.org/0000-0002-5846-1200) (SN)

[0000-0002-0109-0701](https://orcid.org/0000-0002-0109-0701) (SR)

[0000-0002-2233-2239](https://orcid.org/0000-0002-2233-2239) (SJ)

[0000-0001-8361-606X](https://orcid.org/0000-0001-8361-606X) (MN)

[0000-0001-5219-6373](https://orcid.org/0000-0001-5219-6373) (KH)

INTRODUCTION

Candida albicans is a causative pathological agent triggering candidiasis infection. This infection can have implications for a decrease in the mucosal defense system. In addition, *C. albicans* has a faster spread if the biological conditions of the oral cavity are not balanced. Changes in temperature, salivary pH, and hormonal disturbances can trigger the development of the fungus, thereby exacerbating the infection (Jabra-Rizk et al., 2016).

Several factors can increase the risk of the growth of the fungus *C. albicans* in the oral cavity. One of the most common causes is overuse of antibiotics, comorbidities such as cancer, AIDS, and diabetes, and wearing dentures that cause chronic irritation. Candidiasis can be fatal if it reaches the bloodstream or vital organs such as the heart, but it is rare, except in chronic cases. Decreased immunity further exacerbates this infection (Singh et al., 2014).

Candida albicans have some virulence factors involved in the pathogenesis of infection. Hydrophilic and hydrophobic properties are two virulent factors that often work when applied to disease pathogenesis. In addition, adhesion factors and biofilm formation are reported to contribute together with hydrophilic properties when penetrating and infecting the host mucosa (Gani et al., 2017a). Increased biofilm formation helps *C. albicans* spread and form new communities to expand the infection. In addition, the intensity of the stress response and changes in metabolism are indicators of the virulence of *C. albicans* working in increasing disorder (Gulati and Nobile, 2016).

Previous studies of antifungal drugs prevent adhesion to the mucosal host by reducing biofilm formation. One side of the drug must have hydrophilic properties with the cell surface, and it must also have hydrophobic receptors to prevent interactions between pathogens, including *C. albicans*, and host cells, affecting the virulence properties of *C. albicans* cells (Parente-Rocha et al., 2017). In addition, the target antifungal drugs also aim to increase the stress response and changes in cell metabolism so that they can interfere with the cellular activity of *C. albicans* (Chen et al., 2020). Antioxidants sourced from medicinal plants are reported to have a high ability effect on causing damage to cell membranes and disrupting the cellular respiration system, thus disrupting the inter and intercellular communication system. This failure disrupts the mechanism of cell genetic information, which is the forerunner to a decrease in cell function, which ultimately causes stress and cell death. *Moringa oleifera* Lam. (family *Moringaceae*) has been reported to

have fungistatic properties through the membrane synthesis pathway (Unuofin and Lebelo, 2020).

Moringa species contain alkaloids, saponins, tannins, steroids, phenolic acids, glucosinolates, flavonoids, and terpenes. The genus' phytochemical variety makes it useful in pharmacology (Abd Rani et al., 2018). *Moringa oleifera* produces the antifungal protein known as Mo-CBP3, which has been explored as a possible candidate for eradicating fungal illnesses (Pinto et al., 2015). The chemical compounds hexadecanoic acid and oleic acid from *M. oleifera* were reported to have antifungal properties (Lin et al., 2016).

M. oleifera has been reported to be bacteriostatic and fungistatic. As a medicinal plant, this plant is efficacious in increasing immunity and preventing the development of pathogens (Qwele et al., 2013). Our previous study found that the ethanolic extract of *M. oleifera* had the highest cytotoxicity against *Streptococcus mutans* cells and could increase the hydrophilicity on the surface of *C. albicans* cells. This research is a series of studies focusing on the ability of *M. oleifera* leaves ethanolic extract on the stress response and alterations in *C. albicans* cell metabolism associated with growth and biofilm formation. The result of these studies is a reference for formulating fungistatic antifungal drugs from *M. oleifera* through cellular inhibitory pathways against *C. albicans* in the pathogenesis of oral candidiasis infections.

MATERIAL AND METHODS

Chemical and reagents

In this study were used nystatin from Novell Pharmaceutical Laboratories, Bogor, Indonesia, chemical and reagent products from Sigma-Aldrich, an affiliate of Merck KGaA, Darmstadt, Germany, among others, ethanol, alcohol, peptone, and Sabouraud dextrose agar, crystal violet 1%, sodium dodecyl sulfate, and phosphate-buffered saline.

This study used *M. oleifera* as the test material and *C. albicans* as the research subject. The interaction between *M. oleifera* and *C. albicans* was evaluated for stress response and metabolic changes confirmed by growth and biofilm formation on the 96-well plate. Assessment of the functional static properties of *M. oleifera* using concentrations of 6.25%, 12.5%, 25%, and 50%, and nystatin was used as a positive control, and incubation time of 24, 48, and 72 h was used as a determining variable to measure the period of action of *M. oleifera* causing decreased growth and biofilm formation of *C. albicans*.

Plant material

The *M. oleifera* was collected from the District of Aceh Besar, Province of Aceh, Indonesia (5.603444, 95.405863). A sample with voucher number Co2021 was deposited in the Laboratory of Oral Biology, Dentistry Faculty, Universitas Syiah Kuala, Darussalam, Banda Aceh, Indonesia. The plant material was extracted in the Chemical Laboratory, Faculty of Mathematics and Natural Science, Universitas Syiah Kuala, Darussalam Banda Aceh, Indonesia.

Extraction and GC-MS of *Moringa oleifera*

The *M. oleifera* leaves of 1 kg were washed and dried for two days till wilted. The *M. oleifera* leaves were blended into powder and stored airtight, then soaked in 100 mL of 96% ethanol in a clean flat-bottomed glass container. They changed the solvent for three days and remade residue and filtrate. The filtrate is concentrated with a rotary vacuum evaporator at 50°C and 75 mmHg to acquire the extract (Soraya et al., 2021). The GC-MS analysis of *M. oleifera* leaf ethanol extract was performed using a Shimadzu Japan QP2010PLUS with a polymethyl silicon-coated fused GC column (2010). Conditions: 80–200°C, 5 min, 200°C for 20 min. The flame ionization detector (FID) temperature was 300°C, the injection temperature was 220°C, and the nitrogen carrier gas was a flow rate of 1 mL/min. 116.9 kPa. 30 m column, 0.25 mm diameter, 50 mL/min flow rate (Yusuf et al., 2021). The chromatograms obtained from GC-MS should be displayed in Fig. 1.

Bioactivity assay compounds of *Moringa oleifera*

Utilizing the Molinspiration property engine software, the efficiency and molecular properties were calculated. Chemdraw professional v.16 Cambridge software was used to create the Compound structures, and Chem3D v.16 Cambridge software was used to convert the 2D designs into 3D. In addition, the software mol-inspiration properties engine v. 2018.10 was used to calculate the bioactivity and molecular scores of *M. oleifera* compounds. The structure of the two compounds is transferred to the mol-inspiration canvas in the first step. Clicking the computed properties button then calculates the number of atoms, molecular weight, particles, and essential groups such as N, O, OH, and NH. In the second phase, the effectiveness of the two test materials in response to the host or infectious agent was determined (Alberga et al., 2018).

Culture and growth assay

ATCC 10231 *Candida albicans* was recultured on Sabouraud Dextrose Agar medium for 48 h in an aerobic atmosphere at 37°C. Then, one colony was re-

cultured in Peptone medium for 48 h at 37°C. The McFarland 0.5 (1.5×10^8 CFU/mL) was then standardized. The amount of *M. oleifera* gel administered to *C. albicans* was then determined to observe its effect on the stress response and changes in cell metabolism, growth, and biofilm formation (Gani et al., 2017b).

In a 96-well plate, 50 μ L of a *C. albicans* suspension was put in each well (triple series). After that, it was allowed to sit at room temperature for 15 min before one hundred microliters of *M. oleifera* extract were added to each of the different concentrations. In addition, the *C. albicans* suspension was modified with the test material for 10 min at a speed of 200 rpm. After that, the cultures were grown in an aerobic environment at 37°C for 24, 48, and 72 h. Furthermore, the growth inhibition was assessed by spectrophotometry (xMark™ Microplate Absorbance Spectrophotometer, Biorad, California, USA) based on turbidity. Scale for rating growth based on optical density (OD). OD 0.05 nm (<150 CFU/mL), 0.08-0.1 nm (McFarlan 0.5; <300 CFU), OD 0.11-0.29 nm (McFarland 1; 300-600 CFU), OD 0.3-0.49 nm (McFarland 2; 600-1200 CFU). These scales were adopted by McFarland Standard for *in vitro* use only, catalog No. TM50-TM60, Scott Sutton, Measurement of Microbial Cells by Optical Density, 2011 (Sutton, 2011).

Biofilm assay

A total of 25 μ L of saliva was collected on a 96-well plate for 15 min. Then discarded and continued with the addition of 30 μ L *C. albicans* on the bottom, which had been covered with saliva on each well-plate and allowed to stand for 15 min. Then, *M. oleifera* extract gel was added to each well of 100 μ L and then adapted to room temperature for 10 min at 300 rpm. The interaction of *C. albicans* biofilm formation ability was assessed after incubation for 24 and 48 and 72 h at 37°C. Assessment of the inhibition of biofilm formation begins by removing the mixed solution of *C. albicans* and the test material. Then added, 1% sodium dodecyl sulfate (SDS) solution 100 μ L for 15 min and discarded and washed once with phosphate-buffered saline (PBS). Then, 150 μ L of glycerol was added to each well-plate and washed once with 200 μ L of PBS. Visualization of the biofilm mass by adding 150 μ L of 1% crystal violet for 10 min, then washing with 76% ethanol (150 μ L) and PBS once. Quantitative data reading with spectrophotometry at 520 nm (Asmah et al., 2022).

Preparation of whole-cell *Candida albicans*

The fungus *C. albicans*, equivalent to McFarlan 0.5 (<300 CFU/mL) taken 50 μ L, was included in the test material with various concentrations. Furthermore, adaptation was carried out at room temperature for

15 min at a speed of 200 ×g. The whole cell extraction process of *C. albicans* was adopted from He et al. (2009) research with several modifications. Homogenization of the fungus with the test material was checked based on changes in turbidity. Furthermore, the mixture of the two was incubated for 48 h in an aerobic atmosphere. The whole cell *C. albicans* extraction process was initiated by centrifugation for 5 min at 2000 rpm (separating *C. albicans* cells in the test material). Then the supernatant (*C. albicans* cells) was collected and incubated for 30 min at 37°C, vortexed again for 30 s, and 0.1 M HCl (200 mL in 1 mL) was added. Then it was vortexed for 30 s and incubated at 37°C for 5 min. Then it was centrifuged at 300 ×g for 10 min. The precipitate was collected, 500 mL PBS was added, vortexed for 30 s, and centrifuged 3000 ×g for 10 min. It was followed by washing the residue by adding 70% ethanol (1:3) and centrifuging 2000 ×g for 15 min. Furthermore, the deposit was collected as an extract whole (surface protein and endo-protein). Next, a 2 mL sample buffer solution was added. Then, the FTIR (IRPrestige 21, Merk Shimadzu, Kyoto, Japan) device examined the stress response and metabolic changes of *C. albicans* cells under *M. oleifera*.

Stress response and metabolism changes assay

Fourier transform infrared spectroscopy (FTIR) assessment of the stress response of *C. albicans* under the influence of saliva was analyzed based on the functional groups contained in the sample. This analysis uses FTIR-ATR with a wave number of 4000 cm⁻¹-400 cm⁻¹, producing transmittance spectra. In the first stage, an example of *C. albicans* is placed on the surface of a transparent Infrared (IR) prism with a refractive index that is always higher than the sample (the refractive index of bacterial cells is estimated at 1.39). The radiation beam is directed at one of the prism walls for the prism-sample interface at an angle higher than the barrier. Under these conditions, complete reflection occurs on the side of the internal prism. The reflected light exits through the walls of the second prism, where the light intensity and absorption spectrum are recorded (Alvarez-Ordóñez et al., 2011). Stress response assessment based on window peaks. The absorbance value of the intensity (wave) with the area value of each peak becomes the basis for measuring the stress response and changes in the metabolism of *C. albicans* cells. Stress response and metabolic changes are assessed based on changes in constituent cell elements such as lipids, proteins, carbohydrates, and nucleic acids.

The reading of the stress response under the influence of *M. oleifera* used the ATR-FTIR (cm⁻¹) mode approach. Assessment of bacterial stress response based on FTIR assessment adopted from Zarnowiec et

al. (2015) based on windows zone (W). W1- lipids (3,000-2,800 cm⁻¹), W2- proteins (1,700-1,500 cm⁻¹), W3-nucleic acids (1,500-1200 cm⁻¹), W4-carbohydrates (1200-900 cm⁻¹), W5 -fingerprints regions (900-500).

Statistical analysis

Data on growth and biofilm formation were analyzed by One Way ANOVA. At the same time, data on stress response and changes in metabolism were described descriptively based on the frequency value of each concentration of *M. oleifera*. The limit of significance (p<0.05).

RESULTS

Fig. 1 shows the chromatogram of *M. oleifera*. As much as there 17 chemical compounds were exposed based on the GC spectrum, and we reported 13 chemical compounds that were believed to have potential as antioxidants compounds and worked to prevent the development of fungi, including *C. albicans* (Senthilkumar et al., 2015). The peaks show up at 5.452 min of retention until 33.82 min. The GC analysis was in tandem with a mass spectrometer to obtain the relative mass of each compound. The data from the instrument was then compared with the NIST compound database. A total of 13 combinations from varied compound classes were detected and tabulated in Fig. 2. which reports the GC-MS results of several chemical compounds contained in the extract of *M. oleifera* and also reports the rate times and percentage of these chemical compounds of *M. oleifera*.

Fig. 3 shows that the stress indicator experienced by *C. albicans*, based on changes in absorbance (peak), occurred due to changes in nucleic acid synthesis, thus disrupting lipid synthesis, protein, and carbohydrates. At the same time, areal fingerprints can be ascertained that there is a change in virulence properties due to the stress response. The authors assume that the changes in the fingerprints introduced can indicate the assessment of growth and biofilm formation by *C. albicans*. Although these results are debatable because this study correlates with the inhibition of growth and formation of *C. albicans* biofilms, the fingerprints area can certainly be used as a reference for assessing the virulence properties of the pathogen.

Fig. 4 reports the results of the evaluation and analysis of FTIR data based on the intensity value (wavelength) given to *C. albicans* cells that have been affected by *M. oleifera* extract gel. At the same time, the area value is the change in the metabolic properties of *C. albicans* cells after interacting with *M. oleifera*. The intensity and area values in the FTIR assessment of biological products must be synergistic. The small-

er the wave given, the higher the energy produced to obtain the target value (changes in cell metabolism). This study showed a close relationship between the intensity and area values for *C. albicans* cells, which were affected by *M. oleifera* extract concentrations of 6.25%, and 25%. In contrast, 50% and the nystatin group had similarities in inducing changes in the metabolism of *C. albicans* cells.

In Fig. 5, it was reported that at all concentrations of *M. oleifera*, it caused a loss of *C. albicans* biofilm mass or caused a decrease in the morphology of the

biofilm mass. It can be assumed that *M. oleifera* has disrupted the metabolism of *C. albicans* cells to reduce biofilm formation. It is related to the stress response experienced by *C. albicans* during interaction with the test material (Table 1). Fig. 6 shows that *M. oleifera* caused changes in *C. albicans* to growth and biofilm formation at all incubation times. These findings indicate that the inhibition of biofilm formation affects the growth inhibition of *C. albicans*. It means that disrupting the formation of biofilms has interfered with the growth of *C. albicans*.




Figure 1. Gas Chromatography spectrum of *Moringa oleifera*.

Three significant components with a quantity greater than 5 percent are compounds (glycerol, 2,3-dihydro-3,5-dihydroxy-6-methyl-4H-pyran-4-one, and 1,3,4,5-tetrahydroxy-cyclohexane carboxylic acid.




Figure 2. Active compound structures of *M. oleifera* identified by GC-MS.




Figure 3. The FTIR was performed on the stress response of *C. albicans* after interacting with *M. oleifera* (4000-400 wavenumbers; cm^{-1}). In biological assessment, the representation of stress response values is obtained from the frequency range (wavenumbers/ cm^{-1}) based on window 1 (lipid), window 2 (protein), window 3 (nucleic acid), window 4 (carbohydrates), window 5 (fingerprints regions). Changes in absorbance of *C. albicans* cells affected by *M. oleifera* and nystatin from each window were used as a measurement value to analyze the response of *C. albicans* to the environment (stress response).




Figure 4. The FTIR graph pattern of metabolic changes of *C. albicans* after being induced with *M. oleifera* at a frequency range of 4000-500 cm^{-1} . The value of intensity (wavelength) and area (interaction) between *M. oleifera* and *C. albicans* became a reference for changes in *C. albicans* cell metabolism. Concentration groups of 25% and 6.25% had a better ability to interfere with the metabolism of *C. albicans*.

Table 2 shows the percentage quantity of each chemical compound contained in *M. oleifera*. 1,3,4,5-tetrahydroxy-cyclohexanecarboxylic acid compounds have the highest amount among other chemical compounds (43.04%). The assessment of drug properties using the Molinspiration property engine v2018.10 software. The topological polar surface area (TPSA), milogP, molecular weight, and working volume assess work as a drug ingredient. The TPSA value

shows that all TSPA values are below 140 \AA^2 , meaning that all these compounds have excellent absorption values. While the milogP value, all chemical compounds were below 4, except for Hexadecanoid compounds, which had a milogP of 7.06. In addition, all these compounds have a volume above $75 \text{ mm}^3/\text{mol}$. The volume quality of the compounds correlates with pharmacokinetic activity.





Table 1 shows that the ethanolic extract of *M. oleifera* at all concentrations can increase the stress response of *C. albicans*. At a concentration of 25%, it had a better ability to increase the stress response of *C. albicans* and was similar to the positive control group of nystatin (74%). Table 3 shows the incubation times of 24, 48, and 72 h showing the growth intensity of *C. albicans* with an average OD of 0.06-0.02 nm. This value equals <150 CFU/mL or McFarland 0.5 (1.5 ×

10⁸). Based on the One Way ANOVA analysis results, the concentration of *M. oleifera* affected the growth of *C. albicans* at incubation times of 24, 48, and 72 h. All concentrations showed similar properties in reducing the growth of *C. albicans* at OD values of 0.02-0.05, meaning that *C. albicans* colonies ranged below 300 CFU/mL. This ability indicates that *M. oleifera* is fungistatic.

Table 4 shows that *M. oleifera* extracts can reduce biofilm formation (biofilm static). In general, all concentrations of *M. oleifera* extract gel can reduce biofilm on a strong scale (OD > 0.4), except at 72 h, there is a moderate decrease. OD (0.2-0.39), except 25% concentration. Assessment of the biofilm inhibition scale using the reference of Syafriza et al. (2021). The anti-

biofilm assessment according to OD spectrophotometry, OD 0.4 (strong); OD=0.2-3.9 (moderate); OD=0.05-0.1 (low); OD < 0.05 (no biofilm formation). Fig. 4 reports that *M. oleifera* can degrade *C. albicans* biofilm at all concentrations with various frequencies. As a comparison, *C. albicans* has a substantial quantity of biofilm mass.

Table 1. The FTIR peak assessment of the stress response of *C. albicans* cells after interaction with *M. oleifera*.

<i>M. oleifera</i>	Frequency range (cm ⁻¹)	Bands	Absorbance (A)		Area (%)			Stress response	Comments	
			Value	Corr	Freq	Value	Corr			Freq
Conc _{50%}	3240.41-451.34	16	10.56	0.166	14%	795	6.46	21%	65%	Increase
Conc _{25%}	3240.41-462.92	17	9.39	0.197	13%	489	9.84	13%	75%	Increase
Conc _{12.5%}	3240.41-470.63	15	5.56	0.297	8%	1,028	10.25	27%	66%	Increase
Conc _{6.25%}	3240.41-470.63	15	14.59	0.156	20%	721	6.26	19%	62%	Increase
Nystatin	3240.41-451.34	20	11.84	0.103	16%	381	6.59	10%	74%	Increase
<i>C. albicans</i>	3240.41-462.92	17	21.95	0.143	30%	463	5.96	12%	58%	Standard

Table 2. The assessment of medicinal properties of chemical compounds of *M. oleifera*.

NO	Compounds of <i>M.oleifera</i>	GC-MS (Percent area)	Drugs properties assessment			
			milogP	TPSA(Å ²)	MW (kDa)	Vol (mm ³ /mol)
1	Alpa-butyrolactone	2.06	0.90	29.6	86.09	79.95
2	1,3-Cyclopentanedione	4.90	-0.68	34.145	98.1	90.17
3	Glycerol	8.48	-1.60	60.68	92.09	87.12
4	Piperidine-1,2,6-trimethyl-cis	1.33	1.67	12.03	113.2	131.38
5	1,2-Epoxy cyclohexane	2.34	1.55	12.53	98.14	100.8
6	Benzeneacetaldehyde	3.05	1.94	17.07	120.15	119.83
7	2-Pyrrolidinone	2.02	-0.18	29.1	85.11	83.58
8	2,3-Dihydro-3,5-dihydroxy-6-methyl-4H-pyran-4-one	8.00	-0.46	66.76	144.13	123.4
9	Benzeneacetonitrile,4-hydroxy-	4.87	1.42	44.02	133.15	125.72
10	1,2,3,4-Tetrahydro-cyclopentan[b]indole	1.11	2.47	12.03	159.23	158.82
11	1,3,4,5-Tetrahydroxy-cyclohexane carboxylic acid	43.04	-1.76	97.98	176.17	153.44
12	Hexadecanoic acid	3.04	7.06	37.3	256.43	291.42
13	n-cbz-beta-alanine	2.05	1.06	75.63	223.23	201.82

Table 3. The growth inhibition of *C. albicans* by *M. oleifera*.

<i>M. oleifera</i>	N	24 h				48 h				72 h			
		OD 620 nm	SD	Colony (CFU/mL)	Freq	OD 620 nm	SD	Colony (CFU/mL)	Freq	OD 620 nm	SD	Colony (CFU/mL)	Freq
Conc _{50%}	3	0.04	0.002	<150	12%	0.05	0.002	<150	11%	0.06	0.009	<200	13%
Conc _{25%}	3	0.02	0.003	<100	6%	0.05	0.003	<150	11%	0.04	0.002	<150	10%
Conc _{12.5%}	3	0.03	0.003	<150	9%	0.04	0.003	<150	11%	0.0	0.005	<150	10%
Conc _{6.25%}	3	0.03	0.004	<150	9%	0.06	0.004	<200	13%	0.05	0.004	<150	10%
C+ (nystatin)	3	0.05	0.016	<300	15%	0.09	0.003	<150	19%	0.10	0.007	≤300	21%
<i>C. albicans</i>	3	0.16	0.01	300-600	48%	0.17	0.01	300-600	36%	0.17	0.02	300-600	35%
*p-value		0.025				0.0421				0.031			

*One Way ANOVA; SD: Standard Deviation; Freq: Frequency.

Table 4. Inhibition of biofilm formation of *C. albicans* by *M. oleifera*.

<i>M. oleifera</i>	N	24 h			48 h			72 h		
		OD (520 nm)	SD	Freq	OD (520 nm)	SD	Freq	OD (520 nm)	SD	Freq
Conc _{50%}	3	0.412	0.011	18%	0.512	0.003	20%	0.311	0.019	19%
Conc _{25%}	3	0.501	0.002	22%	0.523	0.003	21%	0.422	0.022	25%
Conc _{12.5%}	3	0.411	0.005	18%	0.411	0.002	16%	0.302	0.015	18%
Conc _{6.25%}	3	0.403	0.024	18%	0.411	0.005	16%	0.315	0.014	19%
Nystatin	3	0.450	0.012	20%	0.691	0.002	27%	0.212	0.027	19%
*p-value		0.062			0.041			0.050		

*One Way ANOVA; SD: Standard Deviation; Freq: Frequency.

DISCUSSION

This study demonstrates and evaluates the potential for the fungistatic capabilities of *M. oleifera* to influence the stress response and metabolic alterations in *C. albicans* cells. Changes in the development and ability of *C. albicans* to build biofilms verified this effect. Figs. 1 and 2 depict the GC spectra of *M. oleifera* with the number of chemical components as a reference for evaluating its fungus function. Glycerol, 2,3-dihydro-3,5-dihydroxy-6-methyl-4H-pyran-4-one, and 1,3,4,5-tetrahydroxy-cyclohexanecarboxylic acid are three chemical compounds derived from *M. oleifera* reported having antifungal potential (Chakraborty et al., 2021; Senthilkumar et al., 2015). This glycerol molecule increases the permeability of fungal cells and inhibits the chlamydospore system as a toxin against host cells (Chen et al., 2018). This compound is reported as an antioxidant that influences the development of bacteria and fungi (Ruta and Farcasanu, 2021).

M. oleifera can increase the stress response of *C. albicans* (Table 1). It is indicated by changes in *C. albicans* cell metabolism (Fig. 3), which has implications for decreased growth and increased degradation of biofilm mass (Fig. 5 and Table 4). Changes in *C. albicans* cell metabolism after exposure to *M. oleifera* (antifungal) cause disruption of the protein synthesis system and response to virulence proteins against antifungals (da Silva Neto et al., 2020). This failure causes a decrease in adaptation to extreme environments so that cells undergo biological and physical changes. Biologically, the cell wall changes, and a leak occurs, allowing antifungal compounds to enter the cell membrane (Araújo et al., 2019). Physically, the antifungal effect of *M. oleifera* causes an excessive change in the hydrophobic pressure response of the membrane, causing *C. albicans* cells to have difficulty growing and forming quorum sensing as a prerequisite for infection (Araújo et al., 2019).

The ability of *M. oleifera* to cause an increase in stress response and changes in the metabolism of *C. albicans* is related to the cellular nature of the action of the chemical compound *M. oleifera* in increasing its effect on *C. albicans* cells. The analysis of drug properties of *M. oleifera* using the Molinspiration software (Table 2) shows that *M. oleifera* has a topological polar surface area (TPSA) value below 140 Å², meaning that all these compounds have excellent absorption values (Prasanna and Doerksen, 2009). The absorbance of these compounds is related to the ability of the active compound to penetrate the damaged target or the pharmacokinetic target of the drug. So it can be assumed that all these compounds have a perfect effect on reducing growth and inhibiting the virulence of *C. albicans* biofilm formation. The TPSA assessment relates a molecule of the active compound as the sum of the surfaces on all polar atoms or molecules, primarily oxygen and nitrogen, including the hydrogen atoms attached to the compounds (Lim et al., 2018).

The chemical compound of *M. oleifera* evaluated a milogP value, below 4, except for the hexadecanoid compounds, which had a milogP of 7.06. The LogP value of the chemical compound *M. oleifera* is considered to be active. Suppose the LogP is more significant than one or less than 4 in this value range. In that case, the *M. oleifera* compound is considered to have optimal physicochemical properties for oral drugs (Van De Waterbeemd and Gifford, 2003). Czyski (2019) reported that the compound's Log P value indicates the drug's permeability to reach the target cells and tissues in the body. This ability is related to the volume of each *M. oleifera* compound, which has a volume above 75 mm³/mol, meaning that the total volume of the active compound can be in line with the pharmacodynamic and pharmacokinetic properties of the drug.

The *M. oleifera* has chemical compounds with varying molecular weights 80-256 kDa (Table 2). High molecular weight can be considered to increase chemical resistance - up to a certain point and be an indica-

tor of the viscosity properties of the drug (Broseta et al., 1990). The increase in the viscosity of the *M. oleifera* compound can complicate the process of destroying the cell structure of *C. albicans*. Chae et al. (2005) reported that molecular weight could be considered absorption in the intestine. Using lower molecular weight standards of chemical compounds provides good fungistatic properties because the absorption of chemical compounds co-occurs with a decrease in infectious agents because the function of active compounds guarantees or controls changes in the virulence properties of pathogenic cells (Bickerton et al., 2012).

Cellularly, it is inevitable that the compounds contained in *M. oleifera* have the potential to interfere with changes in cell components such as nucleic acids, lipids, proteins, and carbohydrates. This element is reported as a building block for the integrity of a cell (Erca et al., 2021). Changes in these four chemical elements can be used as a basis for referring to the stress response of cells (Fig. 3). These findings show that the cellular activity of *C. albicans* tremendous changes after interacting with *M. oleifera* is a nucleic acid, lipid, protein, and carbohydrate. Fulda et al. (2010) reported that changes always influence environmental cell activity. The response to these changes manifests in cell defense by expressing several proteins, lipids, and carbohydrates. This increase in activity can be used as a response to environmental changes (stress response) to adjust to the environment (LaDage, 2015).

Alterations in the stress response are linked to metabolic alterations, which can affect cell proliferation and the growth and biofilm formation of *C. albicans* (Li et al., 2015). The data in Table 3 shows a decrease in the growing intensity of *C. albicans* after interaction with *M. oleifera*, which was related to the inhibition of biofilm formation by *C. albicans* (Table 4). This decrease is related to the role of active compounds that interfere with the protein synthesis of *C. albicans*. As a result, it causes a change in the response of the cell membrane to the surface environment, causing leakage of the membrane and releasing intracellular fluid so that the cell becomes lysed and undergoes premature death before dividing (Rock and Kono, 2008).

The decrease in biofilm formation is related to the active compound *M. oleifera* increasing the pressure on the membrane permeability of pathogens, thereby disrupting the hydrophobicity of the membrane to form biofilms (Soraya et al., 2022). Ponde et al. (2021) reported that the hydrophobicity of the cell surface of *C. albicans* can be a determinant of the formation of quorum sensing on the mucosal surface and that the hydrophobicity is a determinant of the development

of biofilms on the mucous membrane. In addition, *M. oleifera* could degrade and reduce the morphology of the biofilm mass (Fig. 4). Mishra et al. (2020) reported that phytochemical compounds of herbal plants play an essential role in inhibiting bacterial adhesion and suppressing genes associated with biofilm formation. According to Nobile et al. (2008), this phenomenon reported that many antifungal drug compounds work to describe the structure of compounds (proteins) that make up biofilms, such as HWP1 or ALS1-3 proteins. The decomposition of biofilm protein compounds can cause a decrease in the biofilm matrix, causing mass degradation and even lowering the tension of spreading biofilms by disrupting communication between pathogenic cells during quorum sensing (Tits et al., 2020).

M. oleifera was responsible for the decline in biofilm production, which corresponded with the organism's slower rate of growth (Fig. 6). It indicates that the height of the biofilm generated has a significant role in determining the development of *C. albicans*, which *M. oleifera* influences. The fungistatic effect of *M. oleifera* and reducing the virulence of *C. albicans* also interfere with protein synthesis. It can limit adaptability to the antifungal environment, disrupting growth and interaction with the surrounding environment (Santos et al., 2021). Meanwhile, *C. albicans* cells cannot increase or make morphological changes at this stage. The antifungal activity of *M. oleifera* can cause *C. albicans* cells to die prematurely. This potential can serve as a reference for using active material to limit the infection caused by *C. albicans*, which plays a role in the pathogenesis of oral candidiasis.

CONCLUSION

The extract of *M. oleifera* has increased the metabolism changes (stress response) of *C. albicans* cells, which have implications for decreasing growth and inhibiting biofilm formation for 24, 48, and 72 h.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

ACKNOWLEDGMENTS

The Universitas Syiah Kuala, Ministry of Education, Culture, Research, and Technology, granted the research. Number: 146/UN11/SPK/PNBP/2022 dated 11 February 2022.

REFERENCES

- Abd Rani NZ, Husain K, Kumolosasi E (2018) *Moringa* genus: A review of phytochemistry and pharmacology. *Front Pharmacol* 9: 108. <https://doi.org/10.3389/fphar.2018.00108>
- Alberga D, Trisciuzzi D, Montaruli M, Leonetti F, Mangiatordi GF, Nicolotti O (2018) A new approach for drug target and

- bioactivity prediction: The multifingerprint similarity search algorithm (mussel). *J Chem Inf Model* 59: 586–596. <https://doi.org/10.1021/acs.jcim.8b00698>
- Alvarez-Ordóñez A, Mouwen D, Lopez M, Prieto M (2011) Fourier transform infrared spectroscopy as a tool to characterize molecular composition and stress response in foodborne pathogenic bacteria. *J Microbiol Methods* 84: 369–378. <https://doi.org/10.1016/j.mimet.2011.01.009>
- Araújo NM, Dias LP, Costa HP, Sousa DO, Vasconcelos IM, de Morais GA, Oliveira JT (2019) CITI, a Kunitz trypsin inhibitor purified from *Cassia leiandra* Benth. seeds, exerts a candidicidal effect on *Candida albicans* by inducing oxidative stress and necrosis. *Biochim Biophys Acta Biomembr* 1861: 183032. <https://doi.org/10.1016/j.bbmem.2019.183032>
- Asmah N, Suniarti DF, Bachtiar EW, Margono DA, Gani BA (2022) Chemical compounds anti-bacterial of *Citrus aurantifolia* ethanol extract to inhibit the early biofilm formation and growth of *Enterococcus faecalis* root canal isolate. *Res J Pharm Technol* 15: 2667–2674. <https://doi.org/10.52711/0974-360X.2022.00446>
- Bickerton GR, Paolini GV, Besnard J, Muresan S, Hopkins AL (2012) Quantifying the chemical beauty of drugs. *Nat Chem* 4: 90–98. <https://doi.org/10.1038/nchem.1243>
- Broseta D, Fredrickson GH, Helfand E, Leibler L (1990) Molecular weight and polydispersity effects at polymer-polymer interfaces. *Macromolecules* 23: 132–139. <https://doi.org/10.1021/ma00203a023>
- Chae SY, Jang MK, Nah JW (2005) Influence of molecular weight on oral absorption of water soluble chitosans. *J Control Release* 102: 383–394. <https://doi.org/10.1016/j.jconrel.2004.10.012>
- Chakraborty S, Majumder S, Ghosh A, Saha S, Bhattacharya M (2021) Metabolomics of potential candidates conferring antioxidant property to varied polar and non-polar solvent extracts of *Edgaria darjeelingensis* CB Clarke. *Bull Natl Res Cent* 45: 48. <https://doi.org/10.1186/s42269-021-00503-3>
- Chen X, Zhang Z, Chen Z, Li Y, Su S, Sun S (2020) Potential antifungal targets based on glucose metabolism pathways of *Candida albicans*. *Front Microbiol* 11: 296. <https://doi.org/10.3389/fmicb.2020.00296>
- Chen Y, Mallick J, Maqnas A, Sun Y, Choudhury BI, Côte P, Yan L, Ni TJ, Li Y, Zhang D, Rodríguez-Ortiz R (2018) Chemogenomic profiling of the fungal pathogen *Candida albicans*. *Antimicrob Agents Chemother* 62: e02365-17. <https://doi.org/10.1128/AAC.02365-17>
- Czyrski A (2019) Determination of the lipophilicity of ibuprofen, naproxen, ketoprofen, and flurbiprofen with thin-layer chromatography. *J Chem* 2019: 3407091. <https://doi.org/10.1155/2019/3407091>
- da Silva Neto JX, da Costa HPS, Vasconcelos IM, Pereira ML, Oliveira JTA, Lopes TDP, Dias LP, Araújo NMS, Moura LFWG, Van Tilburg MF (2020) Role of membrane sterol and redox system in the anti-candida activity reported for Mo-CBP2, a protein from *Moringa oleifera* seeds. *Int J Biol Macromol* 143: 814–824. <https://doi.org/10.1016/j.ijbiomac.2019.09.142>
- Ercan K, Geceşefa OF, Taysi ME, Ali Ali OA, Taysi S (2021) *Moringa oleifera*: A review of its occurrence, pharmacological importance and oxidative stress. *Mini Rev Med Chem* 21: 380–396. <https://doi.org/10.2174/1389557520999200728162453>
- Fulda S, Gorman AM, Hori O, Samali A (2010) Cellular stress responses: Cell survival and cell death. *Int J Cell Biol* 2010: 214074. <https://doi.org/10.1155/2010/214074>
- Gani BA, Alghassani AQ, Mubarak Z, Bachtiar EW, Bachtiar BM (2017a) Potential of cigarette smoke condensate to increase biofilm formation of *Candida albicans* isolate ATCC 10261. *J Syiah Kuala Dent Soc* 2: 33–39.
- Gani BA, Bachtiar EW, Bachtiar BM (2017b) The role of cigarettes smoke condensate in enhanced *Candida albicans* virulence of salivary isolates based on time and temperature. *J Int Dent Med Res* 10: 769–777.
- Gulati M, Nobile CJ (2016) *Candida albicans* biofilms: Development, regulation, and molecular mechanisms. *Microbes Infect* 18: 310–321. <https://doi.org/10.1016/j.micinf.2016.01.002>
- He Y-H, Tian X-B, Wan H-C, Wen Y-L, Zhang F-F, Ma Q-R (2009) Study on protein extraction methods for *Streptococcus mutans*. *Hua Xi Kou Qiang Yi Xue Za Zhi* 27: 100–103.
- Jabra-Rizk MA, Kong EF, Tsui C, Nguyen MH, Clancy CJ, Fidel PL, Noverr M (2016) *Candida albicans* pathogenesis: Fitting within the host-microbe damage response framework. *Infect Immun* 84: 2724–2739. <https://doi.org/10.1128/iai.00469-16>
- LaDage LD (2015) Environmental change, the stress response, and neurogenesis. *Integr Comp Biol* 55: 372–383. <https://doi.org/10.1093/icb/icv040>
- Li P, Seneviratne CJ, Alpi E, Vizcaino JA, Jin L (2015) Delicate metabolic control and coordinated stress response critically determine antifungal tolerance of *Candida albicans* biofilm persisters. *Antimicrob Agents Chemother* 59:6101–6112. <https://doi.org/10.1128/aac.00543-15>
- Lim J, Ryu S, Kim JW, Kim WY (2018) Molecular generative model based on conditional variational autoencoder for de novo molecular design. *J Cheminform* 10: 31. <https://doi.org/10.1186/s13321-018-0286-7>
- Lin HC, Kuo Y-L, Lee W-J, Yap H-Y, Wang S-H (2016) Antidermatophytic activity of ethanolic extract from *Croton tiglium*. *BioMed Res Int* 2016: 3237586. <https://doi.org/10.1155/2016/3237586>
- Mishra R, Panda AK, De Mandal S, Shakeel M, Bisht SS, Khan J (2020) Natural anti-biofilm agents: Strategies to control biofilm-forming pathogens. *Front Microbiol* 11: 566325. <https://doi.org/10.3389/fmicb.2020.566325>
- Nobile CJ, Schneider HA, Nett JE, Sheppard DC, Filler SG, Andes DR, Mitchell AP (2008) Complementary adhesion function in *C. albicans* biofilm formation. *Curr Biol* 18: 1017–1024. <https://doi.org/10.1016/j.cub.2008.06.034>
- Parente-Rocha JA, Bailão AM, Amaral AC, Taborda CP, Pácez JD, Borges CL, Pereira M (2017) Antifungal resistance, metabolic routes as drug targets, and new antifungal agents: an overview about endemic dimorphic fungi. *Med Inflamm* 2017: 9870679. <https://doi.org/10.1155/2017/9870679>
- Pinto CE, Farias DF, Carvalho AF, Oliveira JT, Pereira ML, Grangeiro TB, Freire JE, Viana DA, Vasconcelos IM (2015) Food safety assessment of an antifungal protein from *Moringa oleifera* seeds in an agricultural biotechnology perspective. *Food Chem Toxicol* 83: 1–9. <https://doi.org/10.1016/j.fct.2015.05.012>
- Ponde NO, Lortal L, Ramage G, Naglik JR, Richardson JP (2021) *Candida albicans* biofilms and polymicrobial interactions. *Crit Rev Microbiol* 47: 91–111. <https://doi.org/10.1080/1040841X.2020.1843400>
- Prasanna S, Doerksen R (2009) Topological polar surface area: A useful descriptor in 2D-QSAR. *Curr Med Chem* 16: 21–41. <https://doi.org/10.2174/092986709787002817>
- Qwele K, Hugo A, Oyedemi S, Moyo B, Masika P, Muchenje V (2013) Chemical composition, fatty acid content and antioxidant potential of meat from goats supplemented with moringa (*Moringa oleifera*) leaves, sunflower cake and grass hay. *Meat Science* 93: 455–462. <https://doi.org/10.1016/j.meatsci.2012.11.009>
- Rock KL, Kono H (2008) The inflammatory response to cell death. *Annu Rev Pathol* 3: 99–126. <https://doi.org/10.1146/annurev.pathmechdis.3.121806.151456>

- Ruta LL and Farcasanu IC (2021) Coffee and yeasts: From flavor to biotechnology. *Fermentation* 7: 9. <https://doi.org/10.3390/fermentation7010009>
- Santos LMM, Silva PM, Moura MC, Carvalho Junior AR, Amorim PK, Procópio TF, Coelho L, Silva LCN, Paiva PMG, Santos NDL and Napoleão TH (2021) Anti-Candida activity of the water-soluble lectin from *Moringa oleifera* seeds (WSMoL). *J Mycol Med* 31: 101074. <https://doi.org/10.1016/j.mycmed.2020.101074>
- Senthilkumar S, Sivakumar T, Arulmozhi K and Mythili N (2015) Gas chromatography-mass spectroscopy evaluation of bioactive phytochemicals of commercial green teas (*Camellia sinensis*) of India. *Asian J Pharm Clin Res* 8: 278–282.
- Singh A, Verma R, Murari A, Agrawal A (2014) Oral candidiasis: An overview. *J Oral Maxillofac Pathol* 18: 81–85. <https://doi.org/10.4103/0973-029X.141325>
- Soraya C, Alibasyah ZM, Gani, BA (2021) Biomass index and viscosity values of *Moringa oleifera* influenced by *Enterococcus faecalis*. *J Syiah Kuala Dent Soc* 6: 1–5. <https://doi.org/10.24815/jds.v6i1.21885>
- Soraya C, Syafriza D, Gani BA (2022) Antibacterial effect of *Moringa oleifera* gel to prevent the growth, biofilm formation, and cytotoxicity of *Streptococcus mutans*. *J Int Dent Med Res* 15: 1053–1061.
- Sutton S (2011) Measurement of microbial cells by optical density. *J Valid Technol* 17: 46–49.
- Syafriza D, Sutadi H, Primasari A and Siregar Y (2021) Spectrophotometric analysis of *Streptococcus mutans* growth and biofilm formation in saliva and histatin-5 relate to pH and viscosity. *Pesqui Bras Odontopediatria Clin Integr* 21: e0018. <https://doi.org/10.1590/pboci.2021.004>
- Tits J, Cammue BPA, Thevissen K (2020) Combination therapy to treat fungal biofilm-based infections. *Int J Mol Sci* 21: 8873. <https://doi.org/10.3390/ijms21228873>
- Unuofin JO, Lebelo SL (2020) Antioxidant effects and mechanisms of medicinal plants and their bioactive compounds for the prevention and treatment of type 2 diabetes: An updated review. *Oxid Med Cell Longev* 2020: 1356893. <https://doi.org/10.1155/2020/1356893>
- Van De Waterbeemd H, Gifford E (2003) ADMET *in silico* modelling: towards prediction paradise? *Nat Rev Drug Discov* 2: 192–204. <https://doi.org/10.1038/nrd1032>
- Yusuf H, Husna F, Gani BA (2021) The chemical composition of the ethanolic extract from *Chromolaena odorata* leaves correlates with the cytotoxicity exhibited against colorectal and breast cancer cell lines. *J Pharm Pharmacogn Res* 9: 344–356. https://doi.org/10.56499/jppres20.969_9.3.344
- Zarnowicz P, Lechowicz L, Czerwonka G, Kaca W (2015) Fourier transform infrared spectroscopy (FTIR) as a tool for the identification and differentiation of pathogenic bacteria. *Curr Med Chem* 22: 1710–1718. <https://doi.org/10.2174/0929867322666150311152800>

AUTHOR CONTRIBUTION:

Contribution	Gani BA	Soraya C	Sugijaman VK	Batubara FY	Syafriza D	Naliani S	Rezeki S	Jakfar S	Nazar M	Hayati K
Concepts or ideas	x									
Design	x	x		x	x		x	x	x	
Definition of intellectual content	x	x	x	x			x	x	x	
Literature search	x	x		x	x		x	x	x	x
Experimental studies	x				x					
Data acquisition	x		x	x	x	x		x	x	x
Data analysis	x									
Statistical analysis	x									
Manuscript preparation	x			x	x	x				
Manuscript editing	x			x	x	x		x	x	x
Manuscript review	x	x	x	x	x	x	x	x		x

Citation Format: Gani BA, Soraya C, Sugijaman VK, Batubara FY, Syafriza D, Naliani S, Rezeki S, Jakfar S, Nazar M, Hayati K (2023) Fungistatic effect of *Moringa oleifera* Lam. on the metabolism changes of *Candida albicans*. *J Pharm Pharmacogn Res* 11(1): 179–190. https://doi.org/10.56499/jppres22.1533_11.1.179

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Open Access: This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, duplication, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.