



SURAT TUGAS
No. 819A/FKG-UKM/X/2022

Yang bertanda tangan di bawah ini

Nama : Dr. Ignatius Setiawan, drg., MM.
Jabatan : Dekan Fakultas Kedokteran Gigi Universitas Kristen Maranatha

Dengan ini menugaskan kepada :

No	Nama Dosen	NIK
1.	Dr. Vinna Kurniawati Sugiawan, drg., M.Kes., PBO., CMC	120005
2.	Riani Setiadhi, drg., Sp.PM (K)	120058

Melaksanakan publikasi jurnal : *Antibacterial Potential Ethanol Extract of Beluntas Leaves (Pluchea indica L) to Streptococcus Sanguinis* di Journal of Syiah Kuala Dentistry Society pada Tahun 2022.

Demikian agar tugas ini dilaksanakan dengan sebaik-baiknya.

Bandung, 11 Oktober 2022
Dekan Fakultas Kedokteran Gigi
Universitas Kristen Maranatha

Dr. Ignatius Setiawan, drg., MM..
NIK. 120010

FAKULTAS
KEDOKTERAN GIGI



E-ISSN: 22502-0412

JDS

**Journal of Syiah Kuala
Dentistry Society**



**PUBLISHER
FKG USK PRESS**



PUBLISHER
FKGS USK PRESS



E-ISSN: 22502-0412

JDS (Journal of Syiah Kuala Dentistry Society) Phone: +62 651-7555183 Email: jds.fkg@usk.ac.id

[HOME](#) [ABOUT](#) [LOGIN](#) [REGISTER](#) [CATEGORIES](#) [SEARCH](#) [CURRENT](#) [ARCHIVES](#) [ANNOUNCEMENTS](#) [REVIEWER](#) [EDITORS](#) [ONLINE FIRST](#)

Home > Archives > **Vol 7, No 2 (2022)**

Vol 7, No 2 (2022)

December 2022

Table of Contents

Articles

Inhibitory test of turmeric leaves extract (*Curcuma Longa. L*) against the growth of *Streptococcus mutans* bacterial growth in vitro 73-77

Gostry Aldica DOHUDE, Hendry RUSDY, Olivia Avriyanti HANAFIAH, Preetypal Kaur JOHAL
10.24815/jds.v7i2.30229

Antibacterial potential ethanol extract of beluntas leaves (*Pluchea indica L*) to *Streptococcus sanguinis* 78-85

Wilma ADIWIJAYA, Riani SETIADHI, Vinna Kurniawati SUGIAMAN
10.24815/jds.v7i2.30230

Effectiveness of ethanol extract of kirinyuh leaf (*Chromolaena odorata L*) on the increase the fibroblasts and angiogenesis in nose mucosa of rabbit (*Iepus curpaneums*) 86-92

Teuku Husni TR, Novina RAHMAWATI, Irnawati IRNAWATI
10.24815/jds.v7i2.30231

Root morphology analysis of posterior teeth using intraoral periapical radiograph 93-97

Fitri Yunita BATUBARA, Dewi KARTIKA
10.24815/jds.v7i2.30232

The effect of ethanol extract of kirinyuh leaf (*Chromolaena odorata L*) on the allergy healing of rabbit (*Iepus curpaneums*) 98-106

Teuku Husni TR, Ikbal ISMAIL, Keishya MAURIEZA
10.24815/jds.v7i2.30233

Potential of chitosan oligosaccharide gel as a cavity cleanser against adhesive restoration adhesive on the cavity wall 107-112

Madya Theresa TARIGAN, Trimurni ABIDIN
10.24815/jds.v7i2.30234

Level of Knowledge on Radiation Protection in Roentgen Photo-taking among Clinical Dental Students 113-117

Kemala HAYATI, Herwanda HERWANDA, Yusmar YUSMAR
10.24815/jds.v7i2.30235

Association between the number of bilateral free-end posterior tooth loss and mastication performance in RSGM USU patients 118-124

Susanna Kristina Yanti SILALAH, Ismet Danial NASUTION
10.24815/jds.v7i2.30236

The relationship between static and dynamic occlusion based on the relationship between anterior and posterior teeth and the occlusion scheme in dentistry students 125-131

Riri Harliani SIHOTANG, Ariyani ARIYANI
10.24815/jds.v7i2.30237

Comparative Study of Micro-Leakage Between Glass Ionomer Cement Restoration Materials and Alkasises on Cavitas Class I (GV. Black) 132-137

Viona DIANSARI, Diana Setya NINGSIH, Sri FITRIYANI, Iin SUNDARI, Subhaini SUBHAINI,
M. Nur Hafidul Ghani GIBRAN

[FOCUS AND SCOPE](#)

[AUTHOR GUIDELINES](#)

[PUBLICATION ETHICS](#)

[OPEN ACCESS POLICY](#)

[PEER REVIEW PROCESS](#)

[ONLINE SUBMISSION](#)

[COPYRIGHT NOTICE](#)

[CONTACT US](#)

[JOIN US REVIEWER](#)


Citedness in
Scopus®



CERTIFICATE SINTA 5



TEMPLATE & STATEMENT



PLAGIARISM TOOL



REFERENCES TOOL



USER

Username

Password

Remember me

JOURNAL CONTENT

Bacterial profile in children with early childhood caries

138-149

Sinta Dwi ARDIYANTINI, Hening Tjaturina PRAMESTI, Meirina GARTIKA
10.24815/jds.v7i2.30239

The Streptococcus mutans ability to survive in biofilms and during dental caries formation: scoping review

150-158

Anisauma UTAMANINGYAS, Hening Tjaturina PRAMESTI, Felisha Febriane BALAFIF
10.24815/jds.v7i2.30295

Thermographic study of the maxillofacial area: the possibilities and prospects in modern dentistry

159-165

Abdillah Imron NASUTION, Mikhail Nikolaevich PANKOV
10.24815/jds.v7i2.30446

Indexed by:

Tools:

All papers published in FKG USK Press are licensed under a **Creative Commons Attribution-ShareAlike 4.0 International License**.

Social Media: Facebook Twitter Instagram YouTube

Search

Search Scope
All

- Browse
- By Issue
 - By Author
 - By Title
 - Other Journals
 - Categories

INFORMATION

- For Readers
- For Authors
- For Librarians

VISITORS

Visitors See m

73,188	95	37
5,591	78	36
697	68	33
576	68	32
296	50	30
279	49	30
198	46	29
130	46	28
113	46	22
113	45	21
103	42	21
99	40	15
98	39	
95	38	

FLAG COU

Hirtab.com

Vis. today 14
Visits 4 524
Pag. today 41

CURRENT ISSUE

1.0
 2.0
 1.0

JDS Journal of Syiah Kuala Dentistry Society

PUBLISHER
FKG USK PRESS

Crossref SINTA

E-ISSN: 22502-0412

JDS (Journal of Syiah Kuala Dentistry Society) Phone: +62 651-7555183 Email: jds.fkg@usk.ac.id

HOME ABOUT LOGIN REGISTER CATEGORIES SEARCH CURRENT ARCHIVES ANNOUNCEMENTS REVIEWER EDITORS ONLINE FIRST

Home > **Editorial Team**

Editorial Team

Editor in Chief

Assoc Prof. Basri A. Gani



Department of Oral Biology, Dentistry Faculty, Universitas Syiah Kuala, Darussalam, Banda Aceh, Aceh Indonesia

Managing Editor

Assoc Prof. Subhaini Jakfar



Departement of Dental Material, Dentistry Faculty, Universitas Syiah Kuala, Darussalam, Banda Aceh, Aceh Indonesia

International Editorial Board

Prof. Boy M. Bachtiar



Oral Biology, Dentistry, University of Indonesia, Indonesia

Prof. Talat Al-Gunaid



Orthodontic, Dentistry, University of Taibah, Saudi Arabia

Assoc Prof. Tzu Chieh Lin



Biomedical Engineering, National Taiwan University, Taiwan

Assoc Prof. Shahida Mohd-Said



Periodontics, Dentistry, University of Kebangsaan Malaysia, Malaysia

Assoc Prof. Nirun Intarut



Epidemiology, University of Mahasarakham, Thailand

Editors

Prof. Zaki Mubarak



Microbiology, Dentistry, University of Syiah Kuala, Indonesia

FOCUS AND SCOPE

AUTHOR GUIDELINES

PUBLICATION ETHICS

OPEN ACCESS POLICY

PEER REVIEW PROCESS

ONLINE SUBMISSION

COPYRIGHT NOTICE

CONTACT US

JOIN US REVIEWER

Citedness in
Scopus®



CERTIFICATE SINTA 5




TEMPLATE & STATEMENT



PLAGIARISM TOOL



REFERENCES TOOL



USER

Username

Password

Remember me

Prof. Anton Rahardjo



Public Health in Dentistry, University of Indoensia, Indonesia

English Language Advisor

Kismullah Abdul Muthalib, Ph.D



Departemen English Language, Faculty of Teacher Training and Education, Universitas Syiah Kuala, Darussalam, Banda Aceh, Indonesia

Technical Team

Husnurrizal



Veterinary, University of Syiah Kuala, Indonesia

Agustiar

Dentistry, University of Syiah Kuala, Indonesia

Indexed by:

Tools:



All papers published in FKG USK Press are licensed under a **Creative Commons Attribution-ShareAlike 4.0 International License**.

Social Media: [Facebook](#) [Twitter](#) [Instagram](#) [YouTube](#)

[Login](#)

JOURNAL CONTENT

Search

Search Scope

All ▾

[Search](#)

Browse

- [By Issue](#)
- [By Author](#)
- [By Title](#)
- [Other Journals](#)
- [Categories](#)

INFORMATION

- [For Readers](#)
- [For Authors](#)
- [For Librarians](#)

VISITORS

Visitors See m

	73,188		95		37
	5,591		78		36
	697		68		33
	576		68		32
	296		50		30
	279		49		30
	198		46		29
	130		46		28
	113		46		22
	113		45		21
	103		42		21
	99		40		15
	98		39		
	95		38		

Histats.com

Vis. today	14
Visits	4 524
Pag. today	40

CURRENT ISSUE

ATOM	1.0
RSS	2.0
RSS	1.0

Antibacterial potential ethanol extract of beluntas leaves (*Pluchea indica* L) to *Streptococcus sanguinis*

Wilma ADIWIJAYA¹, Riani SETIADHI², Vinna Kurniawati SUGIAMAN^{3*}

¹ Faculty of Dentistry, Maranatha Christian University, Bandung, West Java, Indonesia

² Department of Oral Medicine, Faculty of Dentistry, Maranatha Christian University, Bandung, West Java, Indonesia

³ Departement of Oral Biology, Faculty of Dentistry, Maranatha Christian University, Bandung, West Java, Indonesia

* Corresponding Author E-mail: vinnakurniawati@yahoo.co.id

Received; June 12, 2022, Revised; August 18, 2022, Accepted; October 22, 2022

ABSTRACT. Recurrent aphthous stomatitis (RAS) is one of the most common oral diseases in the community, with a prevalence of 5-66%, with one of the predisposing factors being *Streptococcus sanguinis*. Treatment for RAS has been symptomatic and supportive, including antiseptic mouthwash such as chlorhexidine gluconate 0.2% or topical corticosteroids (triamcinolone acetonide 0.1% in Orabase). However, these drugs have some side effects. Treating herbal ingredients such as Beluntas leaves low prices and minimal side effects. The active compounds in Beluntas leaves are phenols, tannins, flavonoids, saponins, triterpenoids, essential oils, terpenoids, and many compounds known to have antibacterial activity. Methods: This study aimed to determine the minimum inhibitory level (MIC), and minimum killing rate (MBC) of 96% ethanol extract of Beluntas leaves on the growth of *Streptococcus sanguinis*. MIC was measured by broth microdilution technique with DMSO solvent 10% and eight concentrations of beluntas extract. Chlorhexidine gluconate 0.2% was used as a positive control for the comparison compound. Furthermore, the MBC test was carried out using the total plate count method for treatments that gave the MIC value. One Way Anova analysis with Post Hoc Tukey was used to determine the significant difference between treatments. Results: The ethanol extract of Beluntas leaves (*Pluchea indica* L) has a Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) for the growth of *Streptococcus sanguinis* with a minimum inhibitory concentration of 3.95 g/mL and a minimum concentration of 7.8 g/mL. Conclusion: The ethanol extract of Beluntas leaves (*Pluchea indica* L) has the potential as an antibacterial against *Streptococcus sanguinis*.

KEYWORDS: Recurrent aphthous stomatitis, *Streptococcus sanguinis*, ethanol extract of beluntas leaves, Minimum Inhibitory Concentration, Minimum Bactericidal Concentration

INTRODUCTION

Recurrent aphthous stomatitis (RAS) is a problem that often arises in dentistry and is one of the most common oral diseases in the community, with a prevalence of 66%. RAS is characterized by recurrent ulcers on the patient's oral mucosa and can be painful. The prevalence of oral ulceration worldwide is as much as 4%, where RAS is the disease with the greatest prevalence of 25% and is experienced by most women in the second and third decades.^{1,2}

The clinical manifestations of RAS are ulcers, single or multiple, shallow, oval, and pain.³ Prodromal symptoms appear before the onset of RAS, including discomfort and redness for 1-3 days.⁴ The exact etiology of RAS has not been determined so the research focused on the predisposing factors

for RAS. Several factors are considered predisposing factors for RAS, including immunological abnormalities, genetics, systemic factors, endocrine system, stress, smoking cessation, allergic factors, and microorganisms. The main microorganisms associated with the formation of RAS are pleomorphic transitional L α -hemolytic *Streptococcus* and *Streptococcus sanguinis*.⁵

One of the efforts to manage RAS is by symptomatic and supportive treatment because, until now, the etiology of RAS is still unknown. Therefore, treatment is only aimed at curing complaints. The goal of symptomatic treatment is to reduce symptoms and the number and size of ulcers. Currently used drugs include: antiseptic mouthwash (chlorhexidine gluconate 0.2%) to reduce the

How to cite this article: Adiwijaya W et al, Antibacterial potential ethanol extract of beluntas leaves (*Pluchea indica* L) to *Streptococcus sanguinis*. JDS. 2022; 7(2): 78-85

duration and discomfort of RAS or topical corticosteroids (triamcinolone acetonide 0.1% in Orabase).⁶ Chlorhexidine gluconate 0.2% is the gold standard, effective against gram-negative and positive and facultative aerobes and anaerobes. However, its use in more than two weeks has side effects, including causing burning of the oral mucosa, impaired taste, tooth staining, erosion of the oral surface, and dryness of the oral cavity.⁷

Therefore, it is necessary to develop an alternative therapy by utilizing herbal plants in curing ulcers in RAS. It is intended so that patients can recover without side effects at a relatively low cost.⁸ The development of a back-to-nature lifestyle has made herbal plants increasingly widespread by the community as alternative materials for treating or preventing certain diseases. The community considers that herbal ingredients are easier to obtain, the price is more affordable, and the side effects are minimal, so it is safer for consumption.⁹ One of the natural ingredients that can be used in traditional medicine is the beluntas plant.¹⁰ The leaves, flowers, and roots of beluntas can be used for therapy, but the beluntas leaves are the part that has the highest biologically active component.¹¹

Beluntas (*Pluchea indica* L) is a wild plant in dry areas on hard and rocky soils or grown as a hedge.¹² This plant has a distinctive aromatic odor and bitter

taste. The part used from this plant is the leaves and roots, which are efficacious for eliminating body odor and bad breath, increasing appetite, overcoming digestive disorders in children, relieving pain in rheumatism, and so on. The active compounds in beluntas leaves are flavonoids, triterpenoids, phenols, and essential oil derivatives.¹³ Beluntas leaves have antibacterial properties due to the content of phenolics, triterpenoids, and tannins that can cause bacterial cell death.^{14,15}

Based on several studies that have been carried out, the ethanol extract of beluntas leaves (*Pluchea indica* L) has been shown to inhibit bacterial growth. These bacteria include *Streptococcus mutans*, *Staphylococcus aureus*, *Bacillus subtilis*, and *Pseudomonas aeruginosa*.

Based on the explanation above and there is no research on *Streptococcus sanguinis* bacteria, so the authors are interested in research to determine the antibacterial effect of beluntas leaves through testing the Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) of the ethanol extract of the beluntas leaves on the growth of *Streptococcus sanguinis* which is one of the bacteria that causes recurrent aphthous stomatitis.

MATERIALS AND METHODS

Determination and Ethanol Extract of Beluntas Leaves

Beluntas leaves were obtained from the Subang Tropical Balitbu plantation, which was then determined at the Biology Laboratory of Padjadjaran University. Extract beluntas leaves with 96% ethanol and let stand for three days, then the extract is filtered using filter paper to obtain the filtrate. Then the filtrate was evaporated using a rotary evaporator at a temperature of 70°C until the consistency was like a paste.

The ethanol extract of beluntas leaves was put into a 15 mL tube with as much as 0.5 grams and dissolved in 10% DMSO to reach a concentration of 500 g/mL. The extract was diluted using a multichannel Micropipette so that the final concentrations were 250g/ml, 125g/ml, 62.5g/ml, 31.3g/ml, 15.6g/ml, 7.8g/ml, 3.95g/ml, and 1.95g/ml.

Preparation of *Streptococcus sanguinis* inoculum

The bacteria used in this study is *Streptococcus sanguinis* ATCC 10556. Take two oses and suspension in BHI-B, incubated for 24 hours. Then standardized with 0.5 McFarland solution, the total

density of bacteria used is 1.5×10^8 CFU/ml. Check using a spectrophotometer with a wavelength of 600nm, then dilute to 10^5 .

Minimum Inhibitory Level (MIC) and Minimum Killing Rate (KBM)

The method used in this study followed the method developed by CLSI (Clinical Laboratory Standard Institute) with a slight modification, namely adding 2% sucrose to BHI-A and BHI-B media.

This study consisted of ten treatments, namely eight treatments of ethanol extract of beluntas leaves with concentrations of 250g/ml; 125g/ml; 62.5g/ml; 31.3g/ml; 15.6g/ml; 7.8g/ml; 3.95g/ml, and

1.95g/ml; positive control (chlorhexidine gluconate 0.2%); and negative control (DMSO).

200 L/mL of BHI-Broth media was added to all 96 well plates and added 200 l of ethanol extract of beluntas leaves of each concentration, and 10 l of bacterial suspension, with the format of media+sample (negative control), media+solvent (solvent control), media+sample+bacteria (test sample), media+solvent+bacteria (positive control). Incubate at 37°C for 16-20 hours. Insert the well plate into a spectrophotometer with a wavelength of 600 nm to see the minimum inhibitory concentration using the absorbance method.

Take 200 L of 0.2% Chlorhexidine gluconate and put it into the well plate with 3 repetitions, 10 l

of bacterial suspension 10⁵ for two repetitions, and one repetition plate without bacteria as a positive control. Incubate for 24 hours at 37°C with 5% CO₂.

96-well plates indicated as MIC and KBM were inoculated into sterile BHI-Agar medium in Petri dishes and incubated at 37°C with 5% CO₂ for 24 hours, calculating the number of colonies in each petri dish from each treatment with various concentrations of extract using a colony counter.

The number of colonies contained in BHI-Agar media was used to determine MIC, and MBC was calculated using a Colony counter. The calculation of the number of colonies formed is called the total plate count (TPC), with colony forming units/mL (CFU/mL) calculated by the formula:

$$\text{TPC value (CFU/ml)} = \frac{\text{Number of colonies formed} \times \text{Dilution factor}}{\text{The volume of bacterial colonies that is included in the criteria}}$$

The formula measures the calculation of the percentage of bactericidal power:

$$\% \text{ Kill Efficiency} = \frac{\text{TPC bacteria control (CFU/ml)} - \text{TPC treatment (CFU/ml)} \times 100\%}{\text{Mean bacterial colony control}}$$

RESULTS

The Beluntas plants' determination results at the Biology Laboratory of Padjadjaran University, Jatinangor, and West Java showed that the sample used was Beluntas (*Pluchea indica* L). Phytochemical tests were carried out at the Central Laboratory of

Padjadjaran University, Jatinangor, West Java. Table 1 shows the results of qualitative phytochemical tests taken from the ethanol extract of beluntas leaves.

Table 1. Qualitative Phytochemical Test Results

No	Secondary Metabolite	Methods	Results
1	Phenolic	Reagent FeCl ₃ 5%	++
2	Tanin	Reagent FeCl ₃ 1%	++
3	Flavonoid	a. Reagent HCL concentrated + Mg	-
		b. Reagent H ₂ SO ₄ 2N	-
		c. Reagent NaOH 10	+
4	Saponin	Heated	+
5	Triterpenoid	Reagent H ₂ SO ₄ concentrated + CH ₃ COOH	++
	dan Steroid	anhydrous	-
6	Alkaloid	Reagent Dragendorff	+

Description: +: Little; ++: Medium; +++: Much ; -: None

Minimum Inhibitory Level (MIC)

The results of the MIC test observed the growth of *Streptococcus sanguinis*, which was indicated by turbidity at the bottom of the well in several treatments compared to negative controls.

The results of visual observations can be seen in the good extract of beluntas at a concentration of 1.95g/mL. There is turbidity at the bottom which is a culture of *Streptococcus sanguinis*. There was slight

turbidity at well 3.9g/mL, while at wells 7.8 to 250, visually, no difference in turbidity was observed.

Solvent control and clear positive control showed no contamination in the experiment.



Figure 1. The plate of minimum inhibitory content test results

The average results of the absorbance of the effect of the ethanolic extract of beluntas leaves on the number of colonies of *Streptococcus sanguinis* using spectrophotometry with a wavelength of

600nm, resulting in quantitative data that confirmed that the MIC was right at a concentration of 3.9 g/mL. (Table 2)

Table 2. The absorbance of ethanol extract of beluntas leaves against *Streptococcus sanguinis*

Extract Concentration	Media control	Average Absorbance Bacteria+ Sample	Difference	Note(s)
250µg/mL	2,039	2,018	0,021	
125µg/mL	1,4203	1,3964	0,023	
62.5µg/mL	0,8977	0,8568	0,04	
31.25µg/mL	0,5208	0,4938	0,27	
15.62µg/ mL	0,312	0,2874	0,25	
7.8µg/mL	0,1873	0,1789	0,84	
3.9µg/mL	0,1245	0,114	0,1	MIC
1.95µg/mL	0,0795	0,0833	-0,0035	

The results of the positive control test (chlorhexidine gluconate 0.2%) obtained the average number of colonies of *Streptococcus sanguinis* bacteria

with three repetitions after 24 hours of incubation was 0 CFU/mL with an average bactericidal power of 100%.

Minimum Bactericidal Concentration (MBC)

The results of the MBC test showed that at a concentration of 7.8g/mL, the ethanol extract of the tested beluntas leaves was not observed for the growth of *Streptococcus sanguinis*. To count the

number of bacteria at each test concentration, the number of bacteria was calculated using the total plate count technique with a colony counter, as shown in Figure 2 and Table 3.




Figure 2. The results of the calculation of the number of bacteria using the Colony Counter
Notes: (A) Number of *Streptococcus sanguinis* Bacteria Colonies at the Concentration of Ethanol Extract of Beluntas Leaves 7.8g/mL (KBM). (B) Number of *Streptococcus sanguinis* Bacteria Colonies at Concentration of Ethanol Extract of Beluntas Leaves 3.9g/mL (MIC)

Table 3. Number of *Streptococcus sanguinis* Bacteria Colonies After 24 hours of incubation at various concentrations

Extract Concentration	Mean Colony (CFU/mL)	Mean kill Efficiency	Note(s)
Group A (250 µg/mL)	0	100%	
Group B (125 µg/mL)	0	100%	
Group C (62.5 µg/mL)	0	100%	
Group D (31.25 µg/mL)	0	100%	
Group E (15.62 µg/mL)	0	100%	
Group F (7.8 µg/mL)	0	100%	MBC
Group G (3.9 µg/mL)	12,9x 10 ¹	99%(0,9999612)	MIC
Group H (1.95 µg/mL)	27,3 x 10 ⁵	99%(0,9999178)	

Colony Number Normality Test Results

The normality test results using Shapiro-Wilk showed that two variables were tested, namely concentrations of 3.95g/mL and 1.95g/mL. It is known that the significance value is 0.147 and 0.114,

where the value is more significant than with a value of 0.05. Thus, it can be concluded that the data is normally distributed.

One-Way ANOVA Analyses

The results of the One Way ANOVA test to determine whether the data has an average difference or cannot be known from the significance value of the number of colonies then compared with the significance value, with a test decision, if the value (Sig) > 0.05, then there is no difference in average if value (Sig) < 0.05 then there is an average

difference. The results of the ANOVA test above show the F value for the number of colonies of 22.882 and the significance (p-value) of 0.000 is less than 0.05, so it can be concluded that at concentrations of 3.95 and 1.95, there was a different effect on the number of colonies.

How to cite this article: Adiwijaya W et al, Antibacterial potential ethanol extract of beluntas leaves (*Pluchea indica L*) to *Streptococcus sanguinis*. JDS. 2022; 7(2): 78-85

Tukey's Post-Hoc Test Results

Based on the results of the Tukey Post Hoc test in appendix 3, the number of colonies at a concentration of 250 was the same as a concentration of 125, 62.5, 31.3, 15.6, 7.8, and 1.95 because they were in the same subset. Meanwhile, the number of colonies at a concentration of 3.95 was not the same

as that of 125, 62.5, 31.3, 15.6, 7.8, and 1.95 because they were indifferent subsets. The comparison results can be seen through the mean difference column, obtained at a concentration of 3.95 significantly different ($p < 0.05$).

DISCUSSION

Based on the research results presented in **Table 2**, it was found that the ethanol extract of beluntas leaves had an inhibitory and killing effect on the growth of *Streptococcus sanguinis* bacteria and there was MIC at a concentration of 3.9g/mL with 99% killing power and MBC was found at a concentration of 7.8g/mL. mL with 100% killing power. The positive control group of chlorhexidine gluconate at a concentration of 0.2% had an inhibitory and killing effect on *Streptococcus sanguinis* bacteria with an average killing power of 100%. Therefore, it can be said that the ethanol extract of beluntas leaves at a concentration of 7.8g/mL had the same antibacterial effect as 0.2% chlorhexidine gluconate. The statistical test value obtained a P-value of 0.00, which states that the MIC and MBC values significantly affect the growth of *Streptococcus sanguinis* bacteria.

These results follow previous research, which stated that the ethanol extract of beluntas leaves (*Pluchea indica* L) could inhibit the growth of bacteria, *Staphylococcus aureus*, *Bacillus subtilis*, and *Pseudomonas aeruginosa*.^{16,17} Based on the results of qualitative phytochemical tests, active compounds in beluntas leaves, such as phenols, tannins, flavonoids, saponins, triterpenoids, and alkaloids. Phenol compounds have antibacterial properties due to the presence of hydroxyl and carbonyl groups that can interact with bacterial cells through hydrogen bonds. It will cause the coagulation of proteins and bacterial cell membranes, which will cause bacteria to lyse.¹⁸ Phenol can also cause hyperpolarization of the cytoplasmic membrane and increase the instability membrane. It causes membrane dysfunction and bacterial cell death.^{14,15}

Tannins are one type of compound that belongs to the polyphenol group and has soluble properties in water and organic solvents. Tannins can be used as antibacterial because they have an antiseptic phenol group, so they can be used as an antibacterial component through hydrogen bonds which will form complex compounds with proteins and will cause bacterial cell proteins to be denatured so that bacterial metabolism is disrupted.^{14,15,19}

Flavonoids also have a role as antibacterial compounds by inhibiting cell membrane function, nucleic acid synthesis, and inhibiting energy metabolism. The position in inhibiting the synthesis of nucleic acids occurs through the accumulation of nucleic acid bases, thereby inhibiting the formation of DNA and RNA because rings A and B of flavonoid compounds play an important role in the process of intercalation or hydrogen bonding. The result of flavonoid interaction will also cause damage to the permeability of the bacterial cell wall. Flavonoids will also form complex extracellular compounds and soluble proteins that cause damage to cell membranes and the release of intracellular compounds, thus causing cell function to be inhibited. In addition, energy metabolism will also be hampered due to the inhibition of oxygen used by bacteria. It can occur by preventing the formation of energy in the cytoplasmic membrane and inhibiting the motility of bacteria which play a role in antimicrobial activity and extracellular proteins.^{20,21}

Other components, namely saponins, have broad biological activity and act as antibacterial and antifungal agents. Saponins can cause leakage of bacterial cell membranes, resulting in damage to membrane permeability which can interfere with the survival of bacteria. When the stability of the bacterial cell membrane is disturbed, it will cause the cytoplasm to come out of the cell. It occurs when saponins bind to the cytoplasmic membrane through the outer membrane, which will cause the stability of the bacterial cell membrane to be disturbed and result in bacterial cell lysis and then death.^{21,22}

Triterpenoid compounds are also reported to have antibacterial activity. Triterpenoids will react with porins (transmembrane proteins) on the outer wall of the bacterial cell membrane causing the formation of strong polymer bonds. This condition will cause damage to the porin, which is the entrance and exit of the compound. Furthermore, this causes the permeability of the bacterial cell wall to decrease, the bacterial cell to lysis, and the death of the bacterial cell.^{14,15,23,24}

Alkaloids also have antibacterial properties that cause bacterial cell death because the cell wall layer is not completely formed by interfering with the peptidoglycan constituent components of bacterial cells by interfering with the formation of cross-bridges of peptidoglycan constituent components in bacterial cells. Alkaloids can also inhibit enzymes that play a role in the DNA replication process, inhibiting bacterial growth

because bacteria cannot divide by inhibiting DNA replication.^{25,26}

Because beluntas leaves have an active biological content that acts as an antibacterial, the ethanol extract of beluntas leaves can be developed and used as natural antibacterial ingredient because there is a significant value at a concentration of 7.8g/mL with the number of colonies of *Streptococcus sanguinis* bacteria 0 CFU/mL.

CONCLUSION

The ethanol extract of beluntas leaves (*Pluchea indica* L) has the potential as an antibacterial against *Streptococcus sanguinis*.

REFERENCES

- [1]. Amtha R, Marcia M, Aninda AI. Plester sariawan efektif dalam mempercepat penyembuhan stomatitis aftosa rekuren dan ulkus traumatikus. Yogyakarta: Majalah Kedokteran Gigi Indonesia. 2017;3(2):7.
- [2]. Apriasari ML, Tuti, H. Stomatitis aftosa rekuren oleh karena anemia. J Dentofasial. 2010; 9(1): 39-46.
- [3]. Langlais RL, Miller CS. Kelainan Rongga Mulut yang Lazim. Jakarta: Hipokrates. 2012. p. 1.
- [4]. Odell E. Cawson's Essentials of Oral Pathology and Oral Medicine. New York: Elsevier. 2017;9(1):220-221.
- [5]. Ghom AG. Textbook of Oral Medicine. New Delhi: Jaypee Brothers Medical Publishers; 2014: 360.
- [6]. Widyastutik O, Permadi A. Penatalaksanaan Nyeri pada Stomatitis Aftosa Rekuren. Jakarta: Journal FKG UNEJ. 2011; 4(3): 396-397.
- [7]. Dutt P, Rathore PK, Khurana D. Chlorhexidine- An antiseptic in periodontics. IOSR-JDMS. 2014;13(9): 85-8.
- [8]. Sari LORK. Pemanfaatan Obat Tradisional Dengan Pertimbangan Manfaat Dan Keamanannya. Depok: Journal Pharmaceutical Sciences and Research. 2016; 3(1): 1-7.
- [9]. Saparinto C, Susiana R. Grow Your Own Medical Plant. Yogyakarta: Lily publisher. 2016: 456-473.
- [10]. Amilah S, Ajiningrum PS. Uji Efektivitas Daya Hambat Sari Daun Pegagan (*Centella asiatica*) dan Daun Beluntas (*Plunchea indica* Less) terhadap Pertumbuhan *Mycobacterium* Tuberculosis. STIGMA Journal science. 2015; 8(2): 6-11.
- [11]. Muta'ali R, Kristanti IP. Pengaruh Ekstrak Daun Beluntas (*Pluchea indica*) terhadap Mortalitas dan Perkembangan Larva *Spodoptera litura* F. Sumatera: Journal Sains dan seni ITS. 2015; 4(2): 6-7.
- [12]. Arishandi Y, Andriani Y. Tanaman Obat Plus Pengobatan Alternatif. Jakarta: Setia Kawan. 2010.
- [13]. Silalahi M. Keanekaragaman tumbuhan bermanfaat di pekarangan oleh Etnis Sunda Di Desa Sindang Jaya Kabupaten Cianjur Jawa Barat. Jurnal Pendidikan Matematika Dan IPA. 2019; 10(1): 88-104.
- [14]. Raharjo I, Horsten S. *Pluchea indica* in: Plant Resources of South East Asia. Bogor Indonesia: JLCH van Valkenburg and N Bunyapraphatsara. 2002; 12(2): 441-443.
- [15]. Gayatri DAAM, Ernawati DK, Widhiartini IAA. Uji Efektifitas Ekstrak Etanol Daun Beluntas Terhadap Pertumbuhan Bakteri *Streptococcus pyogenes* ATCC 19615 secara in vitro. Bali: Jurnal Medika Udayana. 2020; 10(1): 7-11.
- [16]. Nahak MM, Tedjasulaksana R, Dharmawati IGGA. Khasiat Ekstrak Daun Beluntas untuk Menurunkan Jumlah Bakteri pada Saliva. Journal Interdental Kedokteran gigi. 2017; 5(3): 139-42
- [17]. Manu RRS. Aktivitas Antibakteri Ekstrak Etanol Daun Beluntas (*Pluchea indica* L.) Terhadap *Staphylococcus Aureus*, *Bacillus Subtilis* Dan *Pseudomonas Aeruginosa*. Surabaya: Jurnal Ilmiah Mahasiswa Universitas Surabaya. 2013; 2(1): 1-10.

- [18]. Ndruru ESC. Perbandingan Efektivitas Berkumur Larutan Probiotik dan Klorheksidin Terhadap Akumulasi Plak dan Jumlah Streptococcus Mutans pada Anak Usia 12-15 Tahun Di Yayasan Sos Children's Village. Medan: Fakultas Kedokteran Gigi Universitas Sumatera Utara. 2021.
- [19]. Utomo SB. Uji Aktivitas Antibakteri Senyawa C-4-Metoksifenil kaliks Resorsinarena Termodifikasi Hexadecyltrimethylammonium-Bromide Terhadap Bakteri Staphylococcus Aureus dan Escherichia Coli. Surakarta: Journal UNS. 2018.
- [20]. Nomer NM, Duniaji AS, Nocianitri KA. Kandungan senyawa flavonoid dan antosianin ekstrak kayu secang (caesalpinia sappan L.) serta aktivitas antibakteri terhadap vibrio cholerae. Bali: Journal Ilmu dan Teknologi Pangan. 2019; 8(2): 216-225.
- [21]. Haqi HD. Aktivitas Antibakteri Ekstrak Metanol Serbuk Biji Kluwih (Artocarpus Communis J.R. & G) Terhadap Pertumbuhan Methicillin Resistant *Staphylococcus Aureus* (MRSA). Yogyakarta: Fakultas Kedokteran Gigi Universitas Muhammadiyah. 2018.
- [22]. Pincus. Clinical Diagnosis and Management by Laboratory Methods. New York: Elsevier. 2011.
- [23]. Makkar HPS, Becker K. Do Tannins In Leaves Of Trees And Shrubs From African And Himalayan Regions Differ In Level And Activity Agroforestry System. Heidelberg: Agroforestry Systems. 2003.
- [24]. Retnowati Y, Bialangi N, Posangi NW. Pertumbuhan Bakteri Staphylococcus aureus Pada Media Yang Diekspos Dengan Infus Daun Sambiloto (*Andrographis paniculata*). Gorontalo: Jurnal Saintek. 2011; 6(2).
- [25]. Taufiq S, Umi Y, Siti, H. Uji Aktivitas Antibakteri Ekstrak Etanol Biji Buah Pepaya (*Carica papaya L.*) terhadap *Escherichia coli* dan *Salmonella typhi*. Bandung: Prosiding Penelitian Spesia Unisba. 2015.
- [26]. Eranawati & Sari K. Kandungan senyawa kimia dan aktivitas antibakteri ekstrak kulit buah alpukat (*Persea americana P.Mill*) terhadap bakteri *Vibrio alginolyticus*. Jurnal Kajian Veteriner. 2015; 3(2): 203-211.