

BUKTI KORESPONDENSI

ARTIKEL JURNAL NASIONAL TERAKREDITASI


Judul Artikel : Antibacterial Potential Ethanol Extract of Beluntas Leaves (*Pluchea indica* L) to *Streptococcus sanguinis*

Jurnal : Journal of Syiah kuala Dentistry Society

Penulis : Wilma Adiwijaya, Riani Setiadhi, Vinna Kurniawati Sugiaman

No	Perihal	Tanggal
1.	Register pada Journal of Syiah kuala Dentistry Society	17 Juli 2022
2.	Bukti konfirmasi submit artikel dan artikel yang disubmit	17 Juli 2022
3.	Bukti melakukan review yang pertama	17 Juli 2022
4.	Bukti konfirmasi submit revisi pertama yang telah direvisi	20 Juli 2022
5.	Bukti melakukan review yang kedua	-
6.	Bukti konfirmasi submit artikel yang telah revisi kedua	-
7.	Bukti konfirmasi artikel diterima	10 Desember 2022
8.	Bukti Galery Proof Manuscript	Desember 2022
9.	Bukti Publikasi Online Artikel	Desember 2022

Register pada Journal of Syiah kuala Dentistry Society (17 Juli 2022)



Dentistry Faculty
Dari: jurnal@unsyiah.ac.id
Kepada: Vinna Kurniawati Sugiaman

Min, 17 Jul 2022 jam 15:22 ☆

Vinna Kurniawati Sugiaman


You have now been registered as a user with Journal of Syiah Kuala Dentistry Society. We have included your username and password in this email, which are needed for all work with this journal through its website. At any point, you can ask to be removed from the journal's list of users by contacting me.

Username: vinna
Password: ukm12345*

Thank you,
Dentistry Faculty

Journal of Syiah Kuala Dentistry Society
<http://jurnal.unsyiah.ac.id/JDS>

Bukti konfirmasi submit artikel dan artikel yang disubmit (17 juli 2022)



Antibacterial Potential Ethanol Extract of Beluntas Leaves (*Pluchea indica* L) to *Streptococcus sanguinis*

Wilma Adiwijaya¹, Riani Setiadhi², Vinna Kurniawati Sugiaman^{3*}

¹ Faculty of Dentistry, Maranatha Christian University, Bandung, West Java, Indonesia

²Departement of Oral Medicine, Faculty of Dentistry, Maranatha Christian University, Bandung, West Java, Indonesia

³Departement of Oral Biology, Faculty of Dentistry, Maranatha Christian University, Bandung, West Java, Indonesia

* Correspondence:

Vinna Kurniawati Sugiaman

Faculty of Dentistry, Maranatha Christian University, Bandung, West Java, Indonesia

Email address: vinnakurniawati@yahoo.co.id.

Phone number: +62 22 2012692 (ext 1906)

ABSTRACT

Background: Recurrent aphthous stomatitis (RAS) is one of the most common oral diseases in the community with a prevalence of 5-66%, with one of the predisposing factors is *Streptococcus sanguinis*. Treatment for RAS

has been symptomatic and supportive, including antiseptic mouthwash such as chlorhexidine gluconate 0.2% or topical corticosteroids (triamcinolone acetonide 0.1% in Orabase), however these drugs have some side effects. Treatment using herbal ingredients such as beluntas leaves relatively low prices and minimal side effects. The active compounds contained in Beluntas leaves are phenols, tannins, flavonoids, saponins, triterpenoids, essential oils, terpenoids, and many compounds are known to have antibacterial activity. Methods: This study aimed to determine the minimum inhibitory level (MIC) and minimum killing rate (MBC) of 96% ethanol extract of beluntas leaves on the growth of *Streptococcus sanguinis*. MIC was measured by broth microdilution technique with DMSO solvent 10% and 8 concentrations of beluntas extract. Chlorhexidine gluconate 0.2% was used as a positive control for the comparison compound. Furthermore, the MBC test was carried out using the total plate count method for treatments that gave the MIC value. One Way Anova analysis with Post Hoc Tukey was used to determine the significant difference between treatments. Results: The ethanol extract of beluntas leaves (*Pluchea indica* L) has a Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) for the growth of *Streptococcus sanguinis* with a minimum inhibitory concentration of 3.95 g/mL and a minimum concentration of 7.8 g/mL. Conclusion: The ethanol extract of beluntas leaves (*Pluchea indica* L) has potential as an antibacterial against *Streptococcus sanguinis*.

Keywords: Recurrent aphthous stomatitis, *Streptococcus sanguinis*, ethanol extract of beluntas leaves, Minimum Inhibitory Concentration, Minimum Bactericidal Concentration.

Introduction

Recurrent aphthous stomatitis (RAS) is a problem that often arises in dentistry and is one of the most common oral diseases in the community with a prevalence of 66%. RAS is characterized by recurrent ulcers on the patient's oral mucosa and can be painful. The prevalence of oral ulceration worldwide is known to be as much as 4%, where RAS is the disease with the greatest prevalence of 25% and is experienced by most women in the second and third decades.^{1,2}

The clinical manifestations of RAS are ulcers, single or multiple, shallow, oval, and pain.³ Prodromal symptoms appear before the onset of RAS including discomfort and redness for 1-3 days.⁴ The exact etiology of RAS has not been determined so the research focused on the predisposing factors for RAS. Several factors are considered as predisposing factors for RAS, including immunological abnormalities, genetics, systemic factors, endocrine system, stress, smoking cessation, allergic factors, and microorganisms. The main microorganisms associated with the formation of RAS are pleomorphic transitional L α -hemolytic *Streptococcus* and *Streptococcus sanguinis*.⁵

One of the efforts to manage RAS is by symptomatic and supportive treatment because until now the etiology of RAS is still unknown. Therefore, treatment is only aimed at curing complaints. The goal of symptomatic treatment is to reduce symptoms, reduce the number and size of ulcers. Currently used drugs include: antiseptic mouthwash (chlorhexidine gluconate 0.2%) to reduce the duration and discomfort of RAS or topical corticosteroids (triamcinolone acetonide 0.1% in Orabase).⁶ Chlorhexidine gluconate 0.2% is the gold standard. which is effective against both gram negative and positive and facultative aerobes and anaerobes. However, its use in more than 2 weeks has side effects, including causing burning of the oral mucosa, impaired taste, tooth staining, erosion of the oral surface, and dryness of the oral cavity.⁷

Therefore, it is necessary to develop an alternative therapy by utilizing herbal plants in curing ulcers in RAS. This is intended so that patients can recover without side effects at a relatively low cost.⁸ The development of a back to nature lifestyle has made the use of herbal plants increasingly widespread by the community as alternative materials for the treatment or prevention of certain diseases. The community considers that herbal ingredients are easier to obtain, the price is more affordable, and the side effects are minimal, so it is safer for consumption.⁹ One of the natural ingredients that can be used as traditional medicine is the beluntas plant.¹⁰ The leaves, flowers, and roots of beluntas can be used for therapy, but the beluntas leaves are the part that has the highest biologically active component.¹¹

Beluntas (*Pluchea Indica* L) is a wild plant in dry areas on hard and rocky soils or grown as a hedge.¹² This plant has a distinctive aromatic odor and bitter taste. The part used from this plant is the leaves and roots which are efficacious for eliminating body odor and bad breath, increasing appetite, overcoming digestive disorders in children, relieving pain in rheumatism, and so on. The active compounds contained in beluntas leaves are flavonoids, triterpenoids, phenols, and essential oil derivatives.¹³ Beluntas leaves have antibacterial properties due to the content of phenolics, triterpenoids, and tannins that can cause bacterial cell death.^{14,15}

Based on several studies that have been carried out, the ethanol extract of beluntas leaves (*Pluchea indica* L) has been shown to inhibit bacterial growth. These bacteria including *Streptococcus mutans*, *Staphylococcus aureus*, *Bacillus subtilis*, and *Pseudomonas aeruginosa* bacteria. Based on the explanation above and there is no research on *Streptococcus sanguinis* bacteria, so the authors are interested in research to determine the antibacterial effect of beluntas leaves through testing the Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) of the ethanol extract of the beluntas leaves on the growth of *Streptococcus sanguinis* which is one of the bacteria that causes recurrent aphthous stomatitis.

Method

Determination and Ethanol Extract of Beluntas Leaves

Beluntas leaves were obtained from the Subang Tropical Balitbu plantation which were then determined at the Biology Laboratory of Padjadjaran University. Extract beluntas leaves with 96% ethanol and let stand for 3 days, then the extract is filtered using filter paper to obtain the filtrate. Then the filtrate was evaporated using a rotary evaporator at a temperature of 70oC until the consistency was like a paste.

The ethanol extract of beluntas leaves was put into a 15mL tube as much as 0.5 grams and dissolved in 10% DMSO to reach a concentration of 500 g/mL. The extract was diluted using a multichannel Micropipette so that the final concentrations were 250 g/ml, 125 g/ml, 62.5 g/ml, 31.3 g/ml, 15.6 g/ml, 7.8 g/ml, 3.95 g/ml, and 1.95 g/ml.

Preparation of *Streptococcus sanguinis* inoculum

The bacteria that will be used in this study is *Streptococcus sanguinis* ATCC 10556. Take 2 oses and suspension in BHI-B, incubated for 24 hours. Then standardized with 0.5 McFarland solution so that the total density of bacteria used is 1.5×10^8 CFU/ml. Check using a spectrophotometer with a wavelength of 600 nm, then diluted to 105.

Minimum Inhibitory Level (MIC) and Minimum Killing Rate (KBM)

The method used in this study was following the method developed by CLSI (Clinical Laboratory Standard Institute) with a slight modification, namely adding 2% sucrose to BHI-A and BH-B media.

This study consisted of ten treatments, namely eight treatments of ethanol extract of beluntas leaves with concentrations of 250 g/ml, 125 g/ml, 62.5 g/ml, 31.3 g/ml, 15.6 g/ml, 7,8 g/ml, 3.95 g/ml, and 1.95 g/ml; positive control (chlorhexidine gluconate 0.2%); and negative control (DMSO).

200 L/mL of BHI-Broth media was added to all 96 well plates and added 200 l of ethanol extract of beluntas leaves of each concentration and 10 l of bacterial suspension, with the format of media + sample (negative control), media + solvent (solvent control), media + sample + bacteria (test sample), media + solvent + bacteria (positive control). Incubate at 37°C for 16-20 hours. Insert the well plate into a spectrophotometer with a wavelength of 600 nm to see the minimum inhibitory concentration using the absorbance method.

Take 200 L of 0.2% Chlorhexidine gluconate and put it into the well plate with 3 repetitions, 10 l of bacterial suspension 105 for 2 repetitions and 1 repetition plate without bacteria as a positive control. Incubate for 24 hours at 37oC with 5% CO2.

Well plates indicated as MIC and KBM were inoculated into sterile BHI-Agar medium in petri dishes and incubated at 37oC with 5% CO2 for 24 hours, calculating the number of colonies in each petri dish from each treatment with various concentrations of extract using a colony counter.

The number of colonies contained in BHI-Agar media was used to determine MIC and MBC was calculated using a Colony counter. The calculation of the number of colonies formed is called the total plate count (TPC) with colony forming units/mL (CFU/mL) calculated by the formula:

$$\text{TPC value (CFU/ml)} = \frac{\text{Number of colonies formed} \times \text{Dilution factor}}{\text{The volume of bacterial colonies that is included in the criteria}}$$

The calculation of the percentage of bactericidal power is measured by the formula:

$$\% \text{ Kill Efficiency} = \frac{\text{TPC bacteria control (CFU/ml)} - \text{TPC treatment (CFU/ml)} \times 100\%}{\text{Mean bacterial colony control}}$$

Results

The results of the determination of Beluntas plants carried out at the Biology Laboratory of Padjadjaran University, Jatinangor, West Java showed that the sample used was Beluntas (*Pluchea Indica* L). Phytochemical tests were carried out at the Central Laboratory of Padjadjaran University, Jatinangor, West Java. Table 4.1 shows the results of qualitative phytochemical tests taken from the ethanol extract of beluntas leaves.

Table 4.1 Qualitative Phytochemical Test Results

No	Secunder Metabolite	Methods	Results
1	Fenolik	Reagent FeCL ₃ 5%	++
2	Tanin	Reagent FeCL ₃ 1%	++
3	Flavonoid	a. Reagent HCL concentrated + Mg	-
		b. Reagent H ₂ SO ₄ 2N	-
		c. Reagent NaOH 10	+
4	Saponin	Heated	+
5	Triterpenoid	Reagent H ₂ SO ₄ concentrated + CH ₃ COOH	++
	dan Steroid	anhydrous	-
6	Alkaloid	Reagent Dragendorff	+

Description: + : Little; ++ : Medium; +++ : Much ; - : None

Minimum Inhibitory Level (MIC)

The results of the MIC test observed the growth of *Streptococcus sanguinis* which was indicated by the presence of turbidity at the bottom of the well in several treatments when compared to negative controls. The results of visual observations can be seen in the well extract of beluntas at a concentration of 1.95 g/mL there is turbidity at the bottom which is a culture of *Streptococcus sanguinis*. There was slight turbidity at well 3.9 g/mL, while at wells 7.8 to 250, visually no difference in turbidity was observed. Solvent control and clear positive control showed no contamination in the experiment.

The results of the positive control test (chlorhexidine gluconate 0.2%) obtained the average number of colonies of *Streptococcus sanguinis* bacteria with 3 repetitions after 24 hours of incubation was 0 CFU/mL with an average bactericidal power of 100%.

Minimum Bactericidal Concentration (MBC)

The results of the MBC test showed that at a concentration of 7.8 g/mL the ethanol extract of the tested beluntas leaves was not observed for the growth of *Streptococcus sanguinis*. To count the number of bacteria at each test concentration, the number of bacteria was calculated using the total plate count technique with a colony counter as shown in Figure 2 and Table 2.



Figure 2. The results of the calculation of the number of bacteria using the Colony Counter

Notes: (A) Number of *Streptococcus sanguinis* Bacteria Colonies at the Concentration of Ethanol Extract of Beluntas Leaves 7.8 g/mL (KBM). (B) Number of *Streptococcus sanguinis* Bacteria Colonies at Concentration of Ethanol Extract of Beluntas Leaves 3.9 g/mL (MIC)

Table 2. Number of *Streptococcus sanguinis* Bacteria Colonies After 24 hours incubation at various concentrations of Beluntas Leaf Ethanol Extract

Extract Concentration	Mean Colony (CFU/mL)	Mean kill Efficiency	Note(s)
Group A (250 µg/mL)	0	100%	
Group B (125 µg/mL)	0	100%	
Group C (62.5 µg/mL)	0	100%	
Group D (31.25 µg/mL)	0	100%	

Group E (15.62 µg/ mL)	0	100%	
Group F (7.8 µg/ mL)	0	100%	KBM
Group G (3.9 µg/mL)	12,9x 10 ¹	99%(0,9999612)	KHM
Group H (1.95 µg/mL)	27,3 x 10 ⁵	99%(0,9999178)	

Colony Number Normality Test Results

The results of the normality test using Shapiro-Wilk showed that there were 2 variables tested, namely concentrations of 3.95 g/mL and 1.95 g/mL. it is known that the significance value is 0.147 and 0.114, where the value is greater than with a value of 0.05, thus it can be concluded that the data is normally distributed.

One Way ANOVA Test Results

The results of the One Way ANOVA test to determine whether the data has an average difference or cannot be known from the significance value of the number of colonies then compared with the significance value, with a test decision, if the value (Sig) > 0.05 then there is no difference in average, if value (Sig) < 0.05 then there is an average difference. The results of the ANOVA test above show the F value for the number of colonies of 22.882 and the significance (p-value) of 0.000 is less than 0.05, so it can be concluded that at concentrations of 3.95 and 1.95 there was a different effect on the number of colonies.

Tukey's Post Hoc Test Results

Based on the results of the Tukey Post Hoc test in appendix 3, the number of colonies at a concentration of 250 was the same as a concentration of 125, 62.5, 31.3, 15.6, 7.8, and 1.95 because they were in the same subset. Meanwhile, the number of colonies at a concentration of 3.95 was not the same as a concentration of 125, 62.5, 31.3, 15.6, 7.8, and 1.95 because they were indifferent subsets. The comparison results can be seen through the mean difference column, obtained at a concentration of 3.95 significantly different (p < 0.05).

Discussion

Based on the research results presented in Table 2, it was found that the ethanol extract of beluntas leaves had an inhibitory and killing effect on the growth of *Streptococcus sanguinis* bacteria and there was MIC at a concentration of 3.9 g/mL with 99% killing power and MBC was found at a concentration of 7.8 g/mL. mL with 100% killing power. The positive control group of chlorhexidine gluconate at a concentration of 0.2% had an inhibitory and killing effect on *Streptococcus sanguinis* bacteria with an average killing power of 100%. Therefore, it can be said that the ethanol extract of beluntas leaves at a concentration of 7.8 g/mL had the same antibacterial effect as 0.2% chlorhexidine gluconate. The statistical test value obtained P-value 0.00 which states that the MIC and MBC values have a significant effect on the growth of *Streptococcus sanguinis* bacteria.

These results are in accordance with previous research which stated that the ethanol extract of beluntas leaves (*Pluchea indica* L) could inhibit the growth of bacteria, *Staphylococcus aureus*, *Bacillus subtilis* and *Pseudomonas aeruginosa*.^{16,17} Based on the results of qualitative phytochemical tests, active compounds contained in beluntas leaves such as phenols, tannins, flavonoids, saponins, triterpenoids, and alkaloids. Phenol compounds have antibacterial properties due to the presence of hydroxyl and carbonyl groups that can interact with bacterial cells through hydrogen bonds so that it will cause coagulation of proteins and bacterial cell membranes which will cause bacteria to lyse.¹⁸ Phenol can also cause hyperpolarization of the cytoplasmic membrane and increase instability membrane. This causes membrane dysfunction and bacterial cell death.^{14,15}

Tannins are one type of compound that belongs to the polyphenol group and has properties that are soluble in water and organic solvents. Tannins can be used as antibacterial because they have an antiseptic phenol group, so that it can be used as an antibacterial component through hydrogen bonds which will form complex compounds with proteins and will cause bacterial cell proteins to be denatured so that bacterial metabolism is disrupted.^{14,15,19}

Flavonoids also have a role as antibacterial compounds by inhibiting cell membrane function, nucleic acid synthesis, and inhibiting energy metabolism. The role in inhibiting the synthesis of nucleic acids occurs through the accumulation of nucleic acid bases thereby inhibiting the formation of DNA and RNA because rings A and B of flavonoid compounds play an important role in the process of intercalation or hydrogen bonding. The result of flavonoid interaction will also cause damage to the permeability of the bacterial cell wall. Flavonoids will also form extracellular complex compounds and soluble proteins that cause damage to cell membranes and the release of intracellular compounds, thus causing cell function to be inhibited. In addition, energy metabolism will also be inhibited due to inhibition of oxygen use by bacteria. This can occur by preventing the formation of energy in the cytoplasmic membrane and inhibiting the motility of bacteria which play a role in antimicrobial activity and extracellular proteins.^{20,21}

Other components, namely saponins, have broad biological activity and act as antibacterial and antifungal agents. Saponins can cause leakage of bacterial cell membranes, resulting in damage to membrane permeability which can interfere with the survival of bacteria. When the stability of the bacterial cell membrane is disturbed, it will cause the cytoplasm to come out of the cell. This occurs when saponins bind to the cytoplasmic membrane through the outer membrane which will cause the stability of the bacterial cell membrane to be disturbed and result in bacterial cell lysis and then death.^{21,22}

Triterpenoid compounds are also reported to have antibacterial activity. Triterpenoids will react with porins (transmembrane proteins) on the outer wall of the bacterial cell membrane causing the formation of strong polymer bonds. This condition will cause damage to the porin which is the entrance and exit of the compound. Furthermore, this causes the permeability of the bacterial cell wall to decrease, the bacterial cell to lysis, and the death of the bacterial cell.^{14,15,23,24}

Alkaloids also have antibacterial properties that cause bacterial cell death because the cell wall layer is not completely formed by interfering with the peptidoglycan constituent components of bacterial cells by interfering with the formation of cross-bridges of peptidoglycan constituent components in bacterial cells. Alkaloids are also able to inhibit enzymes that play a role in the DNA replication process, thereby inhibiting bacterial growth because bacteria cannot divide by inhibiting DNA replication.^{25,26}

Because beluntas leaves have an active biological content that acts as an antibacterial, the ethanol extract of beluntas leaves can be developed and used as natural antibacterial ingredients because there is a significant value at a concentration of 7.8 g/mL with the number of colonies of *Streptococcus sanguinis* bacteria 0 CFU/mL.

Conclusion

The ethanol extract of beluntas leaves (*Pluchea indica* L) has potential as an antibacterial against *Streptococcus sanguinis*.

References

1. Amtha R, Marcia M, Aninda AI. Plester sariawan efektif dalam mempercepat penyembuhan stomatitis aftosa rekuren dan ulkus traumatikus. Yogyakarta: Majalah Kedokteran Gigi Indonesia. 2017;3(2):7.
2. Apriasari ML, Tuti, H. Stomatitis aftosa rekuren oleh karena anemia. J Dentofasial. 2010; 9(1): 39-46.
3. Langlais RL, Miller CS. Kelainan Rongga Mulut yang Lazim. Jakarta: Hipokrates. 2012. p. 1.
4. Odell E. Cawson's Essentials of Oral Pathology and Oral Medicine. New York: Elsevier. 2017;9(1):220-221.
5. Ghom AG. Textbook of Oral Medicine. New Delhi: Jaypee Brothers Medical Publishers; 2014: 360.
6. Widyastutik O, Permadi A. Penatalaksanaan Nyeri pada Stomatitis Aftosa Rekuren. Jakarta: Journal FKG UNEJ. 2011; 4(3): 396-397.
7. Dutt P, Rathore PK, Khurana D. Chlorhexidine- An antiseptic in periodontics. IOSR-JDMS. 2014;13(9): 85-8.
8. Sari LORK. Pemanfaatan Obat Tradisional Dengan Pertimbangan Manfaat Dan Keamanannya. Depok: Journal Pharmaceutical Sciences and Research. 2016; 3(1): 1-7.
9. Saparinto C, Susiana R. Grow Your Own Medical Plant. Yogyakarta: Lily publisher. 2016: 456-473.
10. Amilah S, Ajiningrum PS. Uji Efektivitas Daya Hambat Sari Daun Pegagan (*Centella asiatica*) dan Daun Beluntas (*Pluchea indica* Less) terhadap Pertumbuhan *Mycobacterium Tuberculosis*. STIGMA Journal science. 2015; 8(2): 6-11.
11. Muta'ali R, Kristanti IP. Pengaruh Ekstrak Daun Beluntas (*Pluchea indica*) terhadap Mortalitas dan Perkembangan Larva Spodoptera litura F. Sumatera: Journal Sains dan seni ITS. 2015; 4(2): 6-7.
12. Arishandi Y, Andriani Y. Tanaman Obat Plus Pengobatan Alternatif. Jakarta: Setia Kawan. 2010.
13. Silalahi M. Keanekaragaman tumbuhan bermanfaat di pekarangan oleh Etnis Sunda Di Desa Sindang Jaya Kabupaten Cianjur Jawa Barat. Jurnal Pendidikan Matematika Dan IPA. 2019; 10(1): 88-104.
14. Raharjo I, Horsten S. *Pluchea indica* in: Plant Resources of South East Asia. Bogor Indonesia: JLCH van Valkenburg and N Bunyapraphatsara. 2002; 12(2): 441-443.
15. Gayatri DAAM, Ernawati DK, Widhiartini IAA. Uji Efektifitas Ekstrak Etanol Daun Beluntas Terhadap Pertumbuhan Bakteri *Streptococcus pyogenes* ATCC 19615 secara in vitro. Bali: Jurnal Medika Udayana. 2020; 10(1): 7-11.
16. Nahak MM, Tedjasulaksana R, Dharmawati IGGA. Khasiat Ekstrak Daun Beluntas untuk Menurunkan Jumlah Bakteri pada Saliva. Journal Interdental Kedokteran gigi. 2017; 5(3): 139-42
17. Manu RRS. Aktivitas Antibakteri Ekstrak Etanol Daun Beluntas (*Pluchea Indica* L.) Terhadap *Staphylococcus Aureus*, *Bacillus Subtilis* Dan *Pseudomonas Aeruginosa*. Surabaya: Jurnal Ilmiah Mahasiswa Universitas Surabaya. 2013; 2(1): 1-10.
18. Ndruru ESC. Perbandingan Efektivitas Berkumur Larutan Probiotik dan Klorheksidin Terhadap Akumulasi Plak dan Jumlah *Streptococcus Mutans* pada Anak Usia 12-15 Tahun Di Yayasan Sos Children's Village. Medan: Fakultas Kedokteran Gigi Universitas Sumatera Utara. 2021.
19. Utomo SB. Uji Aktivitas Antibakteri Senyawa C-4-Metoksifenil kaliks Resorsinarena Termodifikasi Hexadecyltrimethylammonium -Bromide Terhadap Bakteri *Staphylococcus Aureus* dan *Escherichia Coli*. Surakarta: Journal UNS. 2018.
20. Nomer NM, Duniaji AS, Nocianitri KA. Kandungan senyawa flavonoid dan antosianin ekstrak kayu secang (*caesalpinia sappan* L.) serta aktivitas antibakteri terhadap *vibrio cholerae*. Bali: Journal Ilmu dan Teknologi Pangan. 2019; 8(2): 216-225.
21. Haqi HD. Aktivitas Antibakteri Ekstrak Metanol Serbuk Biji Kluwih (*Artocarpus Communis* J.R. & G) Terhadap Pertumbuhan Methicillin Resistant *Staphylococcus Aureus* (MRSA). Yogyakarta: Fakultas Kedokteran Gigi Universitas Muhammadiyah. 2018.
22. Pincus. Clinical Diagnosis and Management by Laboratory Methods. New York: Elsevier. 2011.

23. Makkar HPS, Becker K. Do Tannins In Leaves Of Trees And Shrubs From African And Himalayan Regions Differ In Level And Activity Agroforestry System. Heidelberg: Agroforestry Systems. 2003.
24. Retnowati Y, Bialangi N, Posangi NW. Pertumbuhan Bakteri Staphylococcus aureus Pada Media Yang Diekspos Dengan Infus Daun Sambiloto (*Andrographis paniculata*). Gorontalo: Jurnal Saintek. 2011; 6(2).
25. Taufiq S, Umi Y, Siti, H. Uji Aktivitas Antibakteri Ekstrak Etanol Biji Buah Pepaya (*Carica papaya L.*) terhadap *Escherichia coli* dan *Salmonella typhi*. Bandung: Prosiding Penelitian Spesia Unisba. 2015.
26. Eranawati & Sari K. Kandungan senyawa kimia dan aktivitas antibakteri ekstrak kulit buah alpukat (*Persea americana P.Mill*) terhadap bakteri *Vibrio alginolyticus*. Jurnal Kajian Veteriner. 2015; 3(2): 203-211.

Bukti melakukan review yang pertama (17 Juli 2022)

Peer Review	
Round 1	
Review Version	26973-87879-1-RV.DOCX 2022-07-17
Initiated	—
Last modified	—
Uploaded file	None

Bukti konfirmasi submit revisi pertama yang telah direvisi (Juli 2022)

Antibacterial Potential Ethanol Extract of Beluntas Leaves (*Pluchea indica L*) to *Streptococcus sanguinis*

Wilma Adiwijaya¹, Riani Setiadhi², Vinna Kurniawati Sugiaman^{3*}

¹Fakultas Kedokteran Gigi, Universitas Kristen Maranatha, Indonesia

²Departemen Ilmu Penyakit Mulut, Fakultas Kedokteran Gigi, Universitas Kristen Maranatha, Indonesia

³Departemen Oral Biologi, Fakultas Kedokteran Gigi, Universitas Kristen Maranatha, Indonesia

* Correspondence: vinnakurniawati@yahoo.co.id

ABSTRACT

Introduction: Recurrent aphthous stomatitis is one of the most common oral diseases in the community with a prevalence of 5-66%, the exact cause is unknown but one of the predisposing factors is *Streptococcus sanguinis*. Treatment for RAS has been symptomatic and supportive, including antiseptic mouthwash such as chlorhexidine gluconate 0.2% or topical corticosteroids (triamcinolone acetonide 0.1% in Orabase), however these drugs have some side effects. Treatment using herbal ingredients such as beluntas leaves can cure RAS with relatively low

prices and minimal side effects. Beluntas leaves (*Pluchea Indica L*) contain active compounds such as phenols, tannins, flavonoids, saponins, triterpenoids, essential oils, terpenoids, and alkaloids. Many compounds of this group are known to have antibacterial activity. Methods: This study aimed to determine the minimum inhibitory level (MIC) and minimum killing rate (MBC) of 96% ethanol extract of beluntas leaves on the growth of *Streptococcus sanguinis*. MIC was measured by broth microdilution technique with DMSO solvent 10% and 8 concentrations of beluntas extract. Chlorhexidine gluconate 0.2% was used as a positive control for the comparison compound. Furthermore, the MBC test was carried out using the total plate count method for treatments that gave the MIC value. One Way Anova analysis with Post Hoc Tukey was used to determine the significant difference between treatments. Results: The ethanol extract of beluntas leaves (*Pluchea indica L*) has a Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) for the growth of *Streptococcus sanguinis* with a minimum inhibitory concentration of 3.95 g/mL and a minimum concentration of 7.8 g/mL. Conclusion: The ethanol extract of beluntas leaves (*Pluchea indica L*) has potential as an antibacterial against *Streptococcus sanguinis*.

Keywords: Recurrent aphthous stomatitis, *Streptococcus sanguinis*, ethanol extract of beluntas leaves, Minimum Inhibitory Concentration, Minimum Bactericidal Concentration.

Introduction

Recurrent aphthous stomatitis (RAS) is a problem that often arises in dentistry and is one of the most common oral diseases in the community with a prevalence of 66%. RAS is characterized by recurrent ulcers on the patient's oral mucosa and can be painful. The prevalence of oral ulceration worldwide is known to be as much as 4%, where RAS is the disease with the greatest prevalence of 25% and is experienced by most women in the second and third decades.^{1,2}

The clinical manifestations of RAS are ulcers, single or multiple, shallow, oval, and pain.³ Prodromal symptoms appear before the onset of RAS including discomfort and redness for 1-3 days.⁴ The exact etiology of RAS has not been determined so the research focused on the predisposing factors for RAS. Several factors are considered as predisposing factors for RAS, including immunological abnormalities, genetics, systemic factors, endocrine system, stress, smoking cessation, allergic factors, and microorganisms. The main microorganisms associated with the formation of RAS are pleomorphic transitional L α -hemolytic *Streptococcus* and *Streptococcus sanguinis*.⁵

One of the efforts to manage RAS is by symptomatic and supportive treatment because until now the etiology of RAS is still unknown. Therefore, treatment is only aimed at curing complaints. The goal of symptomatic treatment is to reduce symptoms, reduce the number and size of ulcers. Currently used drugs include: antiseptic mouthwash (chlorhexidine gluconate 0.2%) to reduce the duration and discomfort of RAS or topical corticosteroids (triamcinolone acetonide 0.1% in Orabase).⁶ Chlorhexidine gluconate 0.2% is the gold standard, which is effective against both gram negative and positive and facultative aerobes and anaerobes. However, its use in more than 2 weeks has side effects, including causing burning of the oral mucosa, impaired taste, tooth staining, erosion of the oral surface, and dryness of the oral cavity.⁷

Therefore, it is necessary to develop an alternative therapy by utilizing herbal plants in curing ulcers in RAS. This is intended so that patients can recover without side effects at a relatively low cost.⁸ The development of a back to nature lifestyle has made the use of herbal plants increasingly widespread by the community as alternative materials for the treatment or prevention of certain diseases. The community considers that herbal ingredients are easier to obtain, the price is more affordable, and the side effects are minimal, so it is safer for consumption.⁹ One of the natural ingredients that can be used as traditional medicine is the beluntas plant.¹⁰ The leaves, flowers, and roots of beluntas can be used for therapy, but the beluntas leaves are the part that has the highest biologically active component.¹¹

Beluntas (*Pluchea Indica* L) is a wild plant in dry areas on hard and rocky soils or grown as a hedge.¹² This plant has a distinctive aromatic odor and bitter taste. The part used from this plant is the leaves and roots which are efficacious for eliminating body odor and bad breath, increasing appetite, overcoming digestive disorders in children, relieving pain in rheumatism, and so on. The active compounds contained in beluntas leaves are flavonoids, triterpenoids, phenols, and essential oil derivatives.¹³ Beluntas leaves have antibacterial properties due to the content of phenolics, triterpenoids, and tannins that can cause bacterial cell death.^{14,15}

Based on the research that has been done, it shows that the ethanol extract of the leaves of beluntas (*Pluchea indica* L) can inhibit the growth of *Streptococcus mutans*, *Staphylococcus aureus*, *Bacillus subtilis*, and *Pseudomonas aeruginosa* bacteria. Based on the explanation above and there is no research on *Streptococcus sanguinis* bacteria, so the authors are interested in research to determine the antibacterial effect of beluntas leaves through testing the Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) of the ethanol extract of the beluntas leaves on the growth of *Streptococcus sanguinis* which is one of the bacteria that causes recurrent aphthous stomatitis.

METHOD

Determination and Ethanol Extract of Beluntas Leaves

Beluntas leaves were obtained from the Subang Tropical Balitbu plantation which were then determined at the Biology Laboratory of Padjadjaran University. Extract beluntas leaves with 96% ethanol and let stand for 3 days, then the extract is filtered using filter paper to obtain the filtrate. Then the filtrate was evaporated using a rotary evaporator at a temperature of 70°C until the consistency was like a paste.

The ethanol extract of beluntas leaves was put into a 15mL tube as much as 0.5 grams and dissolved in 10% DMSO to reach a concentration of 500 g/mL. The extract was diluted using a multichannel Micropipette so that the final concentrations were 250 g/ml, 125 g/ml, 62.5 g/ml, 31.3 g/ml, 15.6 g/ml, 7.8 g/ml, 3.95 g/ml, and 1.95 g/ml.

Preparation of Streptococcus sanguinis inoculum

The bacteria that will be used in this study is Streptococcus sanguinis ATCC 10556. Take 2 oses and suspension in BHI-B, incubated for 24 hours. Then standardized with 0.5 McFarland solution so that the total density of bacteria used is 1.5×10^8 CFU/ml. Check using a spectrophotometer with a wavelength of 600 nm, then diluted to 105.

Minimum Inhibitory Level (MIC) and Minimum Killing Rate (KBM)

The method used in this study was following the method developed by CLSI (Clinical Laboratory Standard Institute) with a slight modification, namely adding 2% sucrose to BHI-A and BH-B media.

This study consisted of ten treatments, namely eight treatments of ethanol extract of beluntas leaves with concentrations of 250 g/ml, 125 g/ml, 62.5 g/ml, 31.3 g/ml, 15.6 g/ml, 7,8 g/ml, 3.95 g/ml, and 1.95 g/ml; positive control (chlorhexidine gluconate 0.2%); and negative control (DMSO).

200 L/mL of BHI-Broth media was added to all 96 well plates and added 200 l of ethanol extract of beluntas leaves of each concentration and 10 l of bacterial suspension, with the format of media + sample (negative control), media + solvent (solvent control), media + sample + bacteria (test sample), media + solvent + bacteria (positive control). Incubate at 37°C for 16-20 hours. Insert the well plate into a spectrophotometer with a wavelength of 600 nm to see the minimum inhibitory concentration using the absorbance method.

Take 200 L of 0.2% Chlorhexidine gluconate and put it into the well plate with 3 repetitions, 10 l of bacterial suspension 105 for 2 repetitions and 1 repetition plate without bacteria as a positive control. Incubate for 24 hours at 37oC with 5% CO2.

Well plates indicated as MIC and KBM were inoculated into sterile BHI-Agar medium in petri dishes and incubated at 37oC with 5% CO2 for 24 hours, calculating the number of colonies in each petri dish from each treatment with various concentrations of extract using a colony counter.

The number of colonies contained in BHI-Agar media was used to determine MIC and MBC was calculated using a Colony counter. The calculation of the number of colonies formed is called the total plate count (TPC) with colony forming units/mL (CFU/mL) calculated by the formula:

$$\text{TPC value (CFU/ml)} = \frac{\text{Number of colonies formed} \times \text{Dilution factor}}{\text{The volume of bacterial colonies that is included in the criteria}}$$

The calculation of the percentage of bactericidal power is measured by the formula:

$$\% \text{ Kill Efficiency} = \frac{\text{TPC bacteria control (CFU/ml)} - \text{TPC treatment (CFU/ml)} \times 100\%}{\text{Mean bacterial colony control}}$$

RESULTS

The results of the determination of Beluntas plants carried out at the Biology Laboratory of Padjadjaran University, Jatinangor, West Java showed that the sample used was Beluntas (*Pluchea Indica* L).

Phytochemical tests were carried out at the Central Laboratory of Padjadjaran University, Jatinangor, West Java. Table 4.1 shows the results of qualitative phytochemical tests taken from the ethanol extract of beluntas leaves.

Table 4.1 Qualitative Phytochemical Test Results

No	Secunder Metabolite	Methods	Result s
1	Fenolik	Reagent FeCl ₃ 5%	++
2	Tanin	Reagent FeCl ₃ 1%	++
3	Flavonoid	a. Reagent HCL concentrated + Mg b. Reagent H ₂ SO ₄ 2N c. Reagent NaOH 10	- - +
4	Saponin	Heated	+
5	Triterpenoid	Reagent H ₂ SO ₄ concentrated + CH ₃ COOH	++
	dan Steroid	anhydrous	-
6	Alkaloid	Reagent Dragendorff	+

Description:

+ : Little

++ : Medium

+++ : Much

- : None

Extract Concentration	Media control	Average Absorbance Bacteria+ Sample	Difference	Note(s)
250 µg/mL	2,039	2,018	0,021	
125 µg/mL	1,4203	1,3964	0,023	
62.5 µg/mL	0,8977	0,8568	0,04	
31.25 µg/mL	0,5208	0,4938	0,27	
15.62µg/ mL	0,312	0,2874	0,25	
7.8 µg/mL	0,1873	0,1789	0,84	
3.9 µg/mL	0,1245	0,114	0,1	KHM
1.95 µg/mL	0,0795	0,0833	-0,0035	

The results of the positive control test (chlorhexidine gluconate 0.2%) obtained the average number of colonies of *Streptococcus sanguinis* bacteria with 3 repetitions after 24 hours of incubation was 0 CFU/mL with an average bactericidal power of 100%.

Minimum Bactericidal Concentration (MBC)

The results of the MBC test showed that at a concentration of 7.8 g/mL the ethanol extract of the tested beluntas leaves was not observed for the growth of *Streptococcus sanguinis*. To count the number of bacteria at each test concentration, the number of bacteria was calculated using the total plate count technique with a colony counter as shown in Figure 2 and Table 2.



Figure 2. The results of the calculation of the number of bacteria using the Colony Counter

Notes: (A) Number of *Streptococcus sanguinis* Bacteria Colonies at the Concentration of Ethanol Extract of Beluntas Leaves 7.8 g/mL (KBM). (B) Number of *Streptococcus sanguinis* Bacteria Colonies at Concentration of Ethanol Extract of Beluntas Leaves 3.9 g/mL (MIC)

Table 2. Number of *Streptococcus sanguinis* Bacteria Colonies After 24 hours incubation at various concentrations of Beluntas Leaf Ethanol Extract

Extract Concentration	Mean Colony (CFU/mL)	Mean kill Efficiency	Note(s)
250 µg/mL (Kelompok A)	0	100%	
125 µg/mL (Kelompok B)	0	100%	
62.5 µg/mL (Kelompok C)	0	100%	
31.25 µg/mL (Kelompok D)	0	100%	
15.62 µg/ mL (Kelompok E)	0	100%	
7.8 µg/mL (Kelompok F)	0	100%	KBM
3.9 µg/mL (Kelompok G)	12,9x 10 ¹	99%(0,9999612)	KHM
1.95 µg/mL (Kelompok H)	27,3 x 10 ⁵	99%(0,9999178)	

Colony Number Normality Test Results

The results of the normality test using Shapiro-Wilk showed that there were 2 variables tested, namely concentrations of 3.95 g/mL and 1.95 g/mL. It is known that the significance value is 0.147 and 0.114, where the value is greater than with a value of 0.05, thus it can be concluded that the data is normally distributed.

One Way ANOVA Test Results

The results of the One Way ANOVA test to determine whether the data has an average difference or cannot be known from the significance value of the number of colonies then compared with the significance value, with a test decision, if the value (Sig) > 0.05 then there is no difference in average, if value (Sig) < 0.05 then there is an average difference. The results of the ANOVA test above show the F value for the number of colonies of 22.882 and the significance (p-value) of 0.000 is less than 0.05, so it can be concluded that at concentrations of 3.95 and 1.95 there was a different effect on the number of colonies.

Tukey's Post Hoc Test Results

Based on the results of the Tukey Post Hoc test in appendix 3, the number of colonies at a concentration of 250 was the same as a concentration of 125, 62.5, 31.3, 15.6, 7.8, and 1.95 because they were in the same subset. Meanwhile, the number of colonies at a concentration of 3.95 was not the same as a concentration of 125, 62.5, 31.3, 15.6, 7.8, and 1.95 because they were indifferent subsets. The comparison results can be seen through the mean difference column, obtained at a concentration of 3.95 significantly different ($p < 0.05$).

DISCUSSION

Based on the research results presented in Table 2, it was found that the ethanol extract of beluntas leaves had an inhibitory and killing effect on the growth of *Streptococcus sanguinis* bacteria and there was MIC at a concentration of 3.9 g/mL with 99% killing power and MBC was found at a concentration of 7.8 g/mL. mL with 100% killing power. The positive control group of chlorhexidine gluconate at a concentration of 0.2% had an inhibitory and killing effect on *Streptococcus sanguinis* bacteria with an average killing power of 100%. Therefore, it can be said that the ethanol extract of beluntas leaves at a concentration of 7.8 g/mL had the same antibacterial effect as 0.2% chlorhexidine gluconate. The statistical test value obtained P-value 0.00 which states that the MIC and MBC values have a significant effect on the growth of *Streptococcus sanguinis* bacteria.

These results are in accordance with previous research which stated that the ethanol extract of beluntas leaves (*Pluchea indica* L) could inhibit the growth of bacteria, *Staphylococcus aureus*, *Bacillus subtilis* and *Pseudomonas aeruginosa*.¹⁷

Based on the results of qualitative phytochemical tests, beluntas leaves contain active compounds such as phenols, tannins, flavonoids, saponins, triterpenoids, and alkaloids. Phenol compounds have antibacterial properties due to the presence of hydroxyl and carbonyl groups that can interact with bacterial cells through hydrogen bonds so that it will cause coagulation of proteins and bacterial cell membranes which will cause bacteria to lyse.¹⁸ Phenol can also cause hyperpolarization of the cytoplasmic membrane and increase instability membrane. This causes membrane dysfunction and bacterial cell death

Tannins are one type of compound that belongs to the polyphenol group and has properties that are soluble in water and organic solvents. Tannins can be used as antibacterial because they have an antiseptic phenol group, so that it can be used as an antibacterial component through hydrogen bonds which will form complex compounds with proteins and will cause bacterial cell proteins to be denatured so that bacterial metabolism is disrupted.

The mechanism of action of flavonoids as antibacterial compounds is divided into 3, namely: inhibiting nucleic acid synthesis, inhibiting cell membrane function, and inhibiting energy metabolism. The role in inhibiting the synthesis of nucleic acids occurs through the accumulation of nucleic acid bases thereby inhibiting the formation of DNA and RNA because rings A and B of flavonoid compounds play an important role in the process of intercalation or hydrogen bonding. The result of flavonoid interaction will also cause damage to the permeability of the bacterial cell wall. Its role in inhibiting the function of bacterial cell membranes is that flavonoids will form complex compounds from extracellular and dissolved proteins so that the cell membrane will be damaged and intracellular compounds will come out. Another role is to inhibit energy metabolism by inhibiting the use of oxygen by bacteria. This happens by preventing the formation of energy in the cytoplasmic membrane and inhibiting the motility of bacteria that play a role in antimicrobial activity and extracellular proteins.

Other components, namely saponins, have broad biological activity and act as antibacterial and antifungal agents. Saponins can cause leakage of bacterial cell membranes, resulting in damage to membrane permeability which can interfere with the survival of bacteria. When the stability of the bacterial cell membrane is disturbed, it will cause the cytoplasm to come out of the cell. This occurs when saponins bind to the cytoplasmic membrane through the outer membrane which will cause the stability of the bacterial cell membrane to be disturbed and result in bacterial cell lysis and then death.

Triterpenoid compounds are also reported to have antibacterial activity. Triterpenoids will react with porins (transmembrane proteins) on the outer wall of the bacterial cell membrane causing the formation of strong polymer bonds. This condition will cause damage to the porin which is the entrance and exit of the compound. Furthermore, this causes the permeability of the bacterial cell wall to decrease, the bacterial cell to lysis, and the death of the bacterial cell. ^{14,15,23,24}

Alkaloids also have antibacterial properties that cause bacterial cell death because the cell wall layer is not completely formed by interfering with the peptidoglycan constituent components of bacterial cells by interfering with the formation of cross-bridges of peptidoglycan constituent components in bacterial cells. Alkaloids are also able to inhibit enzymes that play a role in the DNA replication process, thereby inhibiting bacterial growth because bacteria cannot divide by inhibiting DNA replication.^{25,26}

Because beluntas leaves have an active biological content that acts as an antibacterial, the ethanol extract of beluntas leaves can be developed and used as natural antibacterial ingredients because there is a significant value at a concentration of 7.8 g/mL with the number of colonies of *Streptococcus sanguinis* bacteria 0 CFU/mL.

CONCLUSION

The ethanol extract of beluntas leaves (*Pluchea indica* L) has potential as an antibacterial against *Streptococcus sanguinis*.

Conflict of interest.....

BIBLIOGRAPHY

27. Amtha R, Marcia M, Aninda AI. Plester sariawan efektif dalam mempercepat penyembuhan stomatitis aftosa rekuren dan ulkus traumatikus. Yogyakarta: Majalah Kedokteran Gigi Indonesia. 2017;3(2):7.
28. Apriasari ML, Tuti, H. Stomatitis aftosa rekuren oleh karena anemia. J Dentofasial. 2010; 9(1): 39-46.
29. Langlais RL, Miller CS. Kelainan Rongga Mulut yang Lazim. Jakarta: Hipokrates. 2012. p. 1.
30. Odell E. Cawson's Essentials of Oral Pathology and Oral Medicine. New York: Elsevier. 2017;9(1):220-221.
31. Ghom AG. Textbook of Oral Medicine. New Delhi: Jaypee Brothers Medical Publishers; 2014: 360.
32. Widyastutik O, Permadi A. Penatalaksanaan Nyeri pada Stomatitis Aftosa Rekuren. Jakarta: Journal FKG UNEJ. 2011; 4(3): 396-397.
33. Dutt P, Rathore PK, Khurana D. Chlorhexidine- An antiseptic in periodontics. IOSR-JDMS. 2014;13(9): 85-8.
34. Sari LORK. Pemanfaatan Obat Tradisional Dengan Pertimbangan Manfaat Dan Keamanannya. Depok: Journal Pharmaceutical Sciences and Research. 2016; 3(1): 1-7.
35. Saparinto C, Susiana R. Grow Your Own Medical Plant. Yogyakarta: Lily publisher. 2016: 456-473.
36. Amilah S, Ajiningrum PS. Uji Efektivitas Daya Hambat Sari Daun Pegagan (*Centella asiatica*) dan Daun Beluntas (*Plunchea indica* Less) terhadap Pertumbuhan *Mycobacterium Tuberculosis*. STIGMA Journal science. 2015; 8(2): 6-11.
37. Muta'ali R, Kristanti IP. Pengaruh Ekstrak Daun Beluntas (*Pluchea indica*) terhadap Mortalitas dan Perkembangan Larva Spodoptera litura F. Sumatera: Journal Sains dan seni ITS. 2015; 4(2): 6-7.
38. Arishandi Y, Andriani Y. Tanaman Obat Plus Pengobatan Alternatif. Jakarta: Setia Kawan. 2010.
39. Silalahi M. Keanekaragaman tumbuhan bermanfaat di pekarangan oleh Etnis Sunda Di Desa Sindang Jaya Kabupaten Cianjur Jawa Barat. Jurnal Pendidikan Matematika Dan IPA. 2019; 10(1): 88-104.
40. Raharjo I, Horsten S. *Pluchea indica* in: Plant Resources of South East Asia. Bogor Indonesia: JLCH van Valkenburg and N Bunyapraphatsara. 2002; 12(2): 441-443.
41. Gayatri DAAM, Ernawati DK, Widhiartini IAA. Uji Efektifitas Ekstrak Etanol Daun Beluntas Terhadap Pertumbuhan Bakteri *Streptococcus pyogenes* ATCC 19615 secara in vitro. Bali: Jurnal Medika Udayana. 2020; 10(1): 7-11.
42. Nahak MM, Tedjasulaksana R, Dharmawati IGGA. Khasiat Ekstrak Daun Beluntas untuk Menurunkan Jumlah Bakteri pada Saliva. Journal Interdental Kedokteran gigi. 2017; 5(3): 139-42
43. Manu RRS. Aktivitas Antibakteri Ekstrak Etanol Daun Beluntas (*Pluchea Indica* L.) Terhadap *Staphylococcus Aureus*, *Bacillus Subtilis* Dan *Pseudomonas Aeruginosa*. Surabaya: Jurnal Ilmiah Mahasiswa Universitas Surabaya. 2013; 2(1): 1-10.

44. Ndruru ESC. Perbandingan Efektivitas Berkumur Larutan Probiotik dan Klorheksidin Terhadap Akumulasi Plak dan Jumlah Streptococcus Mutans pada Anak Usia 12-15 Tahun Di Yayasan Sos Children's Village. Medan: Fakultas Kedokteran Gigi Universitas Sumatera Utara. 2021.
45. Utomo SB. Uji Aktivitas Antibakteri Senyawa C-4-Metoksifenil kaliks Resorsinarena Termodifikasi Hexadecyltrimethylammonium -Bromide Terhadap Bakteri Staphylococcus Aureus dan Escherichia Coli. Surakarta: Journal UNS. 2018.
46. Nomer NM, Duniaji AS, Nocianitri KA. Kandungan senyawa flavonoid dan antosianin ekstrak kayu secang (*caesalpinia sappan* L.) serta aktivitas antibakteri terhadap *vibrio cholerae*. Bali: Journal Ilmu dan Teknologi Pangan. 2019; 8(2): 216-225.
47. Haqi HD. Aktivitas Antibakteri Ekstrak Metanol Serbuk Biji Kluwih (*Artocarpus Communis* J.R. & G) Terhadap Pertumbuhan Methicillin Resistant *Staphylococcus Aureus* (MRSA). Yogyakarta: Fakultas Kedokteran Gigi Universitas Muhammadiyah. 2018.
48. Pincus. *Clinical Diagnosis and Management by Laboratory Methods*. New York: Elsevier. 2011.
49. Makkar HPS, Becker K. Do Tannins In Leaves Of Trees And Shrubs From African And Himalayan Regions Differ In Level And Activity Agroforestry System. Heidelberg: Agroforestry Systems. 2003.
50. Retnowati Y, Bialangi N, Posangi NW. Pertumbuhan Bakteri *Staphylococcus aureus* Pada Media Yang Diekspos Dengan Infus Daun Sambiloto (*Andrographis paniculata*). Gorontalo: Jurnal Saintek. 2011; 6(2).
51. Taufiq S, Umi Y, Siti, H. Uji Aktivitas Antibakteri Ekstrak Etanol Biji Buah Pepaya (*Carica papaya* L.) terhadap *Escherichia coli* dan *Salmonella typhi*. Bandung: Prosiding Penelitian Spesia Unisba. 2015.
52. Eranawati & Sari K. Kandungan senyawa kimia dan aktivitas antibakteri ekstrak kulit buah alpukat (*Persea americana* P.Mill) terhadap bakteri *Vibrio alginolyticus*. Jurnal Kajian Veteriner. 2015; 3(2): 203-211.

Bukti melakukan review yang kedua

Bukti konfirmasi submit artikel yang telah revisi kedua

Bukti konfirmasi artikel diterima



Bukti Galery Proof Manuscript



Bukti Publikasi Online Artikel (Desember 2022)

Google Cendekia Antibacterial Potential Ethanol Extract of Beluntas Leaves (Pluchea indica L.)

Artikel

Profil saya Koleksi

iporj.usk.ac.id

Antibacterial potential ethanol extract of beluntas leaves (Pluchea indica L.) to Streptococcus sanguinis

W ADIMAWATI, E SETIADI, V K SUGAMAWATI

Journal of Stomatognathal Dentistry Society, 2022 - jstomat.usk.ac.id

Abstract

Recurrent aphthous stomatitis (RAS) is one of the most common oral diseases in the community, with a prevalence of 5-66%, with one of the predisposing factors being Streptococcus sanguinis. Treatment for RAS has been symptomatic and supportive, including antiseptic mouthwash such as chlorhexidine gluconate 0.2% or topical corticosteroids (flamcinnone acetonide 0.1% in Orabase). However, these drugs have some side effects. Treating herbal ingredients such as Beluntas leaves (Pluchea indica)

TAMPLIKAN LEBIH BANYAK ~

☆ Simpan ID KUP Ditinjau 2 kali Artikel terkait 2 versi 36

Melampirkan hasil terbaik untuk penyelesaian ini. Lihat semua hasil

Filter saja
Sejak 2025
Sejak 2024
Sejak 2021
Rentang khusus...

Urutkan menurut relevansi
Urutkan menurut tanggal

Sembunyikan
Artikel kapian

sertakan poster
 mencakup kutipan