

DAFTAR PUSTAKA

- 1 Kumar V, Abbas AK, Aster JC. *Robbin Basic Pathology*. 2012; 9: 739-751
- 2 Soelistijo SA, Novida H, Rudijanto A, Soewondo P, Suastika K, Manaf A et al. *Konsensus Pengendalian dan Pencegahan Diabetes Melitus Tipe 2 di Indonesia 2015*. 2015 : 1-75
- 3 Federation ID. *IDF Diabetes Atlas Ninth edition 2019*. 2019 doi:10.1289/image.ehp.v119.i03.
- 4 Kemenkes RI. *Laporan Nasional RISKESDAS 2018*. 2018.
- 5 Sheweita SA, Mashaly S, Newairy AA, Abdou HM, Eweda SM. Changes in oxidative stress and antioxidant enzyme activities in streptozotocin-induced diabetes mellitus in rats: Role of alhagi maurorum extracts. *Oxid Med Cell Longev* 2016; 3 : 4-7.
- 6 Karaçil Ermumucu MŞ, Şanlıer N. Black Cumin (*Nigella sativa*) And Its Active Component Of Thymoquinone : Effects On Health. *J Food Heal Sci* 2017. 3(4):170-183.
- 7 Meddah B, Ducroc R, El Abbes Faouzi M, Eto B, Mahraoui L, Benhaddou-Andaloussi A et al. *Nigella sativa* inhibits intestinal glucose absorption and improves glucose tolerance in rats. *J Ethnopharmacol* 2009; 121(3):419-24
- 8 Heshmati J, Namazi N. Effects of black seed (*Nigella sativa*) on metabolic parameters in diabetes mellitus: A systematic review. *Complement. Ther. Med*. 2015; 23 (2): 275-82
- 9 Elsner M, Tiedge M, Guldbakke B, Munday R, Lenzen S. Importance of the GLUT2 glucose transporter for pancreatic beta cell toxicity of alloxan. *Diabetologia* 2002; 45:1542-1549.
- 10 Kaatabi H, Bamosa AO, Badar A, Al-Elq A, Abou-Hozaiifa B, Lebda F et al. *Nigella sativa* improves glycemic control and ameliorates oxidative stress in patients with type 2 diabetes mellitus: Placebo controlled participant blinded clinical trial. *PLoS One* 2015 ;10(2)
- 11 Younus H. Molecular and Therapeutic actions of Thymoquinone: Actions of Thymoquinone. 2018:1-15.
- 12 Drake R, Vogl AW, Mitchell AWM, Tibbitts R, Richardson P. *Gray's Atlas of Anatomy: with Student Consult Online Access*. 2014.
- 13 Newell FW. *Sobotta Atlas of Human Anatomy*. 2011; 23:120-127.
- 14 Eroschenko VP. *Atlas Histologi, diFiore*. 2011; 11:324-329.
- 15 Gartner LP, Hiatt JL. *Introduction to Histology and Basic Histological Techniques*. In: *Color Textbook of Histology*. 2007; 3 :417-422.
- 16 Sherwood L. *Human Physiology 8th edition*. 2013: 751-756.
- 17 Khonsary S. *Guyton and Hall: Textbook of Medical Physiology 2017* : 939-950

- 18 IDAI. Diagnosis dan Tata Laksana Diabetes Melitus Tipe-1 pada Anak dan Remaja. Jakarta 2017.
- 19 World Health Organization (WHO). Definition, Diagnosis and classification of diabetes mellitus and its complications. Part 1: Diagnosis and classifications of diabetes mellitus. Geneva: Department of Noncommunicable Disease Surveillance; 1999.
- 20 Ghayur MN, Gilani AH, Janssen LJ. Intestinal, airway, and cardiovascular relaxant activities of thymoquinone. *Evidence-based Complement Altern Med* 2012 : 1-13.
- 21 Hajhashemi V, Ghannadi A, Jafarabadi H. Black Cumin Seed Essential Oil, as a Potent Analgesic and Antiinflammatory Drug. *Phyther. Res.* 2004. doi:10.1002/ptr.1390.
- 22 Nickavar B, Mojab F, Javidnia K, Roodgar Amoli MA. Chemical Composition of the Fixed and Volatile Oils of *Nigella sativa* L. from Iran. *Zeitschrift fur Naturforsch - Sect C J Biosci* 2003; 58(10): 629-31.
- 23 Heshmati J, Namazi N, Memarzadeh MR, Taghizadeh M, Kolahdooz F. *Nigella sativa* oil affects glucose metabolism and lipid concentrations in patients with type 2 diabetes: A randomized, double-blind, placebo-controlled trial. *Food Res Int* 2015: 87-93.
- 24 Elsner M, Gurgul-Convey E, Lenzen S. Relative importance of cellular uptake and reactive oxygen species for the toxicity of alloxan and dialuric acid to insulin-producing cells. *Free Radic Biol Med* 2006; 41 (5): 825-34.
- 25 Lenzen S. The mechanisms of alloxan- and streptozotocin-induced diabetes. *Diabetologia*. 2008.
- 26 Alarcon-Aguilar F, Vega-Avila E, Almanza-Perez J, Velasco-Lezama R, Vazquez-Carrillo L, Roman-Ramos R. Hypoglycemic effect of *Plantago major* seeds in healthy and alloxan-diabetic mice. *Proc West Pharmacol Soc* 2006.