



SURAT TUGAS
No.417/SUT/FKG-UKM/VI/2024

Yang bertanda tangan di bawah ini

Nama : Dr. Ignatius Setiawan, drg., MM.
Jabatan : Dekan Fakultas Kedokteran Gigi Universitas Kristen Maranatha

Dengan ini menugaskan kepada :

No	NIK	Dosen
1	120005	Dr. drg. Vinna Kurniawati Sugiaman, M.Kes., PBO
2	120012	drg. Rudy Djuanda, Sp.KGA

Melaksanakan publikasi jurnal dengan judul: *Effect of Saga Leaf Extract (Abrus precatorius Linn) in Inhibiting Enterococcus faecalis Bacteria Growth as an Alternative Root Canal Irrigation Material* pada Juni 2024 di Publikasi **European Journal of General Dentistry (EJGD)**

Demikian agar tugas ini dilaksanakan dengan sebaik-baiknya.

Bandung, 20 Juni 2024
Dekan Fakultas Kedokteran Gigi
Universitas Kristen Maranatha


Dr. drg. Ignatius Setiawan, MM.

NIK. 120010


European Journal of GENERAL DENTISTRY

Editorial Board:
Prof. Roberto Falavigna



Scopus
CRPPI



Thieme

[eFirst \(/products/ejournals/issue/eFirst/10.1055/s-00051518\)](/products/ejournals/issue/eFirst/10.1055/s-00051518)[Issue \(/products/ejournals/journal/10.1055/s-00051518\)](/products/ejournals/journal/10.1055/s-00051518)[Most Read \(/products/ejournals/topten/10.1055/s-00051518\)](/products/ejournals/topten/10.1055/s-00051518)**Publication Date: 24 July 2024****Original Article**

Bagio, Dini Asrianti; Warganegara, Ibramanto; Maharti, Ike Dwi; Margono, Anggraini; Kusumasari, Citra; Alinda, Sylva Dinie; Wairooy, Valeria Widita:

Assessing Cell Viability: Comparative Analysis of Calcium Hydroxide, Triple Antibiotic Paste, and Their Synergistic Impact on human Dental Pulp Stem Cells (/products/ejournals/abstract/10.1055/s-0044-1788041)

[HTML \(/products/ejournals/html/10.1055/s-0044-1788041\)](/products/ejournals/html/10.1055/s-0044-1788041)[PDF \(3444 kb\) \(/products/ejournals/pdf/10.1055/s-0044-1788041.pdf\)](/products/ejournals/pdf/10.1055/s-0044-1788041.pdf)**Original Article**

Husein, Adam bin; Seoudi, Sana Mhd. Fouad; El-Damanhoury, Hatem Mostafa; Aziz, Ibrahim Mahmood; Abou Neel, Ensanya Ali:

Hybrid Layer, Shear Bond Strength, and Fracture Patterns of Titanium Dioxide-Doped Phosphate Glass-Filled Universal Dental Adhesives (/products/ejournals/abstract/10.1055/s-0044-1787789)

[HTML \(/products/ejournals/html/10.1055/s-0044-1787789\)](/products/ejournals/html/10.1055/s-0044-1787789)[PDF \(1588 kb\) \(/products/ejournals/pdf/10.1055/s-0044-1787789.pdf\)](/products/ejournals/pdf/10.1055/s-0044-1787789.pdf)**Original Article**

Sari, Pitri Ayu Puspita; Sugiaman, Vinna Kurniawati; Djuanda, Rudy:

Effect of Saga Leaf Extract (*Abrus precatorius* Linn) in Inhibiting *Enterococcus faecalis* Bacteria Growth as an Alternative Root Canal Irrigation Material (/products/ejournals/abstract/10.1055/s-0044-1788258)

[HTML \(/products/ejournals/html/10.1055/s-0044-1788258\)](/products/ejournals/html/10.1055/s-0044-1788258)[PDF \(1395 kb\) \(/products/ejournals/pdf/10.1055/s-0044-1788258.pdf\)](/products/ejournals/pdf/10.1055/s-0044-1788258.pdf)**Publication Date: 15 July 2024****Review Article**

Selmani, Mimoza E.; Duci, Shkelzen B.; Gashi, Nora Asani; Bukleta, Manushaqe Selmani:

Orthodontic and Surgical Management of Impacted Maxillary Canines: A Narrative Review (/products/ejournals/abstract/10.1055/s-0044-1786550)

[HTML \(/products/ejournals/html/10.1055/s-0044-1786550\)](/products/ejournals/html/10.1055/s-0044-1786550)[PDF \(3528 kb\) \(/products/ejournals/pdf/10.1055/s-0044-1786550.pdf\)](/products/ejournals/pdf/10.1055/s-0044-1786550.pdf)**Publication Date: 08 July 2024****Perspective**

Khurshid, Zohaib; Alqurashi, Hatim; Ashi, Heba:

Advancing Environmental Sustainability in Dentistry and Oral Health (/products/ejournals/abstract/10.1055/s-0044-1787099)



HTML (/products/ejournals/html/10.1055/s-0044-1787099)
PDF (1274 kb) (/products/ejournals/pdf/10.1055/s-0044-1787099.pdf)

Publication Date: 02 July 2024

Original Article

Suresh, Balaji; Jeevanandan, Ganesh; Ravindran, Vignesh; Vishwanathaiah, Satish; Panda, Suman; Syed, Ather Ahmed; Altowairqi, Sultan; AlShehri, Aram; Alaqla, Ali; Khanagar, Sanjeev; Maganur, Prabhadevi:

Evaluation of Volumetric Changes between Pediatric Rotary Files and Manual Files during Canal Preparation of Primary Mandibular Canine: A Nano-CT Analysis
(/products/ejournals/abstract/10.1055/s-0044-1786548)



HTML (/products/ejournals/html/10.1055/s-0044-1786548)
PDF (517 kb) (/products/ejournals/pdf/10.1055/s-0044-1786548.pdf)

Publication Date: 26 June 2024

Original Article

Abdullah, Syatirah-Najmi; Zaki, Wan Nur Faizatulkmal Wan Ahmad; Mansor, Syarifah Musyirah Qistina Sayed; Al-Kadhim, Aws Hashim Ali; Ghafar, Siti Aisyah Abd; Hanafiah, Rohazila Mohamad:

Antibacterial Activity of Calcium Hydroxide and Zinc Oxide Combined with Several Solutions against *Enterococcus faecalis* Growth (/products/ejournals/abstract/10.1055/s-0044-1786547)



HTML (/products/ejournals/html/10.1055/s-0044-1786547)
PDF (2655 kb) (/products/ejournals/pdf/10.1055/s-0044-1786547.pdf)

Publication Date: 14 May 2024

Original Article

Al-Zubaydi, Fudhla Sadoon; Obaid, Dina H.; Nahidh, Mohammed; Marrapodi, Maria Maddalena; Ronsivalle, Vincenzo; Cicciù, Marco; Minervini, Giuseppe:

Tanaka and Johnston Space Analysis: Does It Apply to All Populations?
(/products/ejournals/abstract/10.1055/s-0044-1782187)



HTML (/products/ejournals/html/10.1055/s-0044-1782187)
PDF (1171 kb) (/products/ejournals/pdf/10.1055/s-0044-1782187.pdf)

Publication Date: 13 May 2024

Original Article

Damnoensawat, Phatthra; Mitrakul, Kemthong:

Quantitative Analysis of *Bifidobacterium* and *Scardovia wiggisiae* in Dental Plaque from Children in Northern Thailand and Their Association with Caries Factors (/products/ejournals/abstract/10.1055/s-0044-1785473)



HTML (/products/ejournals/html/10.1055/s-0044-1785473)
PDF (422 kb) (/products/ejournals/pdf/10.1055/s-0044-1785473.pdf)

Original Article

Redwan, Hetaf S.; Hussein, Mohamed A.; Abdul-Monem, Mohamed M.:

Effect of Bleaching on Surface Roughness and Color Parameters of Coffee-Stained Nanohybrid Dental Composites with Different Viscosities (/products/ejournals/abstract/10.1055/s-0044-1786752)



HTML (/products/ejournals/html/10.1055/s-0044-1786752)
PDF (1626 kb) (/products/ejournals/pdf/10.1055/s-0044-1786752.pdf)

Publication Date: 09 May 2024

Original Article

Abdul-Monem, Mohamed M.; Hussein, Mohamed A.; Abdelrehim, Mona G.:

Effect of Repolishing on Color Stability, Translucency, and Surface Roughness of Aged Monochromatic Dental Composites (/products/ejournals/abstract/10.1055/s-0044-1786678)



HTML (/products/ejournals/html/10.1055/s-0044-1786678)
PDF (6402 kb) (/products/ejournals/pdf/10.1055/s-0044-1786678.pdf)

Publication Date: 06 May 2024

Original Article

Elsharkawy, Moustafa Ahmed; Ibrahim, Radwa Ali; Hamama, Hamdi Hosni; Mahmoud, Salah Hasab:

Bulk-FillOrmocer versus Methacrylate-Based Resin Composite Restorative Systems: The Effect of Flowable Lining on Two-Year Clinical Performance in Class II Cavities (/products/ejournals/abstract/10.1055/s-0044-1785472)



HTML (/products/ejournals/html/10.1055/s-0044-1785472)
PDF (251 kb) (/products/ejournals/pdf/10.1055/s-0044-1785472.pdf)

Original Article

Sayed, Moustafa Mohammed; Radad, Khaled; Rausch-Fan, Xiaohui; Elheeny, Ahmad:

Histopathological Evaluation of Pulp Reaction to Potassium Nitrate in Polycarboxylate Cement Compared with Mineral Trioxide Aggregate in Immature Dogs' Teeth (/products/ejournals/abstract/10.1055/s-0044-1785475)



HTML (/products/ejournals/html/10.1055/s-0044-1785475)
PDF (3083 kb) (/products/ejournals/pdf/10.1055/s-0044-1785475.pdf)

Publication Date: 02 May 2024

Original Article

Dias, Randi; Kusdhany, Lindawati S.; Indrasari, Muslita:

Contributing Factors of Dysphagia in Indonesian Elderly (/products/ejournals/abstract/10.1055/s-0044-1782184)



HTML (/products/ejournals/html/10.1055/s-0044-1782184)
PDF (888 kb) (/products/ejournals/pdf/10.1055/s-0044-1782184.pdf)

Publication Date: 01 May 2024

Original Article

Salah, Lina; Alkhames, Hussain M.; Alhaj, Zokaa; Al-Jubran, Saleh M. A.; Mohammed, Aseel; Baleis, Hussain H. I.; Almadeh, Naji M. M.; Ibrahim, Amir I. O.:

A Cross-Sectional Cone-Beam Computed Tomography Study of Root Canal Morphology and Configuration of Mandibular Premolars in Saudi and Sudanese Subpopulations (/products/ejournals/abstract/10.1055/s-0044-1785474)



HTML (/products/ejournals/html/10.1055/s-0044-1785474)
PDF (1007 kb) (/products/ejournals/pdf/10.1055/s-0044-1785474.pdf)

Publication Date: 29 April 2024

Original Article

Yasmin, Winarni; Vaithilingam, Rathna Devi; Halib, Nadia; Mohamad Hanafiah, Rohazila; Ramli, Haslinda:
Periodontal Health and Microbiological Characteristics of Siwak (Chewing Stick) and Toothbrush Users:
A Cross-Sectional Study (/products/ejournals/abstract/10.1055/s-0044-1782185)



HTML (/products/ejournals/html/10.1055/s-0044-1782185)
PDF (597 kb) (/products/ejournals/pdf/10.1055/s-0044-1782185.pdf)

European Journal of General Dentistry

Honorary Editor-in-Chief

Nairn H F Wilson

King's College London Dental Institute,
UK

Editors-in-Chief

Necdet Adanir

Department of Restorative Dentistry,
College of Dentistry, King Faisal
University, Al Ahsa, KSA

Zohaib Khurshid

Department of Prosthodontics and
Dental Implantology, College of
Dentistry, King Faisal University,
Al-Ahsa, KSA

Deputy Editor

Muhammad S Zafar

Department of Restorative Dentistry,
College of Dentistry, Taibah University,
Al Madina Al Munawwarra, KSA

Scientific Advisory Board

Amit Arora

Oral Health Services, Sydney Local
Health District and Sydney Dental
Hospital, Surry Hills, NSW, Australia

Bruno Chrcanovic

Department of Prosthodontics, Faculty
of Odontology, Malmö University,
Malmö, Sweden

Carla V. Gurgel

Department of Pediatric Dentistry,
Orthodontics and Public Health, Bauru
School of Dentistry, USP - University of
São Paulo, Bauru, Brazil

Christian Morsczeck

Department of Oral and Maxillofacial
Surgery, University Hospital Regens-
burg, Regensburg, Germany

Clement C. Azodo

Department of Periodontics, Univer-
sity of Benin Teaching Hospital, Benin
City, Nigeria

Dilek Arslantunali Tagtekin

Department of Restorative Dentistry,
Faculty of Dentistry, Marmara Universi-
ty, Istanbul, Turkey

Durga Shanker Gupta

Departments of Oral and Maxillofacial
Surgery, Teerthanker Mahaveer Dental
College and Research Centre, Morad-
abad, Uttar Pradesh, India

Elio Hitoshi Shinohara

Departament of Oral and Maxillofacial
Surgery, Hospital Vivaldo Martins
Simões, Osasco, Brazil

Ella A Naumova

Department of Biological and Mate-
rial Sciences in Dentistry, Faculty of
Health, Witten/Herdecke University,
Witten, Germany

E. Nursen Topcuoglu

Department of Microbiology, Faculty
of Dentistry, Istanbul University, Istan-
bul, Turkey

Filiz Namdar Pekiner

Department of Oral and Maxillofacial
Radiology, Faculty of Dentistry, Mar-
mara University, Istanbul, Turkey

Folakemi A. Oredugba

Department of Child Dental Health,
Faculty of Dental Sciences, College of
Medicine University of Lagos, Lagos,
Nigeria

Francisco Vera-Sempere

Department of Pathology, La Fe Uni-
versity Hospital, Valencia, Spain

Fusun Ozer

Department of Preventive and Re-
storative Sciences, School of Dental
Medicine University of Pennsylvania,
Philadelphia, USA

Hakan Turkkahraman

Department of Orthodontics and Oral
Facial Genetics, School of Dentistry,
Indiana University, Indianapolis,
Indiana, USA

Henk S. Brand

Department of Oral Biochemistry. Aca-
demic Centre for Dentistry Amsterdam
(ACTA), Amsterdam, The Netherlands

Izzet Yavuz

Department of Pedodontics, Faculty of
Dentistry, Dicle University, Diyarbakir,
Turkey

James Bahcall

Department of Endodontics, Univer-
sity of Illinois-Chicago, College of
Dentistry, Chicago, Illinois, USA

Jenny Abanto

Department of Pediatric Dentistry and
Orthodontics, School of Dentistry,
University of São Paulo, São Paulo, SP,
Brazil

Jithendra Ratnayake

Department of Oral Science, Faculty
of Dentistry, University of Otago,
Dunedin, New Zealand

José Renato Cavalcanti de Queiroz

Department of Biotechnology, College
of Health, Potiguar University, Natal,
RN, Brazil

Josué Martos

Department of Semiology and Clinics,
School of Dentistry, Federal University
of Pelotas (UFPEL), Pelotas, Brazil

Karen Vinal-Collier

Department of Dental Public, Leeds,
School of Dentistry, University of
Leeds, UK

Lisa M. Jamieson

Australian Research Centre for Pop-
ulation Oral Health, Adelaide Dental
School, The University of Adelaide,
Adelaide, Australia

Luca Testarelli

Department of Oral and Maxillo-Facial
Sciences, "La Sapienza" University of
Rome, Rome, Italy

Lumnije Kqiku

Department of Dental Medicine and
Oral Health, Division of Restorative
Dentistry, Periodontology and Prost-
hodontics, Medical University of Graz,
Graz, Austria

Martin Gosau

Department of Oral and Maxillofacial
Surgery, University Hospital Ham-
burg-Eppendorf, Hamburg, Germany

Michele Callea

Unit of Dentistry, Bambino Gesù Children Hospital and Research Institute, Rome, Italy

Nidambur Vasudev Ballal

Department of Conservative Dentistry and Endodontics, Manipal College of Dental Sciences, Manipal University, Manipal, Karnataka, India

Nikolaos A. Chrysanthakopoulos

Department of Pathological Anatomy, Medical School, University of Athens, Athens, Greece

Pierre Le Bars

Division of Fixed Prosthodontics, CHU Nantes, Nantes, France

Renato N. Hopp

Department of Oral Diagnosis, Piracicaba Dental School, University of Campinas, Piracicaba, São Paulo, Brazil

Shiu-yin Cho

Department of Health, Fanling School Dental Clinic, School Dental Care Service, Fanling, New Territories, Hong Kong

Sunit Kumar Jurel

Department of Prosthodontics, King George's Medical University, Lucknow, India

Uma Shanker Pal

Department of Oral and Maxillofacial Surgery, King George's Medical University, Lucknow, Uttar Pradesh, India

Vytaute Peciuliene

Institute of Odontology, Faculty of Medicine, Vilnius University, Vilnius, Lithuania

Wolfgang H. Arnold

Department of Biological and Material Sciences in Dentistry, School of Dentistry, Faculty of Health, Witten/Herdecke University, Germany

Xiaoli Gao

Faculty of Dentistry, National University of Singapore, Singapore, Singapore

Yogesh Kini

Department of Oral and Maxillofacial Surgery, D.Y. Patil Dental College, Navi Mumbai, India

© 2021. European Journal of General Dentistry. All rights, including the rights of publication, distribution, and sales, as well as the right to translation, are reserved. No part of this work covered by the copyrights hereon may be reproduced or copied in any form or by any means – graphic, electronic, or mechanical, including photocopying, recording, taping, or information and retrieval systems - without written permission of the publisher.

European Journal of General Dentistry is published three times per year in January, May, and September by Thieme Medical and Scientific Publishers Private Limited, A – 12, Second Floor, Sector - 2, Noida - 201301, Uttar Pradesh, India. Tel: +91-120-4556600, Fax: +91-120-455-6649.

Subscription: Open access journals available online for free at <http://open.thieme.com>.

Advertisers contact: Marketing, Thieme Publishers Delhi, A-12, Second Floor, Sector -2, Noida-201301, Uttar Pradesh, marketing@thieme.in.

European Journal of General Dentistry is indexed in SCOPUS, EBSCO, ProQuest, Google Scholar. Thieme Medical Publishers is a member of the CrossRef initiative.

Editorial comments should be sent to journals@thieme.com. The content of this journal is available online at www.thieme-connect.com/Products. Visit our website at www.thieme.com and the direct link to this journal at www.thieme.com/ejgd.

Typesetting: DiTech Process Solutions Pvt. Ltd. Mumbai, India

Printing and Binding: Sai Printo Pack (Pvt) Ltd.

Printed in India



Effect of Saga Leaf Extract (*Abrus precatorius* Linn) in Inhibiting *Enterococcus faecalis* Bacteria Growth as an Alternative Root Canal Irrigation Material

Pitri Ayu Puspita Sari¹ Vinna Kurniawati Sugiama¹ Rudy Djuanda¹

¹ Faculty of Dentistry, Maranatha Christian University, Bandung, West Java, Indonesia

² Department of Oral Biology, Faculty of Dentistry, Maranatha Christian University, Bandung, West Java, Indonesia

³ Department of Conservative Dentistry and Endodontic, Faculty of Dentistry, Maranatha Christian University, Bandung, West Java, Indonesia

Address for correspondence Vinna Kurniawati Sugiama, drg, MKes, PBO, CMC, Department of Oral Biology, Faculty of Dentistry, Maranatha Christian University, Bandung, West Java 40164, Indonesia (e-mail: vinnakurniawati@yahoo.co.id).

Eur J Gen Dent

Abstract

Objective This study aims to examine the antibacterial effect of saga leaf extract on *Enterococcus faecalis* bacteria, which causes root canal treatment failure.

Materials and Methods This research was conducted using a laboratory experimental method with saga leaf extract at 3.125, 6.25, 12.5, 25, 50, and 100% concentrations. Two percent chlorhexidine acts as the positive control, and 100% dimethyl sulfoxide as the negative control. The diameter of the inhibition zones was measured using the well diffusion test method.

Statistical Analysis Data from the measurement of inhibition zone diameter were obtained and tested statistically using the normality test (Shapiro–Wilk), homogeneity test (Levene), parametric test (one-way analysis of variance), and further test (post hoc test).

Results The largest inhibition zone diameter observed in this study was 9.46 mm at 100% concentration; however, it was not bigger than the positive control, which was measured at 16.55 mm. The research data were analyzed based on the classification of Davis and Stout inhibition zones.

Conclusion This study concludes that saga leaf extract has an antibacterial effect on the growth of *E. faecalis* bacteria.

Keywords

- ▶ *Enterococcus faecalis*
- ▶ *Abrus precatorius* Linn
- ▶ root canal treatment
- ▶ well diffusion
- ▶ antibacterial

Introduction

Pulp necrosis and irreversible pulpitis are the most common cases found in dentistry. They can be prevented and treated with root canal treatment.^{1,2} Root canal treatment is a dental procedure that aims to maintain or improve the condition of infected tooth to be accepted biologically by the tissues. Successful root canal treatment depends on the ability to eliminate microorganisms from the root canal system and the reinfection prevention.^{3,4}

The most common root canal microorganisms isolated from infections are anaerobic bacteria, one of which is

Enterococcus faecalis bacterium. It is considered to be the main cause of root canal abnormalities with a prevalence value of 77%.^{5,6} *Enterococcus faecalis* is the dominant species of facultative anaerobic gram-positive cocci that exist in pairs, singles, short chains, oval or rounded egg shaped.^{7,8} *Enterococcus faecalis* can survive and multiply in the root canal with poor nutrient, high pH (alkaline) up to 11.5, and any help from other bacteria.⁹ The resilience of *E. faecalis* to survive in an unfavorable root canal environment causes this bacterium to be a pathogen that leads to root canal treatment failure. The failure can be prevented by using an appropriate irrigation with low toxicity solution.⁷ An ideal irrigation

DOI <https://doi.org/10.1055/s-0044-1788258>.
ISSN 2320-4753.

© 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

material should be nontoxic, able to dissolve organic and inorganic tissues, prevent smear layers formation during root canal preparation, and have antimicrobial properties.⁴

Two percent chlorhexidine (CHX) was suggested as a root canal irrigation agent because its antimicrobial effect can effectively protect the root canal after root canal treatment at that concentration. However, 2% CHX is not recommended as the first option for irrigation agents since it causes discoloration and allergic reactions when used repeatedly over a long period of time.^{10,11} The damaging effects of these irrigation agents have encouraged people to search alternative treatments, such as herbal medicine. One herbal medicinal plant often utilized as traditional medicine by the community is saga leaves.⁸

Abrus precatorius Linn is known as cough medicine and treatment for stomatitis, pharyngitis, and tonsillitis.¹² According to research, it has been reported that saga leaves (*A. precatorius* Linn) have preclinical anti-inflammatory properties. Saga leaf extract contains flavonoids, terpenoids, tannins, alkaloids, and saponin compounds that are effective as antibacterials by inhibiting cyclooxygenase and lipoxygenase.^{13,14} The antibacterial activity of saga leaves has been tested and proven effective against *Streptococcus mutans* bacteria, a gram-positive facultative anaerobic bacteria.¹⁵ This study was conducted to determine the effect of different saga leaf concentrations extract in inhibiting *E. faecalis* bacteria growth.

Materials and Methods

This was a laboratory experimental study with in vitro posttest only control group design by giving different treatments to *E. faecalis* bacteria with saga leaf extract (*A. precatorius* Linn) in various concentrations with the well diffusion test method. The formation of inhibition zone was calculated in each treatment. This research was conducted at Aretha Medika Utama Laboratory, Bandung.

Plant Determination

The saga leaves were obtained from a local herbal plantation in South Bogor subdistrict, Bogor city, West Java, which was then identified at the Biology Laboratory, Padjadjaran University, Jatinangor, West Java, to obtain the determination of *A. precatorius* Linn.

Concentration Series Preparation

This study was treated with a positive control in the form of 2% CHX, negative control in the form of 100% dimethyl sulfoxide (DMSO), and saga leaf extract (*A. precatorius* Linn) with concentrations of 3.125, 6.25, 12.5, 25, 50, and 100%. Calculation of dilution of saga leaf extract solution and 100% DMSO were performed to make series of concentrations. The varying concentration of extracts used are as follows:

100% saga leaf extract: stock solution.

50% saga leaf extract: 500 μ L stock solution + 500 μ L DMSO 100% (Solution A).

25% saga leaf extract: 500 μ L solution A + 500 μ L DMSO 100% (Solution B).

12.5% saga leaf extract: 500 μ L solution B + 500 μ L DMSO 100% (Solution C).

6.25% saga leaf extract: 500 μ L solution C + 500 μ L DMSO 100% (Solution D).

3.125% saga leaf extract: 500 μ L of solution D + 500 μ L of 100% DMSO.

Preparation of Saga Leaf Extract

This research was conducted at the Central Laboratory, Padjadjaran University, Jatinangor, West Java. A total of 1 kg of saga leaves was dried and pulverized using blender to form fine powder; 287 g of saga leaf powder were used in the first maceration process by soaking it with 1 L of 96% ethanol solvent in a tightly closed container. The soaking process was performed for 72 hours and mixed occasionally. The saga leaf solution then proceeds to the filtering stage using a filter paper to separate the solution. The filtrate was separated and placed in a glass bottle. The filtrate was then evaporated using vacuum rotary evaporator with a temperature of 45 to 50°C; 50 rpm speed; 170–180 mbar pressure for roughly 5 hours until a thick saga leaf extract was obtained. Furthermore, the dilution process was performed using 100% DMSO at a concentration of 3.125, 6.25, 12.5, 25, 50, and 100%. Finally, the ethanol extract of saga leaves with various concentrations were placed into closed sterile bottles and stored in the refrigerator at -20°C .

Phytochemical Analysis

Phytochemical analysis test was conducted to test the presence of bioactive compounds, such as tannins, flavonoids, alkaloids, steroids, triterpenoids, phenolics, and saponins from the saga leaf extract. Alkaloids were identified using Dragendorff reagent. Flavonoids were identified using HCl + Mg, H_2SO_4 , and NaOH 10%. Saponins were detected by heating method. Tannins and phenolics were identified using 1 and 5% FeCl_3 reaction, and steroids and triterpenoids were analyzed with $\text{H}_2\text{SO}_4 + \text{CH}_3\text{COOH}$. Qualitative results were expressed as (+) for presence and (–) for absence of phytochemicals.

Culture of *Enterococcus faecalis* Bacteria

The sample in this study was *E. faecalis* bacteria (ATCC 29212) obtained from Aretha Medika Utama Laboratory, Bandung. The study used 19 g of Mueller–Hinton agar (MHA) and 10.5 g of Mueller–Hinton broth (MHB) as the growth media which were measured using an analytical balance. Microwave was used to help dissolve the growth media in 500 mL of ddH₂O. The next step was the sterilization process using an autoclave at 121°C with a pressure of 1.5 atm for 20 minutes under anaerobic conditions. Colony suspension method was used to prepare the bacterial inoculum by inoculating *E. faecalis* colonies that had been cultured for 18 to 24 hours on MHA medium, into MHB medium. The solution's turbidity was adjusted according to the McFarland 0.5 standard solution's turbidity to produce an inoculum with a bacterial count

of ~ 1 to 2×10^8 CFU/mL. Pure culture of bacteria was taken as much as 1 ose and implanted on MHA by swab method.

Diffusion Test

The inoculation process on the test agar plates was performed using the swab method by dipping a sterile cotton swab into the prepared bacterial suspension. The cotton swab was pressed against the tube wall to remove excess suspension, which was then applied to the surface of the MHA medium evenly. The inoculation was allowed to rest at room temperature for 3 to 5 minutes until the suspension was absorbed into the agar. After that, holes were made in the MHA medium using 8-mm-diameter tips. Each hole was filled with 50 μ L of different concentrations of saga leaf extract, starting with 3.125, 6.25, 12.5, 25, 50, and 100%, the positive control 2% CHX, and the negative control DMSO 100%, respectively. In this study, the test agar wells were made for three repetitions. The agar plates were first incubated at 37°C for 24 hours before measuring the diameter of the inhibition zone formed using a caliper.

Statistical Analysis

Data from the measurement of inhibition zone diameter were obtained and tested statistically using the normality test (Shapiro–Wilk), homogeneity test (Levene), parametric test (one-way analysis of variance), and further test (post hoc test).

Results

Phytochemical Examination

The extracted saga leaf plants were then subjected to qualitative phytochemical testing with the following test results.

Based on the results of phytochemical tests, only secondary metabolite compounds flavonoids appeared in low amounts (+) with 10% NaOH testing, while saponins, tannins, steroids, and phenolic appeared in moderate amounts (++).

Measurement of Inhibition Zone Diameter of Saga Leaf Extract against *Enterococcus faecalis*

In this study, antibacterial measurements were calculated from the inhibition zone diameter. The inhibition area measured is a clear zone where *E. faecalis* bacteria did not

grow. The clear zone area was measured using a caliper in millimeters (mm). The results are shown in ►Tables 1 and 2, and ►Figs. 1 and 2.

Based on the tables and figures above, it can be inferred that saga leaf extract shows antimicrobial activity against *E. faecalis* bacteria characterized by the presence of inhibition zone diameter at 100, 50, 25, 12.5, 6.25, and 3.125% concentrations of saga leaf extract. The largest inhibition zone was with 100% concentration with an average diameter of 9.46 mm, and the lowest inhibition zone was with 3.125% concentration with an average diameter of 1.26 mm. The level of inhibition shown by saga leaf extract is directly proportional to the level of concentration where higher extract concentration produces higher inhibition area.

Discussion

Inferring from the data, saga leaf extract shows an effect in inhibiting *E. faecalis* bacteria growth. The diameter of the inhibition zone grows from using the smallest to the largest concentration where the smallest inhibition zone happened at 3.125% concentration with 1.26 mm, whereas the largest inhibition zone was at 100% concentration at 9.46 mm diameter. The result can be explained due to higher concentration of saga leaf extract has more bioactive compounds contained. These bioactive compounds cause the inhibition zone to appear on the bacterial culture media.

The ability of saga leaf extract to inhibit the growth and kill *E. faecalis* bacteria is related to bioactive compounds that has antibacterials properties. The results of qualitative phytochemical tests on saga leaf extract used in this study showed the presence of phenolic compounds, tannins, saponins, steroids in moderate amounts; flavonoids in small amounts; and no alkaloid compounds were detected. There are factors that causes failure to detect alkaloid compounds such as soil type, soil pH, organic matter content, air temperature, and rainfall in the area where the plant grows. These factors can cause content in the plants of each region to have different cadences.^{16,17}

Phenolic compounds are said to prevent systemic inflammation by restoring redox balance by modulating the inflammatory response through mitigating the cytokine

Table 1 Phytochemical test results on saga leaf extract (*Abrus precatorius* Linn)

No.	Metabolite compounds	Test method	Test results
1	Alkaloid	Dragendorff reagent	–
2	Flavonoid	Concentrated HCl and Mg reagent	–
		2N H ₂ SO ₄ reagent	–
		10% NaOH reagent	+
3	Saponin	Heated	++
4	Tannin	1% FeCl ₃ reagent	++
5	Steroid	Concentrated H ₂ SO ₄ reagent and anhydrous CH ₃ COOH	++
6	Triterpenoid	Concentrated H ₂ SO ₄ reagent and anhydrous CH ₃ COOH	–
7	Phenolic	5% FeCl ₃ reagent	++

Table 2 Results of measurement of the diameter of the zone of inhibition of saga leaf extract against *Enterococcus faecalis* bacteria

Treatment	Inhibition zone diameter (mm)			Mean	Standard Deviation	RSD
	P 1	P 2	P 3			
Positive control (2% chlorhexidine)	17.50	17.06	15.10	16.55	1.28	7.72
Negative control (dimethyl sulfoxide 100%)	0.00	0.00	0.00	0.00	0.00	0.00
100% saga leaf extract	9.12	9.69	9.57	9.46	0.30	3.18
50% saga leaf extract	8.08	7.59	7.50	7.72	0.31	4.04
25% saga leaf extract	5.50	4.56	5.06	5.04	0.47	9.33
12.5% saga leaf extract	3.45	3.01	4.09	3.52	0.54	15.44
6.25% saga leaf extract	2.12	2.15	2.19	2.15	0.04	1.63
3.125% saga leaf extract	1.43	1.33	1.08	1.26	0.25	19.72

Abbreviation: RSD, relative standard deviation.

pathway to degrade oxidative stress. However, the actual mechanism of phenolic acid itself in inhibiting bacteria is not fully understood because it has a complex chemical structure. In some studies, it is hypothesized that phenolic acids can damage the electrochemical gradient of the mitochondrial membrane and prevent systemic inflammation.^{18,19}

Saponin compounds are able to react with porins (trans-membrane proteins) on the outer membrane of the bacterial cell wall to form strong polymer bonds that causes porin damage.²⁰ Damage to porins will reduce the permeability of the bacterial cell membrane since it is the entrance and exit of the compound. It causes bacterial cells to lack nutrients, inhibited its growth and die.^{20,21} This helps the entry of tannin and flavonoid compounds to enter bacterial cells.

Tannins activates reverse transcriptase, the adhesions of the microbial cell, DNA topoisomerase enzymes that will interfere with bacterial DNA synthesis, and also attack cell wall polypeptides, causing damage to bacterial cell wall.²² All of this is possible because tannins have a target on the polypeptide wall of bacterial cells, resulting in incomplete cell wall formation and then bacterial cells will die.²⁰


It is reported that steroids act as antibacterial due to lipid membrane correlation and sensitivity to steroid compounds that show leakage in liposomes. Interaction between steroids and cell phospholipid membranes that are permeable to

lipophilic compounds causes integrity of the membrane to decrease and changes to the morphology of the cell membrane, causing bacterial cells to become fragile and lysed.^{23,24}

Flavonoids function as antibacterial by inhibiting bacterial growth by forming complex compounds against extracellular proteins that disrupt the integrity of the bacterial cell membrane. The compound denatured the bacterial cell proteins and damaged the cell membrane to prevent reparation process.²⁵ Furthermore, bacterial mobility is also inhibited by flavonoids due to the presence of flavonoid's hydroxyl groups that cause alterations in organic components and nutrient transport that lead to harmful effects on bacteria.²⁶

Abrus precatorius contains the lethal toxin Abrin, a toxalbumin that inhibits protein synthesis leading to cell death and tissue damage.³ It is necessary to investigate further the toxicity of its active constituents and side effects.

Abrin is a type-II ribosome-inactivating protein from *A. precatorius* seeds that is similar in structure and properties to ricin. It is classified as a Category B bioterrorism warfare agent. Abrin is 75 times more toxic than ricin, and is classified as a Category B biological warfare agent in the United States due to its widespread source, ease of preparation, and lethality. Abrin has a molecular weight of ~63 kDa, consisting of A chain with N-glycosidase activity and a galactose-

**Fig. 1** The zone of inhibition of saga leaf extract against *Enterococcus faecalis* shown by a clear area.




Fig. 2 Comparison of inhibition zone diameter of saga leaf extract against *Enterococcus faecalis* bacteria.

specific lectin B chain. The structure of the abrin B chain consists of lectin that binds to the β -D-galactoside portion of the cell surface and mediates the internalization/endocytosis of the entire toxin into the host cell. There are four isoforms of abrin: abrin-a, abrin-b, abrin-c, and abrin-d. Abrin-a and abrin-d are the most toxic. Abrin-b and abrin-c show weak B-chain lectin activity.^{1,2}

After entering the cell, the A chain can hydrolyze the N-glycosidic bond of the 28S ribosomal RNA of eukaryotic cells and catalyze depurination, causing ribosome inactivation and inhibits protein synthesis, therefore causing cell apoptosis. Due to its high toxicity, consumption by human will cause death due to multiple organ failure. Abrin is highly toxic, with an estimated fatal dose of 0.1 to 1 μ g/kg. It has caused death after intentional or accidental poisoning. Clinically, abrin is considered an immunotoxin that can target cancer cells and can be delivered to the tumor site after combining with antitumor drugs.¹⁻⁴

At the cellular level, abrin inhibits protein synthesis, leading to cell death. Many of the features observed in abrin poisoning can be shown by abrin-induced endothelial cell damage, which leads to increased capillary permeability. This results in fluid and protein leakage and tissue edema (vascular leak syndrome). The most reported cases of human poisoning involve the consumption of jequirity beans, most of which cause gastrointestinal toxicity.²

In previous studies, saga leaf extract (*A. precatorius* Linn) was shown to have antibacterial activity by inhibiting several bacteria, both gram-negative and gram-positive bacteria, such as *S. mutans* bacteria which are included in the gram-positive facultative anaerobic bacteria group. Saga leaf extract (*A. precatorius* Linn) was proven to effectively work as an antibacterial at concentrations of 50, 25, 12.5, and 6.25%. It was also proven at 50% concentration of saga leaf extract (*A. precatorius* Linn) that the inhibition zone was formed to inhibit *Streptococcus mutans* bacteria at a diameter of

10.3 mm. Therefore, the results of this research are comparable to the results of previous studies conducted that shows ethanol extract of saga leaves (*A. precatorius* Linn) works effectively as an antibacterial.

Conclusion

Based on the results of the study, it can be concluded that saga leaf extract (*A. precatorius* Linn) is able to show antimicrobial activity against *E. faecalis* bacteria characterized by the presence of inhibition zones formed in different concentrations of saga leaf extract, starting at the smallest concentration of 3.125% to the largest concentration of 100%. The level of inhibition shown by saga leaf extract is directly proportional to the level of concentration, which means that the higher the concentration of the extract, the higher the inhibition produced. The level of inhibition of the extract is lower than that of CHX solution and is significantly different based on the statistical tests conducted.

Conflict of Interest

None declared.

References

- Khairunnisa K, Yanuar Ichrom Nahzi M, Diana S. Toxicity test of Dayak onion bulbs extract (*Eleutherine Palmifolia* (L) Merr) on artemia salina leach using BSLT method. DENTINO: Jurnal Kedokteran Gigi 2018;III(01):91-95
- Nishio Ayre W, Melling G, Cuveillier C, et al. *Enterococcus faecalis* demonstrates pathogenicity through increased attachment in an ex vivo polymicrobial pulpal infection. Infect Immun 2018;86(05):1-14
- Kuntari LM, Hadriyanto W, Mulyawati E. Perbedaan daya antibakteri klorheksidin 2% dan berbagai konsentrasi sodium hipoklorit kombinasi omeprazole 8,5% terhadap *Enterococcus faecalis*. Jurnal Kedokteran Gigi 2014;5(02):139-149

- 4 Urkande NK, Mankar N, Nikhade PP, Chandak M. Beyond tradition: non-surgical endodontics and Vital Pulp Therapy as a dynamic combination. *Cureus* 2023;15(08):e44134
- 5 Tamara R, Rochyani L, Budi Teguh P. Daya hambat ekstrak teripang emas (*Stichopus hermanii*) terhadap bakteri *Enterococcus faecalis*. *Denta Jurnal Kedokteran Gigi* 2015;9(01):37–47
- 6 Singh H. Scient open access exploring the world of science microbiology of endodontic infections. *J Dent Oral Health* 2016; 2(5.044):3–4
- 7 Soraya C, Sunnati S, Maulina V. Efek antibakteri ekstrak batang serai (*Cymbopogon citratus*) terhadap pertumbuhan *Enterococcus faecalis*. *Cakradonya Dent J* 2016;8(02):69–78
- 8 Sainudeen S, Nair VS, ZARBH M, Abdulla AM, Najeeb CM, Ganapathy S. Can herbal extracts serve as antibacterial root canal irrigating solutions antimicrobial efficacy of *Tylophora indica*, *Curcumin longa*, *Phyllanthus amarus*, and sodium hypochlorite on *Enterococcus faecalis* biofilms formed on tooth substrate: in vitro study. *J Pharm Bioallied Sci* 2020;12(Suppl 1): S423–S429
- 9 Alghamdi F, Shakir M. The influence of *Enterococcus faecalis* as a dental root canal pathogen on endodontic treatment: a systematic review. *Cureus* 2020;12(03):e7257
- 10 Guneser MB, Eldeniz AU. The effect of gelatinase production of *Enterococcus faecalis* on adhesion to dentin after irrigation with various endodontic irrigants. *Acta Biomater Odontol Scand* 2016; 2(01):144–149
- 11 Drews D-J, Nguyen AD, Diederich A, Gernhardt CR. The Interaction of two widely used endodontic irrigants, chlorhexidine and sodium hypochlorite, and its impact on the disinfection protocol during root canal treatment. *Antibiotics* 2023;12(03):589
- 12 Widiyanto E, Santoso DB, Kardiman K, Nugraha AE. Pemberdayaan masyarakat tentang pemanfaatan tanaman saga (*Abrus Precatorius L*) di desa tanahbaru pakisjaya Karawang. *Aksiologi: Jurnal Pengabdian Kepada Masyarakat* 2019;4(01):63
- 13 Wound O, Rabbits T, Cuniculus O, et al. Test the effectiveness of saga leaf extract (*Abrus precatorius* Linn). *J Pharm Sci* 2021;4(02):111–116
- 14 Rambe R, Paramitha R, Ginting E, Caniago MY. Test the effectiveness of saga leaf extract (*Abrus precatorius* Linn) ointment for wound treatment in rabbits (*Oryctolagus Cuniculus*). *J Pharm Pharm Sci* 2021;4(02):111–116
- 15 Andika NA. Aktivitas antibakteri ekstrak dan fraksi aktif daun saga (*Abrus Precatorius L.*) terhadap bakteri *Streptococcus mutans* ATCC 25175. *Warta Bhakti Husada Mulia Jurnal Kesehatan* 2022;9 (02):1–11
- 16 Jansen G, Jürgens HU, Schliephake E, Ordon F. Effect of the soil pH on the alkaloid content of *Lupinus angustifolius*. *Int J Agron* 2013. Doi: 10.1155/2012/269878
- 17 Abror Zamzami M. Pengaruh penyangraian kopi dalam mengembangkan cita rasa, kesehatan dan daya saing kopi Indonesia. *RADAR Plantation* 2021;2(01):1–7
- 18 Gu H, Zhu Y, Peng Y, et al. Physiological mechanism of improved tolerance of *Saccharomyces cerevisiae* to lignin-derived phenolic acids in lignocellulosic ethanol fermentation by short-term adaptation. *Biotechnol Biofuels* 2019;12(01):268
- 19 Cory H, Passarelli S, Szeto J, Tamez M, Mattei J. The role of polyphenols in human health and food systems: a mini-review. *Front Nutr* 2018;5:87
- 20 Rahmawati A, Mayasari D, Narsa AC. Kajian literatur: Aktivitas antibakteri ekstrak herba suruhan (*Peperomia pellucida L.*). *Proceeding of Mulawarman Pharmaceuticals Conferences* [Internet]. 2020;12:117–124
- 21 Saptowo A, Supriningrum R, Supomo S. Uji Aktivitas antibakteri ekstrak kulit batang sekilang (*Embelia borneensis* Scheff) terhadap bakteri *Propionibacterium acnes* dan *Staphylococcus epidermidis*. *Al Ulum Sains Dan Teknologi* 2022;7(02):93–97
- 22 Ramonah D, Ryan Radix Rahardhian M, Nindya Putri C. Determinasi total flavonoid, total fenolik, dan aktivitas antibakteri ekstrak etanol daun insulin (*Smallanthus sonchifolius*) dengan metode perkolasi. *Media Farmasi Indonesia* 2020;15(01):1585–1592
- 23 Huda C, Putri AE, Sari DW, et al. Uji aktivitas antibakteri fraksi dari maserat *Zibethinus folium* terhadap *Escherichia coli*. *Jurnal Sain Health* 2019;3(01):7–14
- 24 Bontjura S, Waworuntu OA, Siagian KV, et al. Uji efek antibakteri ekstrak daun leilem (*Clerodendrum minahassae L*) terhadap bakteri *Streptococcus mutans*. *PHARMACON* 2015;4(04):96–101
- 25 Manik DF, Hertiani T, Anshory H, Mada G. Analisis korelasi antara kadar flavonoid dengan aktivitas antibakteri ekstrak etanol dan fraksi-fraksi daun kersen (*Muntingia calabura L*) terhadap *Staphylococcus aureus*. *KHAZANAH* 2014;6(02):1–11
- 26 Egra S, Rofin M, Adiwena M, Jannah N, Kuspradini H. Tohru Mitsunaga dan. Aktivitas antimikroba ekstrak bakau (*Rhizophora mucronata*) dalam menghambat pertumbuhan *Ralstonia solanacearum* penyebab penyakit layu. *AGROVIGOR: Jurnal Agroekoteknologi* 2019;12(01):26–31