

## BUKTI KORESPONDENSI

### ARTIKEL JURNAL INTERNASIONAL BEREPUTASI


**Judul Artikel** : **Anti-Inflammatory Activity of Sappan Wood Extract Cream (*Caesalpinia sappan*) in Porphyromonas gingivalis-induced Gingivitis Rats Models**

**Jurnal** : Open Veterinary Journal

**Penulis** : Vinna Kurniawati Sugiaman, Jeffrey, Wahyu Widowati, Nindia Salsabila Mia Dewi, Rival Ferdiansyah, Dhenta Maulana Muchtar, Dhanar Septyawan Hadiprasetyo

No	Perihal	Tanggal
1.	Register pada Open Veterinary Journal	02 Mei 2024
2.	Bukti konfirmasi submit artikel dan artikel yang disubmit	06 Mei 2024
3.	Bukti melakukan review yang pertama	20 Juli 2024
4.	Bukti konfirmasi submit revisi pertama yang telah direvisi	03 Agustus 2024
5.	Bukti melakukan review yang kedua	09 Agustus 2024
6.	Bukti konfirmasi submit artikel yang telah revisi kedua	15 Agustus 2024
7.	Bukti konfirmasi artikel diterima	19 Agustus 2024
8.	Bukti Galery Proof Manuscript	21 September 2024
9.	Bukti Publikasi Online Artikel	30 September 2024

**Register pada Open Veterinary Journal (02 Mei 2024)**



**Bukti konfirmasi submit artikel dan artikel yang disubmit (06 Mei 2024)**



Dear Vinna Kurniawati,

Your submission entitled **Anti-Inflammatory Activity of Sappan Wood Extract Cream (*Caesalpinia sappan*) in Porphyromonas gingivalis-induced Gingivitis Rats Models** (Manuscript Number: OVJ-2024-05-154) has been received by **Open Veterinary Journal**.

You could follow status of your manuscript by login to your author account at [www.ejmanager.com](http://www.ejmanager.com).

Thank you for submitting your work to our journal.

Best regards,

Editor  
Open Veterinary Journal  
<http://www.openveterinaryjournal.com>

## **Anti-Inflammatory Activity of Sappan Wood Extract Cream (*Caesalpinia sappan*) in Porphyromonas gingivalis-induced Gingivitis Rats Models**

Vinna Kurniawati Sugiaman<sup>1\*</sup>, Jeffrey Jeffrey<sup>2</sup>, Wahyu Widowati<sup>3</sup>, Nindia Salsabila Mia Dewi<sup>4</sup>, Rival Ferdiansyah<sup>5</sup>, Dhenta Maulana Muchtar<sup>5</sup>, Dhanar Septyawan Hadiprasetyo<sup>4,6</sup>

<sup>1</sup>Faculty of Dentistry, Maranatha Christian University, Jl. Surya Sumantri No. 65 Bandung 40164, Indonesia

<sup>2</sup>Department of Pediatric Dentistry, Faculty of Dentistry, Universitas Jenderal Achmad Yani, Cimahi 40531, Indonesia

<sup>3</sup>Faculty of Medicine, Maranatha Christian University, Jl. Surya Sumantri No. 65 Bandung 40164, Indonesia

<sup>4</sup>Aretha Medika Utama, Biomolecular and Biomedical Research Center, Indonesia, Jl Babakan Jeruk 2 no 9, Bandung 40163, Indonesia

<sup>5</sup>Department of Pharmaceutics, Sekolah Tinggi Farmasi Indonesia, Indonesia, Jl. Soekarno Hatta No.354, Bandung 40266, Indonesia

<sup>6</sup>Faculty of Pharmacy, Universitas Jenderal Achmad Yani, Cimahi 40531, Indonesia

Corresponding author: Vinna Kurniawati Sugiawan, Faculty of Dentistry, Maranatha Christian University,  
Bandung 40164, West Java, Indonesia. Tel: +62 812-2115-650; email:[vinna.ks@dent.maranatha.edu](mailto:vinna.ks@dent.maranatha.edu),  
[vinnakurniawati@yahoo.co.id](mailto:vinnakurniawati@yahoo.co.id)

## ABSTRACT

**Background:** Gingivitis is a dysbiotic condition characterized by persistent inflammation caused by a disease-associated multispecies bacterial population that has established itself in the subgingival region.

**Objectives:** The objectives of this study is to analyze the anti-inflammatory activity of sappan wood extract cream (SWC) in *Porphyromonas gingivalis* induced-periodontitis rats model. **Methods:** In this study, rats were infected to the gingiva with *P. gingivalis* as an inducer for gingivitis rats model for 14 days. The SWC were treated topically into the infected tissue. The inflammation and angiogenesis of gingival tissue was analyzed using histological examination. An immunohistochemistry (IHC) assay was used to analyze the Nuclear Factor kappa-B (NF-kB) protein expression. Masson Trichrome (MT) assay was used to analyze the histopathology of collagen Scores. The Tumor Necrosis Factor- $\alpha$  (TNF- $\alpha$ ), Interleukin-1  $\beta$  (IL-1 $\beta$ ), IL-6, and P38 genes expression were measured using quantitative Real Time-Polymerase Chain Reaction (qRT-PCR). **Result:** The induction of *P. gingivalis* -induced the gingivitis in rats model indicated by reddish gingiva and bacterial plaque. MT tests showed that SWC increased collagen density, HE test showed increased angiogenesis and in contrast, lowered inflammation score. IHC test showed that SWC decreased the expression of NF-kB protein. The TNF- $\alpha$ , IL-1 $\beta$ , IL-6, and P38 gene expression were also decreased due to the SWC treatment. **Conclusions:** Sappan wood extract cream proves that it has anti-inflammatory activity which has the potential to prevent or treat gingivitis

**Keywords:** Antiinflammation, *Caesalpinia sappan*, Cream, Gingivitis, Oxidative Stress.

## Introduction

Periodontitis is a complex chronic illness that is characterized by ongoing interactions between bacteria and inflammatory responses, which are influenced by both hereditary and extrinsic factors. Even though oral bacterial plaque has traditionally been considered the major cause of periodontal diseases (Kinane et al., 2017). Normally, the presence of a stable periodontal microbiota and a controlled host immune response leads to a condition of dynamic balance or homeostasis. Inflammation causes a decrease in fibroblast number and collagen degradation, which can lead to attachment loss and the creation of periodontal pockets (Naruishi, 2022). When a wound occurs in gingivitis, the cells that play a role in forming the ground substance and forming collagen fibers in closing the wound are fibroblast cells (Sardi et al., 2023). Fibroblasts are the dominant cells in the proliferation phase, forming new connective tissue and synthesizing collagen which affects tensile strength in tissue rejuvenation (Hanna & Giacomelli, 1997). Dysbiosis can impair periodontal tissue homeostasis by causing an imbalance within the microbial ecosystem, resulting in a s

ignificant production of Interleukin (IL)-1 members and initiating an over immune response via host receptors (Jia et al., 2019; Papathanasiou et al., 2020).

Cellular immune responses show the role of various cytokines in the process of inflammation of periodontal tissue (periodontitis). Different mediators influence the equilibrium in chronic inflammation. Pro-inflammatory cytokines like as IL-1 $\beta$ , Tumor Necrosis Factor-  $\alpha$  (TNF- $\alpha$ ), and IL-6 cause periodontal inflammation and tissue damage (Cardoso et al., 2018). Similarly, cytokine-based therapies have the potential to enhance both periodontitis and overall health (Cheng et al., 2020). Mitogen-activated protein Kinases (MAPKs), according to some researchers, are the initial signaling mediators of numerous inflammatory stimuli, including IL-1 $\beta$  and TNF- $\alpha$ . Furthermore, P38 MAPK, which plays an important part in the proinflammatory signaling system, can directly or indirectly regulate cytokines. Indeed, P38 MAPK is important in normal immunological and inflammatory responses; several studies have established its

participation in the generation of inflammatory cytokines that contribute to chronic inflammation (Lee & Kim, 2017). Nuclear Factor kappa beta (NF- $\kappa$ B) controls the inflammasomes and stimulates the production of a number of pro-inflammatory genes, including those encode cytokines and chemokines. As a result, abnormal NF- $\kappa$ B activation leads to the pathogenesis of numerous inflammatory illnesses (Liu et al., 2017).

Metronidazole is a popular periodontitis/gingivitis treatment. Metronidazole is effective against *P. gingivalis*, however, it can induce impaired taste, nausea, ataxia, neurotoxicity, xerostomia, brain illness, headache, seizures, and stomach cramps (Nastri et al., 2019; Vaithiyam et al., 2019). Therefore, a new agent for gingivitis treatment with minimum side effects might be a solution. Natural plants/plant extracts may be an alternative agent for treating the diseases. The main effect of the active substances in plant extracts has the potential as a treatment agent because these substances have antimicrobial activity, are useful as antioxidants, and are active components that can increase cell proliferation, angiogenesis and collagen production (Ghosh & Gaba, 2013).

Plant-derived chemicals have been identified as possible therapeutic agents due to their powerful biological capabilities. Polyphenols are advantageous constituents found in several plant-based. They are categorized into different groups, including flavonoids, polyphenolic amides, proanthocyanidins, as well as additional polyphenols like resveratrol and curcumin (Jayusman et al., 2022). *Caesalpinia sappan* L., also known as Brazil or Sappan wood, is a plant native to South East Asia and South India. It is a member of the Legingivainosae family. The heartwood of this plant has been historically utilised as a natural red dye, medicines, a traditional ingredient in food and beverages (Yang et al., 2010).

The examination of sappan wood's chemical constituents were resulting in a numerous structural kinds of phenolic components such as brazilin, chalcones, xanthone, homoisoflavonoids, coumarin, and flavones (Nirmal et al., 2015). Research has demonstrated that sappan wood comprises

diverse bioactive chemicals that possess anti-inflammatory properties. In this study, the sappan wood extract was made into a cream form to facilitate in gingivitis drug delivering. Because the easy to handle and the stabilization of the cream, the use of cream contains extract might be one of the solution for gingivitis treatment orally. Therefore, in this study we examine the antiinflammatory of sappan wood extract (*Caesalpinia sappan* L.) in gingivitis by measuring the activity of antiinflammatory, angiogenesis, NF-kB, collagen, also IL-1 $\beta$ , IL-6, TNF- $\alpha$  and p38 protein expression.

## **Material and Methods**

### ***Sappan Wood Extract Cream Preparation and LC-MS/MS Assay***

The Sappan Wood Extract Cream (SWC) was processed by the Sekolah Tinggi Farmasi Indonesia. The maceration technique was used in the extraction process of Sappan wood (*C. sappan* L). Fresh sappan wood is dried, crushed and filtered using a 60 mesh sieve. Sappan wood powder was then extracted using 96% ethanol for 4 days. Every 24 hours, the filtrate is collected and evaporated to obtain sappan wood extract (Widowati, 2011). The detection of bioactive component in sappan wood extract and determined using a liquid chromatography mass spectrometer (LC-MS/MS) (Widowati et al., 2023). The formulation of the Sappan Wood cream was from (Sayuti, 2015) with slight modifications. The cream preparations was made with 3.5 g Sodium Carboxymethylcellulose (Na CMC), 3.5 g Vaseline Flavum, 0.1 g Methyl Paraben, 2g Glycerin, 1g Tretanolamine, and 60 mg of SWC and Aquadest up to 100 ml.

### ***Animal Grouping and Treatment***

The rats were obtained from iRATco Veterinary Laboratory Services in Bogor, Indonesia, and ranged in weight from 150-200 g with aged 8-12 weeks. Rats were placed in individually ventilated cages with controlled temperatures for 12 hours day/night. Rats were fed a standard basal metabolism for rats with crude protein 18%, fat 6%, fiber 5,5%, and carbohydrate 44,2% ad libitum. Rats in each group were adapted for seven days.

Thirty-two male Sprague Dawley (SD) rats were used in the experiment, and they were randomly divided into 6 groups with four rats in each group. Negative control (NC): normal rats, Positive control (PC): bacterial-induced gingiva, Base cream group (BC): positive control + base cream, Comparative control group (CC): positive control + Kenalog cream, Sappan Wood Cream 1 (SWC1) group: positive control + applied with sappan cream 1 times daily in the morning, Sappan Wood Cream 2 (SWC2) group: positive control + applied with sappan cream 2 times daily in the morning and afternoon.

In the gingivitis induction, the bacteria used was *P. gingivalis* with a concentration of  $2 \times 10^8$  for each rat. The upper gingiva is drilled and then *P. gingivalis* bacteria are injected. The bacteria are injected into the gingivas using the Nichimate Stepper Injections. The volume of bacteria injected was 50  $\mu$ l with a ratio of 1:1. Induction was carried out for 14 days and every 3 days were injected until rats suffered positive gingivitis, characterized by bleeding, swelling, and darkly red gingiva tissue (Wang et al., 2022). After that, the rats were sacrificed for sample collection.

### ***Sample Collection***

The rat was terminated by injection of ketamine (Ikapharmindo) and xylazine (361453, Interchemie) intraperitoneally. Rats were sacrificed after being deeply anesthetized (Widowati et al., 2022). The rat's jaw was then removed and soaked in 10% Formaline fixation solution for two days. The rat jaws were then soaked again in a weak acid solution for the decalcification process for 10 days prior to analysis.

### ***Histopathological Examination***

Histopathological examination was performed using hematoxylin-eosin. The gingival sample was rinsed with 0.9% physiological NaCl and then fixation for three days using 10% formalin. The organs were dehydrated with graded alcohol (70%, 80%, 90%, 95%, 100%) for 2 hours. Subsequently, the organs were cleansed and dipped into the xylol while being continuously agitated. Then the organs were immersed in liquid paraffin at 60°C and preserved until they formed paraffin blocks. The paraffin block is sectioned using a microtome (Leica RM 2135 BioCut Rotary Microtome). The paraffin slices were

placed onto a slide plate and stained for further analysis (Gondokesumo et al., 2019; Widowati et al., 2022).

#### ***Collagen Proportion Examination by Masson Thricome***

The tissue was stained with MT after being submerged in Weigert's Iron Hematoxylin for 10 minutes, washed with distilled water, and then stained with a solution of Biebrich scarlet acid fuschin for 10-15 minutes. The tissue was then washed with distilled water and stained for 10-15 minutes with a phosphomolybdic-phosphotungstic acid solution. The tissue was dyed for 5-10 minutes with aniline blue solution, washed with Aquadest, and then soaked in 1% acetic acid solution. The tissue was washed with distilled water before to being submerged in 95% and 100% ethanol. A coverslip is placed over the preparation, which is then viewed under a microscope. The quantity of collagen proportions was assessed (Suvik and Effendy, 2012).

#### ***Immunohistochemistry Examination***

The Immunohistochemistry assay was carried out using the IHC kit (Elabscience, E-IR-R213) as a secondary antibody. The IHC slide contains gingival tissue were incubated in 3% H<sub>2</sub>O<sub>2</sub> (Elabscience, E-IR-R217C) for 10 minutes and added with Normal Goat Blocking Buffer (Elabscience, E-IR-R217A) at temperature 37°C. The NF-kB Monoclonal antibody (E-AB-22066) was added and incubated overnight at room temperature as a primary antibody. The target proteins were then visualized using Polyperoxidase-anti-Mouse/Rabbit IgG (Elabscience, E-IR-R217B) and DAB substrate was added and incubated for 20 minutes at room temperature. As a counterstaining agent, hematoxylin eosin was used. The stained tissues were analyzed using a Zeiss primostar microscope and photographed with a Lumenera infinity 1-3c (Hidayat et al., 2016; Pujimulyani et al., 2020).

#### ***qRT-PCR examination of TNF- $\alpha$ , IL-1 $\beta$ , IL-6, and P38 gene expression***

The isolation and purification of mRNAs was done using the Direct-zol RNA Miniprep Plus (Zymo, R2073). The mRNAs were reversed into cDNAs using SensiFAST cDNA synthesis kit (BIO-65054). SensiFAST SYBR

Green for DNA band visualization was used for gene optimization performed based on the manufacturer's directions. The Quantitative Polymerase Chain Reaction (qPCR) reaction mix was performed using SensiFAST SYBR No-ROX Kit (Meridian Bioscience, BIO-98005) and analyzed using Thermo Scientific's PikoReal Real-time PCR System (Thermo Fisher). The pre-incubation cycle was set at 95 °C for 5 minutes followed by 40 cycles of denaturation at 95 °C for 30 seconds. Then for each gene, the annealing process was carried out at 55, 53, 54, 58, 55°C with 40 cycles respectively. Pre-elongated and elongated at 72 °C for 1 minute (Hidayat et al., 2016; Pujimulyani et al., 2020; Widowati et al., 2018). The RNA Purity Concentration and primes used in this study was (Macrogen) as shown in Table 1 and Table 2 (Widowati et al., 2023).

### ***Ethical Approval***

This research was approved by the iRATCo Veterinary Laboratory with ethical number 4.2.012-3/KEHI//I/2023.

### ***Statistical Analysis***

The data analysis was based on One-way ANOVA and Tukey Post Hoc ( $p < 0.05$ ) using SPSS software (Widowati et al., 2016).

## **Results**

### ***The Bioactive Compound of Sappan Wood Extract by LC-MS/MS***

The chromatogram peak was used to define the different chemicals based on their respective molecular weights (**Figure 1 and 2**). Based on our study, there are 6 compounds present in sappan wood extract.

The compound detected in sappan wood extract was Chalcone (208.25 g/mol), ProtosappaninA (272.25 g/mol), Brazilin (286.28 g/mol), Sappanone B (302.28 g/mol), CaesalpinJ (316.30 g/mol), and Protosappanin B (304.29 g/mol).

### ***The effect of Sappan Wood Extract Cream on macro gingival rat model***

**Figure 3.** shows the effect of SWC on macro gingival morphology after bacterial induction. Based on the results showed that bacterial induction caused gingivitis in the rat model indicated by gingival bleeding and bacterial dental plaque in rat teeth. However, the group treated with SWC 1 and 2 showed macro improvements in the condition of the gingival. There was tissue repairment every day marked by the reduction of the reddish gingiva. It was able to treat the condition of the gingival tissue in rats, although plaque remained but did not cause dental calculus. These results demonstrate that treatment with SWC cream has anti-inflammatory activity that is able to treat the condition of the gingivas and teeth caused by gingivitis.

***The effect of Sappan wood extract cream towards inflammation and angiogenesis of gingival tissue in gingivitis rat model***

**Figure 4** shows the histopathology of inflammation and angiogenesis of gingival tissue in gingivitis model mice using Haematoxylin and Eosin (HE) staining. The bacterial induction caused the damage of the gingival tissue marked by the redness of gingiva (**Figure 4A**). The Inflammation and angiogenesis is shown in **Figure 4B**. The results of angiogenesis showed that in the PC and BC (base cream) the inflammation Score is increased and the angiogenesis score has no significant differences ( $P < 0.05$ ). Meanwhile, the sappan cream treatment group has a significantly differences ( $P < 0.05$ ) compared to positive control and base cream. Sappan cream treatment was effective in decreasing the inflammation score and in contrast increasing the angiogenesis Score. Therefore, the number of fibroblast cells, increasing new blood vessels and reducing the inflammatory process so it was able to help the rat gingival wound healing process. The best result was found in SWC2 groups with 2 times smears of sappan cream.

***The effect of Sappan wood extract cream towards NF-kB protein expression in gingivitis rat model***

**Figure 5** shows the effect of SWC towards NF-kB protein expression in the gingivitis rat model. The results showed that the gingival tissue treated with SWC is rejuvenated. When compared to SWC1 and

SWC2, a darker color indicates more NF-kB expression, which is a symptom of inflammation. The brown color intensity (light brown) is lower in the SWC2 treatments with 2 times smear (**Figure 5A**). Based on **Figure 5B**, the SWC2 treatment group, showed significant differences compared to positive control group ( $P < 0.05$ ). The NF-kB protein expression in SWC2 was  $23.43 \pm 1.70$  (**Figure 5B**). Indicating that sapan wood extract cream may downregulate NF-kB expression.

#### ***The effect of Sappan wood extract cream towards collagen density in gingivitis rat model***

Based on **Figure 6B**, it shows that the collagen score has been calculated based on the results of gingival histopathology images using the Masson Trichrome method. The collagen proportion was indicated in blue colour (**Figure 6A**). The graph shows that the SWC2 treatment group had the highest collagen score and was not significantly different from the negative control group (normal) ( $P < 0.05$ ). This indicates that SWC2 treatment is effective in increasing collagen in rat gingival tissue and has the potential to act as an anti-inflammatory agent.

#### ***The effect of Sappan wood extract cream towards IL-1 $\beta$ Gene Expression***

The IL-1 $\beta$  gene expression in gingivitis rats is shown in **Figure 7**. The bacterial induced only and base cream treatment groups showed a significant increase in IL-1 $\beta$  expression than negative controls (NC) and sappan wood cream groups (SWC1 and 2). Treatment with SWC with two topicals can significantly decrease the expression of the IL-1 $\beta$  gene expressions in gingivitis rat model ( $P < 0.05$ ). Treatment with SWC with two times smear was also the more effective compared to the one-time treatment significantly ( $P < 0.05$ ).

#### ***The effect of Sappan wood extract cream towards IL-6 Gene Expression***

The IL-6 gene expression after treated using SWC are displayed in **Figure 8**. Sappan cream can suppressed the IL-6 gene expression compared to the positive control group significantly ( $P < 0.05$ ). The SWC2 treatment has shown the lowest IL-6 gene expression ( $5.18 \pm 0.33$ ), that indicated have higher anti-

inflammatory activity. In contrast, the positive and base cream group has the highest IL-6 gene expressions.

#### ***The effect of Sappan wood extract cream towards p38 Gene Expression***

Based on qRT-PCR results showed that sappan cream (SWC) can decreased the expression of p38 gene (**Figure 9**). The SWC treatment with 2 times applied shows the lowest expressions of p38 ( $8.41 \pm 1.46$ ) significantly compared to the PC and BC groups ( $P < 0.05$ ). The sappan cream treatment with 2 times smear of SWC was also the more effective compared to the one-time smear treatment ( $P < 0.05$ ).

#### ***The effect of Sappan wood extract cream towards TNF- $\alpha$ Gene Expression***

The effect of sappan cream (SWC) towards TNF- $\alpha$  gene expression in gingivitis rat model is presented in **Figure 10**. The positive groups increased significantly the TNF- $\alpha$  gene expression compared to NC. The SWC groups treatment was found to statistically significant reduce the TNF- $\alpha$  expression compared to PC ( $P < 0.05$ ). Based on Figure 9, it shows that the gene expression for TNF- $\alpha$  in the SWC2 treatment group (SWC cream 2 times smear) showed the lowest TNF- $\alpha$  gene expression, with a value of  $7.86 \pm 0.43$ .

### **Discussion**

Periodontitis is a highly prevalent inflammatory disorders affecting people of all ages, however it is more common in the elderly (Xu et al., 2020). Periodontitis affects around 50% of adults globally and is most frequent in maturity. The frequency of severe forms is around 10%, with a large rise during the third and fourth decades (Könönen et al., 2019). Categorized as an infectious, long-lasting inflammatory condition affecting the tooth-supporting tissues (Kinane et al., 2017).

The origins and pathophysiology of periodontal disease are explained by the polymicrobial synergy and dysbiosis concept. In a healthy periodontium, bacterial populations are governed by the host's inflammatory response. However, following the development of resistant microorganism such as *Porphyromonas gingivalis*, there is a significant increase in the bacterial population and leading to

dysbiosis. This, in turn, results in the host's immune response becoming ineffective and damaging to the host's tissue (Hajishengallis & Lamont, 2016). In the healthy oral environment, *P. gingivalis* is found in small quantities and activates itself by consuming iron from the blood in swollen gingiva. *P. gingivalis*, once established, can actively contribute to the aggravation of periodontal disease. Because of the expensive expense of synthetic medications, researchers are more identifying bioactive chemicals of natural origin as therapeutics, one of the plant that has the potential to prevent periodontal disease is *Caesalpinia sappan*. Here, our study highlights the potential antiinflammatory activity of sapan wood extract.

The in vivo assessment was done by administering SWC for 14 days to rat that had been induced by *P. gingivalis* bacteria. Hematoxylin Eosin (HE) was done for assess the inflammation and angiogenesis score, Immunohistochemistry (IHC) was done for detecting protein expressions of Nf-KB, and Masson Trichome (MT) staining were used to analyze the collagen proportions in gingivitis rat models.

The gingiva is one of the oral mucosa most susceptible to injury or inflammation. According to our findings, the *P. gingivalis* bacterial induction causes the gingivitis. The occurrence of gingivitis is characterized by bleeding, edema and/or redness of the gingivas. This is in line with our findings that the rat macrogingiva experience reddish gingivas and bacterial plaque and there is repairment of the gingiva together with SWC treatment (Figure 3). The redness gingiva tissue is a sign of inflammation that is mediated to excessive injury (Zhang et al., 2018).

According to our results, the rat treated with SWC with 2 times smear had a lower score of inflammatory, angiogenesis, and expression of NF-kB protein. This finding is correspondent with previous study found that, Sappan wood extract is shown to have a pain-relieving effects, which can also assist in alleviating damaged caused by inflammation (Sa'diyah, 2018). In this study, sappan wood extract is prove to suppress NF-kB, the factors transcription that control the expression of genes associated with inflammation. The IHC assay revealed that sappan cream also can decrease the NF-kB

protein expressions. This means that, the transcription pathway for pro-inflammatory protein related to gingivitis could be suppressed.

As known before that sappan wood extract contains flavonoid polyphenolic compounds. These flavonoid compounds are thought to play a role in the ability to inhibit angiogenesis. The angiogenesis inhibitory action of flavonoids in the methanolic extract of sappan wood is thought to be through several mechanisms (Kolamala & Chava, 2023). In previous research it has been proven that flavonoid compounds, resveratrol and quercetin are able to inhibit aspects of angiogenesis (proliferation, migration of endothelial cells and formation of blood vessel pipes). The angiogenesis process is blocked by the COX-2 which plays a role in the angiogenesis process through the synthesis of prostaglandin (PG) (Kolamala & Chava, 2023). This is in line with our study result showed that the angiogenesis of cell were increased following SWC treatment.

The results of this investigation indicate that SWC increased angiogenesis and collagen proportion in the gingivitis rats induced by *P. gingivalis* induction. The evidence was shown in the collagen density in the gingiva of periodontitis rats treated with SWC, which were higher than that in periodontitis rats given Kenalog cream. Collagen breakdown is a critical physiologic mechanism involved in wound healing and tissue remodeling (Golub & Lee, 2020). The progression of disease is influenced by a number of associated molecular pathways, including growth factors, cytokines, and Matrix Metalloprotease (MMPs), as well as the regulators and inhibitors of these pathways (Binderman et al., 2017). The MMPs generated by inflammatory cells are the main host factor responsible for degradation of marginal gingiva collagen. In chronic periodontitis, the breakdown of the collagenous fibrous tissue of the marginal gingiva speeds up the resorption of alveolar bone (Tewtrakul et al., 2015). A study reported that sappan wood extract acts as a collagen receptor and involved in wound healing. An In vitro study reported that ethanolic sappan wood extract could increased type-I collagen production significantly by the L929 cells (Sobocki et al., 2022).

Gingivitis is a prolonged inflammatory responses that is primarily caused by gram-negative bacteria. The increase in the synthesis and release of different proinflammatory factors, resulting in excessive damage of gingivitis (Zhang et al., 2018). Proinflammatory factors including TNF  $\alpha$  , IL-1, IL-6 and INF- $\gamma$  have been found to be strongly associated with hemostasis, particularly in the development of dental plaques (Sobocki et al., 2022). *P. gingivalis* ' virulent component, a cysteine protease called gingipain, can cleave pro-MMP9 into its active form MMP9 (Zhou et al., 2015). Metalloproteinase (MMP) families degrade the extracellular matrix, basement membrane, and collagen degradation which promotes neoplastic cell invasion. *P. gingivalis* gingipains can also activate PAR2 and PAR4, resulting in I $\kappa$ B phosphorylation, NF- $\kappa$ B nuclear translocation, p38 activation and enhanced pro-MMP9 synthesis (Masi et al., 2011). Besides that, oxidative stress is taking a part in the emergence of periodontal diseases (Tamaki et al., 2014). Oxidative stress can lead to gingival damage due to the excessive production of ROS. This can occur through various pathways, including lipid peroxidation, DNA damage, protein degradation, oxidation of crucial enzymes, and the activation of proinflammatory cytokines (Vij et al., 2023; Zheng et al., 2023).

IL-1 $\beta$  is a pro-inflammatory cytokine that is taking a part in inflammatory mechanisms, this cytokine must be suppressed so that it can reduce the inflammatory process in the body (Rehman & Akash, 2016). TNF- $\alpha$  is regarded as the most important inflammatory cytokine, TNF- $\alpha$  inhibitors are now utilized as treatment medicines for certain disorders (Jang et al., 2021). IL-6 is a vital role in inflammation and its inhibitors are currently being employed as therapeutic medicines for a variety of disorders, with the potential to expand to more in the future, inhibiting IL-6 activity is useful against inflammatory disorders (Hirano, 2021).

SWC has the ability to suppress gene expressions cytokines in the inflammatory process. According to Sa'diyah (2018), secang wood extract reduces levels of pro-inflammatory cytokines IL-1 $\beta$ , IL-6, TNF- $\alpha$  and p38 in serum. These cytokines have a central role in chronic inflammation and tissue

damage. Secang wood extract exhibits anti-inflammatory properties through numerous mechanisms such as inhibition of p38 MAPK pathway and regulation of NF- $\kappa$ B which also inhibits the pro-inflammatory cytokines (Nirmal et al., 2015). This is in line with this study, revealed that sappan cream could decrease the TNF- $\alpha$ , IL-1 $\beta$ , IL-6, and p38 gene expressions compared to the positive controls. This result is in line with the IHC assay showed that Nf-KB protein expression was decreased following by the SWC treatment. The suppression of Nf-KB expression will affect the transcriptional pathway of TNF- $\alpha$ , IL-1 $\beta$ , IL-6, and p38 gene to be expressed in the cells. The proposed mechanism of sappan cream as anti-inflammatory agent is explained in **Figure 11**. However another methods should be added for validation and further experiments to explore the precise mechanism.

Overall, the sappan cream has the best effect on the animal tests compared to the basis cream even the comparative product. This is due to the biologically active compound of sappan wood extract in cream. The active compound contained in cream have a significant effect in the collagen density and angiogenesis, as shown by basis cream who don't have any active compound inside. In contrast, sappan cream contain bioactive compound such as flavonoid that enhanced the regeneration of gingival tissues, reduced inflammation in gingivitis. Further clinical study must be done to confirm its effect on human.

In conclusion, sappan wood extract cream has an anti-inflammatory activity with 2 times smear has the best results to suppress inflammation,, and decrease the expression of IL-1 $\beta$ , IL-6, TNF- $\alpha$  and p38 pro-inflammatory cytokines. In contrast it also has the ability to increase the collagen density and angiogenesis.

### **Acknowledgments**

The authors are thankful to the Maranatha Christian University for provided fund for this research (*Skema Percepatan Guru Besar*) with research grant Number 007/PRJ-GB/UKM/I/2023. We are also grateful to Aretha Medika Utama, Bandung, West Java, Indonesia, and iRATCo Veterinary Laboratory which supplied the laboratory facilities and and their assistance.

### ***Conflict of Interest***

The authors declare that there is no conflict of interest

### **Authors Contributions**

VKS, JJ, WW, RF designed the experiments; RF, DMM, DSH performed experiments and collected data; VKS, NSMD, DMM, DSH analyzed and interpretation of the data; VKS, WW, NSMD, JJ, RF drafted the manuscript; VKS, WW, RF Analyzed the data, Supervised and directed the study; VKS, JJ, WW, NSMD, RF, DMM, and DSH Final approved of the version to be published.

### **Funding**

This research is funded by Maranatha Christian University (*Skema Percepatan Guru Besar*) with research grant Number 007/PRJ-GB/UKM/I/2023.

### **Data Availability**

The data that support the findings of this study are not openly available due to reasons of sensitivity and are available from the corresponding author upon reasonable request.

### **References**

- Binderman, I., Gadban, N., and Yaffe, A. 2017. Extracellular ATP is a key modulator of alveolar bone loss in periodontitis. *Arch. Oral Biol.* 81, 131–135.
- Cardoso, E. M., Reis, C., and Manzanares-Céspedes, M. C. 2018. Chronic periodontitis, inflammatory cytokines, and interrelationship with other chronic diseases. *Postgrad. Med.* 130(1), 98–104.
- Cheng, R., Wu, Z., Li, M., Shao, M., and Hu, T. 2020. Interleukin-1 $\beta$  is a potential therapeutic target for periodontitis: a narrative review. *Int. J. Oral Sci.* 12(1), 1–9.
- Ghosh, P. K., and Gaba, A. 2013. Phyto-extracts in wound healing. *Journal of Pharmacy and Pharm. Sci.* 16(5), 760–820.
- Golub, L. M., and Lee, H.-M. 2020. Periodontal therapeutics Current host-modulation agents and future directions. *Periodontol.* 2000. 82, 186–204.

- Gondokesumo, M. E., Pardjianto, B., Handono, K., Sumitro, S. B., Widowati, W., Wismandanu, O., Hartady, T., Rosdianto, A. M., Goenawan, H., Lesmana, R., Wathoni, N., and Supratman, U. 2019. *Garcinia mangostana* extract enhances skin epithelialization in rat induced burn injury. *Pak. Vet. J.* 39(3), 365–370.
- Hajishengallis, G., and Lamont, R. J. 2016. The polymicrobial synergy and dysbiosis model of periodontal disease pathogenesis. *The Human Microbiota and Chronic Disease: Dysbiosis as a Cause of Human Pathology*, Eds., Nibali, L. and Henderson, B. Willey Publisher. pp: 227–242.
- Hanna, J. R., and Giacobelli, J. A. 1997. A review of wound healing and wound dressing products. *J. Foot. Ankle. Surg.* 36(1), 2–14.
- Hidayat, M., Prahastuti, S., Fauziah, N., Maesaroh, M., Balqis, B., and Widowati, W. 2016. Modulation of adipogenesis-related gene expression by ethanol extracts of detam 1 soybean and jati belanda leaf in 3T3-L1 cells. *Bangladesh J. Pharmacol.* 11(3), 697–702.
- Hirano, T. 2021. IL-6 in inflammation, autoimmunity and cancer. *Int. Immunol.* 33(3), 127–148.
- Jang, D. I., Lee, A. H., Shin, H. Y., Song, H. R., Park, J. H., Kang, T. B., Lee, S. R., and Yang, S. H. 2021. The role of tumor necrosis factor alpha (Tnf- $\alpha$ ) in autoimmune disease and current tnf- $\alpha$  inhibitors in therapeutics. *Int. J. Mol. Sci.* 22(5), 1–16.
- Jayusman, P. A., Nasruddin, N. S., Mahamad Apanidi, N. I., Ibrahim, N., and Budin, S. B. 2022. Therapeutic Potential of Polyphenol and Nanoparticles Mediated Delivery in Periodontal Inflammation: A Review of Current Trends and Future Perspectives. *Front. Pharmacol.* 13(7), 1–17.
- Jia, L., Han, N., Du, J., Guo, L., Luo, Z., and Liu, Y. 2019. Pathogenesis of important virulence factors of *Porphyromonas gingivalis* via toll-like receptors. *Front. cell. infect. microbiol.* 9(7), 1–14.
- Kinane, D. F., Stathopoulou, P. G., and Papapanou, P. N. 2017. Periodontal diseases. *Nat. Rev. Dis.* 3(6), 1–14.

- Kolamala, N., and Chava, V. K. 2023. Collagen Degradation in Periodontal Health and Disease: A Brief Review. *Essent. Dent.* 2(1), 39–44.
- Könönen, E., Gursoy, M., and Gursoy, U. K. 2019. Periodontitis: A multifaceted disease of tooth-supporting tissues. *J. Clin. Med.* 8(8), 1135.
- Lee, J. K., and Kim, N. J. 2017. Recent advances in the inhibition of p38 MAPK as a potential strategy for the treatment of Alzheimer's disease. *Molecules*, 22(8), 1287.
- Liu, T., Zhang, L., Joo, D., and Sun, S. C. 2017. NF- $\kappa$ B signaling in inflammation. *Signal Transduct. Target. Ther.* 2(3), 1-9.
- Masi, S., Salpea, K. D., Li, K., Parkar, M., Nibali, L., Donos, N., Patel, K., Taddei, S., Deanfield, J. E., D'Aiuto, F., and Humphries, S. E. 2011. Oxidative stress, chronic inflammation, and telomere length in patients with periodontitis. *Free Radic. Biol. Med.* 50(6), 730–735.
- Naruishi, K. 2022. Biological Roles of Fibroblasts in Periodontal Diseases. *Cells*, 11(21), 3345.
- Nastri, L., De Rosa, A., De Gregorio, V., Grassia, V., and Donnarumma, G. 2019. A New Controlled-Release Material Containing Metronidazole and Doxycycline for the Treatment of Periodontal and Peri-Implant Diseases: Formulation and in Vitro Testing. *Int. J. Dent.* 2019(1), 9374607.
- Nirmal, N. P., Rajput, M. S., Prasad, R. G. S. V., and Ahmad, M. 2015. Brazilin from *Caesalpinia sappan* heartwood and its pharmacological activities: A review. *Asian Pac. J. Trop. Med.* 8(6), 421–430.
- Papathanasiou, E., Conti, P., Carinci, F., Lauritano, D., and Theoharides, T. C. 2020. IL-1 Superfamily Members and Periodontal Diseases. *J. Dent. Res.* 99(13), 1425–1434.
- Pujimulyani, D., Yulianto, W. A., Setyowati, A., Arumwardana, S., Kusuma, H. S. W., Sholihah, I. A., Rizal, R., Widowati, W., and Maruf, A. 2020. Hypoglycemic activity of curcuma mangga val. Extract via modulation of GLUT4 and ppar- $\gamma$  mrna expression in 3T3-L1 adipocytes. *J. Exp. Pharmacol.* 12, 363–369.
- Rehman, K., and Akash, M. S. H. 2016. Mechanisms of inflammatory responses and development of

- insulin resistance: How are they interlinked?. *J. Biomed. Sci.* 23(1), 1–18.
- Sa'diyah, I. F. 2018. Efek Anti-Angiogenesis Ekstrak Kayu Secang Sebagai Terapi Adjuvant pada Diabetes Retinopati. *Saint. Med.* 14(2), 114–118.
- Sardi, N. W. A., Adnyasari, N. L. P. S. M., and Ekasari, N. P. R. B. 2023. Gel Extraction Of Earthworms (*Lumbricus Rubellus*) To The Number Of Fibrobal Cells In Male Wistar Rats (*Rattus Norvegicus*) Gingival Wound Healing. *IJKG.* 19(1), 34–42.
- Sayuti, N. A. 2015. Formulasi dan Uji Stabilitas Fisik Sediaan Gel Ekstrak Daun Ketepeng Cina (*Cassia alata* L.). *JKI.* 5(2), 74–82.
- Sobocki, B. K., Basset, C. A., and Ka, K. (2022). Molecular Mechanisms Leading from Periodontal Disease to Cancer. *Int. J. Mol. Sci.* 23(2), 970.
- Suvik and Effendy. 2012. the Use of Modified Masson ' S Trichrome Staining in Collagen Evaluation in Wound Healing Study. *Malaysian J. Vet. Res.* 3(1), 39–47.
- Tamaki, N., Cristina Orihuela-Campos, R., Inagaki, Y., Fukui, M., Nagata, T., and Ito, H. O. 2014. Resveratrol improves oxidative stress and prevents the progression of periodontitis via the activation of the Sirt1/AMPK and the Nrf2/antioxidant defense pathways in a rat periodontitis model. *Free Radic. Biol. Med.* 75, 222–229.
- Tewtrakul, S., Tungcharoen, P., Sudsai, T., Karalai, C., Ponglimanont, C., and Yodsaoue, O. 2015. Antiinflammatory and wound healing effects of *Caesalpinia sappan* L. *Phytother. Res.* 29(6), 850–856.
- Vaithiyam, V., Jadon, R. S., Ray, A., Manchanda, S., Meena, V. P., Ranjan, P., and Vikram, N. K. 2019. Metronidazole induced encephalopathy: A rare side effect with a common drug. *Indian J. Radiol. Imaging.* 29, 431–434.
- Vij, T., Anil, P. P., Shams, R., Dash, K. K., Kalsi, R., Pandey, V. K., Harsányi, E., Kovács, B., and Shaikh, A. M. 2023. A Comprehensive Review on Bioactive Compounds Found in *Caesalpinia sappan*.

Molecules, 28(17), 6247.

Wang, Y., Fan, Q., Xu, Y., Zeng, F., Liu, X., Zhao, D., Zhang, L., and Bai, G. 2022. Effect of *Eucommia* water extract on gingivitis and periodontitis in experimental rats. *BMC Oral Health*, 22(1), 1–9.

Widowati, W. 2011. Uji Fitokimia dan Potensi Antioksidan Ekstrak Etanol Kayu Secang (*Caesalpinia sappan* L.). *JMH*. 11(65), 23–31.

Widowati, W., Afifah, E., Mozef, T., Sandra, F., Rizal, R., Amalia, A., Arinta, Y., Bachtiar, I., and Murti, H. 2018. Effects of insulin-like growth factor-induced Wharton jelly mesenchymal stem cells toward chondrogenesis in an osteoarthritis model. *Iran. J. Basic Med. Sci.* 21(7), 745–752.

Widowati, W., Darsono, L., Lucianus, J., Setiabudi, E., Susang Obeng, S., Stefani, S., Wahyudianingsih, R., Reynaldo Tandibua, K., Gunawan, R., Riski Wijayanti, C., Novianto, A., Sari Widya Kusuma, H., and Rizal, R. 2023. Butterfly pea flower (*Clitoria ternatea* L.) extract displayed antidiabetic effect through antioxidant, anti-inflammatory, lower hepatic GSK-3 $\beta$ , and pancreatic glycogen on Diabetes Mellitus and dyslipidemia rat. *J. King Saud Univ. Sci.* 35(4), 102579.

Widowati, W., Darsono, L., Suherman, J., Fauziah, N., Maesaroh, M., and Erawijantari, P. 2016. Anti-inflammatory effect of mangosteen (*Garcinia mangostana* L.) peel extract and its compounds in lps-induced raw264.7 cells. *Nat. Prod. Sci.* 22(3), 147–153.

Widowati, W., Wargasetia, T., Rahardja, F., Gunanegara, R., Priyandoko, D., Gondokesumo, M., Afifah, E., Wijayanti, C., and Rizal, R. 2022. Human Wharton's jelly mesenchymal stem cells inhibit cytokine storm in acute respiratory distress syndrome in a rat model. *Asian Pac. J. Trop. Biomed.* 12(8), 343–350.

Xu, W., Zhou, W., Wang, H., and Liang, S. 2020. Roles of *Porphyromonas gingivalis* and its virulence factors in periodontitis. *Adv. Protein Chem. Struc. Biol.* 120, 45-84.

Yang, D., Zhongzhi, Q., Yanze, L., Guoping, Z., Yong, P., and Peigen, X. 2010. New collection of crude drugs in Chinese Pharmacopoeia 2010 I. *Callicarpa* Linn. And related items. *Chin. Herb. Med.*,

2(4), 272–288.

Zhang, F., Geng, Y., Zhao, H., Wang, H., Zhang, Y., Li, D., Bian, B., and Yang, H. 2018. Effects of Huanglian Jiedu Decoration in Rat Gingivitis. *Evid. Based. Complement. Alternat. Med.* 2018(1), 8249013.

Zheng, X., Chen, J., Rao, N., Yang, C., Liu, J., Zhang, J., and Li, Y. 2023. The Effect of an Extract of Sappanwood, Protosappanin A and Protosappanin B on Osteogenesis in Periodontitis. *Front. Biosci. (Landmark Ed)*. 28(8), 172.

Zhou, Y., Sztukowska, M., Wang, Q., Inaba, H., Potempa, J., Scott, D. A., Wang, H., and Lamont, R. J. 2015. Noncanonical activation of  $\beta$ -catenin by *Porphyromonas gingivalis*. *Am. Soc. Microbiol.* 83(25), 3195–3203.

**Table 1 Concentration and purity of RNA**

Groups	RNA concentration (ng/ml)	RNA purity ( $\lambda$ 260/ $\lambda$ 280 nm)
NC (normal rats)	47.80	2.2111
PC (bacterial-induced rats)	54.88	2.2039
BC (positive control+base cream)	60.65	2.4406
CC (positive control+ Kenalog)	16.23	2.0450
SWC1 (positive control+ sappan cream 1 time daily)	22.88	2.0927
SWC2 (positive control+ sappan cream 2 times daily)	44.05	2.1380

**Table 2. Primer sequence of Rat IL-1 $\beta$ , IL-6, TNF- $\alpha$ , p38 and GADPH**

Gene	Sequence (5' - 3')	Product size (bp)	Annealing ( $^{\circ}$ C)	References
IL-1 $\beta$	5'- AGAATACCACTTGTGGCT-3' 5'- GTGTGATGTTCCCATTAGAC-3'	134	55	NM_031512.2
IL-6	5'- GAGCATTGGAAGTTGGGGTA-3' 5'- TGATGGATGCTTCCAAACTG-3'	230	53	NM_012589.2
TNF- $\alpha$	5'- GAAGACAATAACTGCACCCA-3' 5'- AACCCAAGTAACCCTTAAAGTC-3'	138	54	NM_012675.3
P38	5'-AGATAATGCGTCTGACGGG-3' 5'-AGGGGATTGGCACCAATAAA-3'	139	58	NM_031020.3
GADPH	5'- GAAGGTGAAGGTCGGAGT-3' 5'- GAAGATGGTGATGGGATTC-3'	172	55	NM_001357943.2

**Figure. 1. LC-MS/MS Spectrum of Sappan Wood Extract, mass spectra 50–500 m/z with negative ionisation.**

**Figure. 2. LC-MS/MS Spectrum of Sappan Wood Extract, mass spectra 50–500 m/z with positive ionization.**

**Figure 3. Macrogingival Morphology in gingivitis rat model treated with SWC**

\*NC: Negative Control, PC: Positive Control (Bacterial induced rats), BC: Base Cream (1 x oral base cream applied), CC: Comparative Control (1x kenalog cream applied), SWC1: Sappan Cream 1 (Positive control + 1 times cream applied) , SWC2: Sappan Cream 2 (Positive control + 2 times cream applied).

**Figure 4. Effect of Sappan wood extract cream towards histopathology of inflammation and angiogenesis of gingival tissue in gingivitis rat model (40x magnification)**

\*Data are presented as mean  $\pm$  standard deviation. For each treatment, testing was carried out in four repetitions. \*NC: Negative Control, PC: Positive Control (Bacterial induced rats), BC: Base Cream Control (1 x oral base cream applied), CC: Comparative Control (1x kenalog cream applied), SWC1: Sappan Cream 1 (Positive control + 1 times cream applied) , SWC2: Sappan Cream 2 (Positive control + 2 times cream applied). Different superscripts (a,ab,abc,bcd,c,cd) mark significant differences among treatment for based on Tukey HSD post hoc test ( $P < 0.05$ ). The black arrows (indicate the inflammation site on the gingival tissue). SC: stratun corneum; SS: stratum spinosum; SB: stratum basalis.

**Figure 5. Expression of NF-kB protein in gingiva using immunohistochemical methods**

\*Data are presented as mean  $\pm$  standard deviation. For each treatment, testing was carried out in four repetitions. \* NC: Negative Control, PC: Positive Control (Bacterial induced rats), BC: Base Cream Control (1 x oral base cream applied), CC: Comparative Control (1x kenalog cream applied), SWC1: Sappan Cream 1 (Positive control + 1 times cream applied) , SWC2: Sappan Cream 2 (Positive control + 2 times cream applied). Different superscripts (a,b,c,d) mark significant differences among treatments based on Tukey HSD post hoc test ( $P < 0.05$ ). The black arrows indicate the cell positive with NF-kB protein. SC: stratum corneum; SS: stratum spinosum; SB: stratum basalis.

**Figure 6. Collagen score on gingival histopathology using the Masson Trichrome method**

\*Data are presented as mean  $\pm$  standard deviation. For each treatment, testing was carried out in four repetitions. \* NC: Negative Control, PC: Positive Control (Bacterial induced rats), BC: Base Cream Control (1 x oral base cream applied), CC: Comparative Control (1x kenalog cream applied), SWC1: Sappan Cream 1 (Positive control + 1 times cream applied) , SWC2: Sappan Cream 2 (Positive control + 2 times cream applied). Different superscripts (a,b,c,d) mark significant differences among treatment based on Tukey HSD post hoc test ( $P < 0.05$ ). The black arrows indicate the collagen site on the gingival tissue. SC: stratum corneum; C: collagen; SB: stratum basalis.

**Figure 7. IL-1 $\beta$  gene expression in rat gingival tissue**

\*Data are presented as mean  $\pm$  standard deviation. For each treatment, testing was carried out in four repetitions. \* NC: Negative Control, PC: Positive Control (Bacterial induced rats), BC: Base Cream Control (1 x oral base cream applied), CC: Comparative Control (1x kenalog cream applied), SWC1: Sappan Cream 1 (Positive control + 1 times cream applied) , SWC2: Sappan Cream 2 (Positive control + 2 times cream applied). Different superscripts (a,b,c,d) mark significant differences based on Dunnet T3 test (P<0.05).

**Figure 8. IL-6 gene expression in rat gingival tissue**

\*Data are presented as mean  $\pm$  standard deviation. For each treatment, testing was carried out in four repetitions. \* NC: Negative Control, PC: Positive Control (Bacterial induced rats), BC: Base Cream Control (1 x oral base cream applied), CC: Comparative Control (1x kenalog cream applied), SWC1: Sappan Cream 1 (Positive control + 1 times cream applied) , SWC2: Sappan Cream 2 (Positive control + 2 times cream applied). Different superscripts (a,b,c,d) mark significant differences among treatment based on Tukey HSD post hoc test ( $P < 0.05$ )

**Figure 9. p38 gene expression in rat gingival tissue**

\*Data are presented as mean  $\pm$  standard deviation. For each treatment, testing was carried out in four repetitions. \* NC: Negative Control, PC: Positive Control (Bacterial induced rats), BC: Base Cream Control (1 x oral base cream applied), CC: Comparative Control (1x kenalog cream applied), SWC1: Sappan Cream 1 (Positive control + 1 times cream applied) , SWC2: Sappan Cream 2 (Positive control + 2 times cream applied). Different superscripts (a,b,c,d) mark significant differences among treatments based on Dunnet T3 test ( $P < 0.05$ ).

**Figure 10. TNF- $\alpha$  gene expression in rat gingival tissue**

\*Data are presented as mean  $\pm$  standard deviation. For each treatment, testing was carried out in four repetitions. \* NC: Negative Control, PC: Positive Control (Bacterial induced rats), BC: Base Cream Control (1 x oral base cream applied), CC: Comparative Control (1x kenalog cream applied), SWC1: Sappan Cream 1 (Positive control + 1 times cream applied) , SWC2: Sappan Cream 2 (Positive control + 2 times cream applied). Different superscripts (a,b,c,d) mark significant differences among treatment based Dunnet T3 test.

## **Figure 11. Proposed Mechanism of Sappan Cream in reducing the anti-inflammatory markers of Gingivitis Rat Models**

\**P. gingivalis* injection caused gingival inflammation. It is activated Nf-KB and p38 pathway and increase the pro-MMP9 production. Nf-KB pathway also induce the inflammation by producing the of pro-inflammatory cytokines expressions. The sappan cream treatment could reduce the inflammatory markers evaluated by RT-PCR and Immunohistochemistry.

### **Bukti melakukan review yang pertama (20 Juli 2024)**

=> Reviewer # 1

In the authors' manuscript, sappan cream was used to enhance the regeneration of gum tissue and reduce the inflammatory response to gingivitis. The manuscript is of average quality and the authors need to make a lot of effort if they want to be published in this journal.

1. In the manuscript, SWC exerts anti-inflammatory effects by inhibiting PAR2 and PAR4 activated NF-KB pathway and inflammatory factors, but the authors only illustrated this by qPCR of inflammatory factors and immunohistochemistry of NF-KB. These validation methods are single and the results are not solid enough, and experiments such as WB should be added for validation, while further experiments can be added to explore the mechanism.
2. This manuscript was lacking in drug safety validation; does the drug cause other adverse effects in mice? Evidence of this needs to be demonstrated with routine blood tests and liver and kidney function tests in mice, if possible.
3. In the Materials and Methods section, 32 mice were divided into 8 groups, but in the Results section, the groups were divided into 6, which was contradictory and made the authenticity of the article doubtful.
4. Some of the bar charts in the article only analyzed the trend and did not illustrate the statistical significance, e.g., in 5B, the SWC2-treated group showed a significant decrease in NF-kB protein expression compared with the positive control group, but did not illustrate the statistical significance. Statistical analysis should be added and presented in each bar graph.
5. The authors need to do a better job of verifying their work; the description of the figure in Results 4 does not match the figure legends; Figure 7 has incomplete figure legends.
6. Figure 7 is not sufficiently clear and should be adjusted.

## Bukti konfirmasi submit revisi pertama yang telah direvisi (03 Agustus 2024)



Open Veterinary Journal

Dari: noreply@ejmanager.com

Kepada: vinnakurniawati@yahoo.co.id



Sab, 3 Agu jam 21.32 ★

Dear Vinna Kurniawati,

Your REVISED ARTICLE entitled **Anti-Inflammatory Activity of Sappan Wood Extract Cream (Caesalpinia sappan) in Porphyromonas gingivalis-induced Gingivitis Rats Models** (Mns No:OVJ-2024-05-154) has been received by **Open Veterinary Journal**.

You could follow status of your manuscript by login to your author account at [www.ejmanager.com](http://www.ejmanager.com).

Thank you for submitting your REVISED version of your article.

Best regards,

Editor  
Open Veterinary Journal  
<http://www.openveterinaryjournal.com>

## Bukti melakukan review yang kedua (09 Agustus 2024)



Open Veterinary Journal

Dari: noreply@ejmanager.com

Kepada: vinnakurniawati@yahoo.co.id



Jum, 9 Agu jam 15.43 ★

Dear Vinna Kurniawati,

The revisions for your manuscript titled **-Anti-Inflammatory Activity of Sappan Wood Extract Cream (Caesalpinia sappan) in Porphyromonas gingivalis-induced Gingivitis Rats Models-** and manuscript number (OVJ-2024-05-154) was reviewed by Editorial Board of Open Veterinary Journal and decided that the following revisions should be done:

- Editorial Comments:

Re-write the References list as per OVJ format. For example: Journals names should be abbreviated as that in PubMed. See Instructions for Authors or see other published articles in OVJ as a guide for References writing.

You should send your revised manuscript by journal Submit Article page.

Sincerely yours,

Ibrahim Eldaghayes, Prof.  
Editor-in-Chief  
Open Veterinary Journal  
[ibrahim.eldaghayes@vetmed.edu.ly](mailto:ibrahim.eldaghayes@vetmed.edu.ly)  
<http://www.ejmanager.com/my/ovj>

## Bukti konfirmasi submit artikel yang telah revisi kedua (15 Agustus 2024)



Open Veterinary Journal

Dari: noreply@ejmanager.com

Kepada: vinnakurniawati@yahoo.co.id



Kam, 15 Agu jam 10.23



Dear Vinna Kurniawati,

Your REVISED ARTICLE entitled **Anti-Inflammatory Activity of Sappan Wood Extract Cream (Caesalpinia sappan) in Porphyromonas gingivalis-induced Gingivitis Rats Models** (Mns No:OVJ-2024-05-154) has been received by **Open Veterinary Journal**.

You could follow status of your manuscript by login to your author account at [www.ejmanager.com](http://www.ejmanager.com).

Thank you for submitting your REVISED version of your article.

Best regards,

Editor

Open Veterinary Journal

<http://www.openveterinaryjournal.com>

## Bukti konfirmasi artikel diterima (19 Agustus 2024)

August 19, 2024

Dear Vinna Kurniawati,

I am pleased to inform you that your manuscript titled as "Anti-Inflammatory Activity of Sappan Wood Extract Cream (Caesalpinia sappan) in Porphyromonas gingivalis-induced Gingivitis Rats Models" (Manuscript Number: OVJ-2024-05-154) was accepted for publication in the Open Veterinary Journal.

A Final PDF draft will be prepared in the first three weeks of September and sent to you for a revision before publication.

Your manuscript will be published online in the ninth issue by the end of September 2024.

You may login to your author account page and visit accepted articles section in order to get official/formal acceptance letter as PDF.

Thank you for your interest in Open Veterinary Journal.

Sincerely yours,

*Ibrahim Eldaghayes, Prof.*

*Editor-in-Chief*

*Open Veterinary Journal*

*ibrahim.eldaghayes@vetmed.edu.ly*

## Bukti Galery Proof Manuscript (21 September 2024)

OVJ Open Veterinary Journal

Anna Kusawati - LDGOUT  
Personal Messages(0)

Home · Guide for Authors · Switch Journal · Submit New Manuscript · Status of my Articles

For your questions send email to [ibrahim.eltaghayest@gmail.com](mailto:ibrahim.eltaghayest@gmail.com)

### Author Center

- [My Manuscripts at the Editorial Office](#) (0)
- [My Manuscripts under Review](#) (0)
- [My Incomplete Submissions](#) (0)
- [Manuscripts Waiting for my Review](#) (0)
- [My Accepted Manuscripts \(Including waiting proofs\)](#) (0)
- [Charges / Fees / Payments](#) (Waiting: 0)
- [My Withdrawn or Transferred Manuscripts](#) (0)
- [My Deleted Manuscripts](#) (0)
- [My Rejected Manuscripts](#) (0)
- [My Manuscripts in Showroom](#) (0)  
Let know who of journals see your article(s)
- [My Published Manuscripts](#) (1)

Widhacsi / Enny  
Ranoel

## Bukti Publikasi Online Artikel (30 September 2024)

Google Cendekia

Anti-inflammatory Activity of Sappan Wood Extract Cream (Caesalpinia sappi

Artikel 1 hasil (0,07 dik)

Kapan saja  
Sejak 2024  
Sejak 2023  
Sejak 2020  
Rentang khusus...

Urutkan menurut relevansi  
Urutkan menurut tanggal

Semua jenis  
Artikel kajian

sertakan paten  
 mencakup kutipan  
 Buat notifikasi

Anti-inflammatory activity of sappan wood extract cream (Caesalpinia sappan) in porphyromonas gingivalis-induced gingivitis rats models  
VK Sugiawan, J Jeffrey, W Widowati... - Open Veterinary ..., 2024 - openveterinaryjournal.com  
Background: Gingivitis is a dysbiotic condition characterized by persistent inflammation caused by a disease-associated multispecies bacterial population that has established itself in the subgingival region. Aim: The objectives of this study is to analyze the anti-inflammatory activity of sappan wood extract cream (SWC) in Porphyromonas gingivalis induced-periodontitis rats model. Methods: In this study, rats were infected to the gingiva with P. gingivalis as an inducer for gingivitis rats model for 14 days. The SWC were treated ...  
☆ Simpan 📄 Kutip Artikel terkait 🔗