



सिरीएसआईआर  
CSIR  
भारत का नवाचार इंजन  
The Innovation Engine of India

A CSIR Publication

NIScPR  
National Institute of Science Communication and Policy Research  
सीएसआईआर-निस्र

Indian J Nat Prod Resour


June 2025

Vol. 16, No. 2, pp. 203-336

ISSN : 0976-0504 (print); ISSN 0976-0512 (online)

ijnpr@niscpr.res.in

# Indian Journal of Natural Products and Resources



<https://niscpr.res.in>

CSIR-National Institute of Science Communication and  
Policy Research

New Delhi, INDIA

Online Submission URL: <https://or.niscpr.res.in/index.php/IJNPR>



Prisca, Suzanne, Maribel, Sarah, Kristin, Catherine



Fouad S. Al-Sayed, Maged A. Al-Ghannay, Ali T. Zaki, Shams M. El-Mor, Hani A. Al-Sayid, Nasser A. Al-Sayid



As the business grows, the owner can add more staff and expand the range of services offered.



Yvonne Kucharski/El Seguros, Jeffrey Jeffrey, Philia Sazawa, Wahpis Wokoswa, Elmer Seguyaw, Haggisawon, Vic Assad, Timoth Barting Hagosaw Girmawakel, Rhod Acie

[illegible]

Rajal Manojkumar, Divya Hukla, Kiran Chaitan, V.K. Verma

Shirley Ann, 4 years old, living with mother, 38 years old.

Christian Schermann, Michaeland Huber-Nasse, Eke-Walsche, Jörn H. Dymowski, Wilfried Dier, Rainer  
Frost



The authors V. S. Shcherbakov, T. A. Kuznetsova, S. N.

Departamento de Matemática, Facultad de Ciencias Exactas, Universidad Nacional de Mar del Plata, Mar del Plata, Argentina



Shirley A. Korman, Ph.D.

[Show All Images](#) [»](#)(SIR) Article 28 does not have Article 28(1)(b) of the Charter of Fundamental Rights of the European Union.

Copyright Notice

Authors who publish with IJOPW agree that once published copyright of the article will be transferred to the publisher, with the work simultaneously licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/), that allows others to share the work with an acknowledgment of the work's authorship and initial publication in this journal.

Except where otherwise noted, the Articles on this site are licensed under [Creative Commons License: CC Attribution-ShareAlike 4.0 International](#)



## Editorial Team

### Scientific Editor

Pramila Majumdar, CSIR-NIScPR, India

### Chief Editor

FR Dr S. Ignacimuthu, S J  
Director and Research Supervisor  
Xavier Research Foundation, St Xavier's College  
Palayamkottai 627002, Tamil Nadu, India

### Executive Editor

Dr P. Pushpangadan  
Director General  
Amity Institute For Herbal and Biotech Products  
Development  
3 Ravi Nagar, Ambalamukku, Peroorkada P.O.,  
Thiruvananthapuram 695005, Kerala, India

### Subject Editors

Dr Sharad K Srivastava  
Pharmacognosy Division  
CSIR-National Botanical Research Institute  
Lucknow 226001, India

Dr Navin Kumar Rastogi  
Department of Food Engineering  
CSIR- Central Food Technological  
Research Institute  
Mysore 570020, Karnataka, India

Prof Jyoti Prakash Tamang  
Department of Microbiology  
Sikkim University, Tadong 737102,  
Gangtok, Sikkim, India

Dr Randhir Singh Dahiya  
Department of Pharmacology  
Central University of Punjab  
Ghudda 151401, Bathinda, Punjab, India

Dr Deepak Kumar Semwal  
Faculty of Biomedical Sciences  
Uttarakhand Ayurved University  
Dehradun 248001, Uttarakhand, India

Dr Konnala Naravana Sunil Kumar



SJR

SI

EPI

SCImago

SJR

Enter Journal Title, ISSN or Publisher Name

menu

←

Ads by Google

Ad options

Send feedback

Why this ad? ⓘ

## Indian Journal of Natural Products and Resources



<b>COUNTRY</b>	<b>SUBJECT AREA AND CATEGORY</b>
India	Agricultural and Biological Sciences
Universities and research institutions in India	— Agronomy and Crop Science
	— Food Science
Media Ranking in India	— Plant Science
<b>PUBLISHER</b>	<b>SJR 2024</b>
National Institute of Science Communication and Information Resources (NISCAIR)	<b>0.185</b> <b>Q4</b>
	<b>H-INDEX</b>
	<b>41</b>
<b>PUBLICATION TYPE</b>	<b>ISSN</b>
Journals	09760504, 09760512
<b>COVERAGE</b>	<b>INFORMATION</b>
2006-2024	<a href="#">Homepage</a>
	<a href="#">How to publish in this journal</a>
	<a href="mailto:ijnpr@niscair.res.in">ijnpr@niscair.res.in</a>

Red dragon fruit extract as a natural disclosing agent with antioxidant, antibacterial, and antibiofilm properties against *Streptococcus mutans*

Vinna Kurniawati Sugiaman<sup>1\*</sup>, Jeffrey Jeffrey<sup>2</sup>, Dahlia Sutanto<sup>1</sup>, Wahyu Widowati<sup>3</sup>, Dhanar Septyawan Hadiprasetyo<sup>4,5</sup>,  
Vini Ayuni<sup>5</sup>, Timbul Partogi Haposan Simorangkir<sup>6,7</sup> and Rizal Azis<sup>5,8</sup>

<sup>1</sup>Faculty of Dentistry, <sup>3</sup>Faculty of Medicine, Maranatha Christian University, Bandung 40164, Indonesia

<sup>2</sup>Faculty of Dentistry, <sup>4</sup>Faculty of Pharmacy, Universitas Jenderal Achmad Yani, Cimahi, 40531, Indonesia

<sup>5</sup>Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung 40163, Indonesia

<sup>6</sup>Faculty of Military Pharmacy, Indonesia Defense University, Bogor 16810, Indonesia

<sup>7</sup>Army Health Center Pharmacy Institute, Bandung 40111, Indonesia

<sup>8</sup>Biomedical Engineering, Department of Electrical Engineering, Faculty of Engineering, Universitas Indonesia, Depok 16425, Indonesia

Received 25 November 2024; revised received 16 July 2025; accepted 25 July 2025

Dental caries is a widespread health concern driven by poor oral hygiene and biofilm-forming bacteria *Streptococcus mutans*. While chemical disclosing agents effectively visualize dental plaque, their potential risks, such as mucosal irritation, highlight the need for safer, natural alternatives. Red dragon fruit (*Hylocereus polyrhizus*) contains bioactive compounds (betacyanin and phyllocactin) with antioxidant, antibacterial, and antibiofilm properties, making it a promising candidate. This study employed molecular docking, *in vitro* assays, and formulation development to evaluate its potential as a natural disclosing agent. Docking simulations assessed the binding of betanin and phyllocactin to *S. mutans* biofilm-associated proteins (SrtA and GbpC). The extract was prepared using 70% ethanol through maceration. Antioxidant activity was measured using the DPPH assay; antibacterial and antibiofilm effects were evaluated through disc diffusion, MIC, MBC, and biofilm inhibition tests. Molecular docking showed strong binding affinities (−8.8 to −12.2 kcal/mol). The extract demonstrated antioxidant activity (IC<sub>50</sub>: 158.43 µg/mL), antibacterial activity at 1000 µg/mL with a 5.43 mm inhibition zone, and antibiofilm activity up to 91.57%. The chewable jelly formulation exhibited suitable physical characteristics, though limited colour intensity remains a key challenge. Red dragon fruit extract showed potential as a natural disclosing agent for caries prevention in children, supported by its bioactive properties and docking results. Further studies are recommended to enhance colour intensity and validate its efficacy through *in vivo* applications.

**Keywords:** Antibacterial, Antibiofilm, Antioxidant, Dental caries, Disclosing agent, Red dragon fruit, *Streptococcus mutans*

**IPC code: Int. cl. (2021.01)– A61K 6/00, A61K 36/00, A61K 131/00, A61P 31/00, A61P 39/00**

## Introduction

Tooth decay, or dental caries, remains a widespread chronic condition and a major global public health concern. The World Health Organization (WHO) reports that it impacts between 60% and 90% of school-aged children, as well as the majority of adults. The dental caries prevalence is elevating, especially in certain populations, due to factors such as poor eating habits, lack of oral hygiene, and limited access to dental care<sup>1</sup>. In Indonesia alone, a specific study conducted in East Jakarta found that around 71% 12-year-old school children had experienced caries incidence, on average, shows Decayed-Missing-Filled Teeth (DMFT) index of 2.278<sup>2</sup>.

*Streptococcus mutans* has long been identified as the main bacterium contributed in biofilm formation and the dental caries process. This bacterium metabolizes carbohydrates to produce acids and forms biofilms that adhere firmly to tooth surfaces. The acid produced causes enamel demineralization, thus increasing the risk of caries<sup>3</sup>. In addition, *S. mutans* interacts synergistically with other oral pathogens, further exacerbating the cariogenic process. Various studies have demonstrated that *S. mutans* biofilms are resistant to mechanical removal and host immune defences, making them a persistent source of infection unless adequately managed through preventive measures. Therefore, preventive measures through plaque detection and biofilm reduction are essential in controlling caries progression<sup>4</sup>.

\*Correspondent author  
Email: vinnakurniawati@yahoo.co.id



Biofilm formation in *S. mutans* is mediated by several virulence factors, notably Sortase A (SrtA) and Glucan-Binding Protein C (GbpC). SrtA is a membrane-anchored transpeptidase that attaches surface proteins, including GbpC, to the cell wall via the LPXTG motif<sup>6</sup>. GbpC contributes to biofilm stability by mediating binding to glucans and enhancing adherence to the tooth surface<sup>6-7</sup>. The interaction of these proteins facilitates initial colonization and biofilm maturation, increasing the cariogenic potential of *S. mutans*<sup>8</sup>. Targeting these proteins offers a promising approach for preventing dental plaque and caries.

Using toothpaste can reduce plaque formation, clean the tooth surface, provide a refreshing flavour, and eliminate bad breath. However, there is a lack of public awareness in controlling plaque, which can cause caries. Therefore, early detection of dental plaque using disclosing solution is essential to optimize plaque control and prevent caries. In modern dental practice, disclosing agents are commonly used visual aids to detect the presence of plaque on the tooth surface<sup>9</sup>. Disclosing agents allow dentists and patients to visually identify areas where plaque accumulates, enabling early intervention before conditions like dental caries<sup>10</sup>. However, currently available commercial disclosing agents generally contain synthetic chemicals such as erythrosine and basic fuchsin that can cause various side effects such as skin irritation, respiratory problems, thyroid dysfunction, carcinogenicity, and toxicity<sup>11-12</sup>.

Plant-derived disclosing agents may serve as an effective alternative. Red dragon fruit (*Hylocereus polyrhizus*) is recognized for having betacyanin compounds which makes it a potential candidate as a disclosing agent. Betacyanin from red dragon fruit has potential as a natural colourant and a powerful antioxidant that can protect tissues from free radical damage and reduce oxidative stress that causes inflammation, and is often associated with various oral diseases<sup>13</sup>. In addition, studies show that red dragon fruit extract has antibacterial and antibiofilm properties that are effective against *S. mutans*, thus providing the potential to prevent biofilm formation and plaque development on teeth<sup>14-15</sup>. Betanin, the prominent betacyanin, is known for its capability to scavenge free radicals and has also been shown to improve lipid profiles and exhibit anti-inflammatory effects<sup>11</sup>. Phyllocactin, another bioactive compound present in red dragon fruit, has been associated with

significant antioxidant activity that contributes to the fruit's potential as a natural preservative and health supplement<sup>15</sup>.

Several studies have demonstrated the antibacterial, antioxidant, and antibiofilm properties of red dragon fruit extract. However, research focusing on its use as a plaque disclosing agent remains limited. Furthermore, no studies have evaluated incorporating this extract into a child-friendly, chewable jelly that can visualize plaque and deliver therapeutic benefits. We hypothesized red dragon fruit extract will exhibit notable antibacterial and antibiofilm activity while serving as an effective plaque disclosing agent. This study is expected to provide an alternative disclosing agent based on natural ingredients that is safe, acceptable to children, and effective in preventing dental caries.

## Materials and Methods

### Molecular docking

A molecular docking study was conducted on two bioactive compounds from red dragon fruit extract—betanin and phyllocactin. The compounds were identified through the PubChem database. The target proteins, Sortase A (SrtA) and Glucan-Binding Protein C (GbpC), were retrieved from the RCSB Protein Data Bank. Afterwards, the proteins were prepared using AutoDock Tools, in which water molecules were eliminated and polar hydrogens were added. This resulted in the creation of AutoDock Vina-compatible PDBQT files. Docking simulations were performed in AutoDock Vina, based on the gridbox references in previous SrtA<sup>16</sup> and GbpC<sup>17</sup> studies. An exhaustiveness parameter of 16 and the number of output binding modes set to 5 were used in the docking simulation. For GbpC, the center of the grid box was positioned at  $x = 240.549$ ,  $y = -25.160$ , and  $z = 8.572$ , with a grid box size of  $26.00 \times 18.00 \times 23.25$  Å. For SrtA, the center point of the grid was configured at  $x = -37.000$ ,  $y = -18.200$ , and  $z = 4.300$ , with the same grid box size. The molecular docking results were then visualised using Discovery Studio to display the binding interactions, including hydrogen bonds and hydrophobic contacts.

### Red dragon fruit extraction

Maceration was used as the extraction method. The red dragon fruit (*Hylocereus polyrhizus*) extract powder was obtained from PT. Fathonah Amanah Shidiq Tabligh (PT. FAST), Indonesia. The extraction

process involved macerating the powder in 70% ethanol. After the first 24 hours, the liquid extract (filtrate) was collected, and fresh 70% ethanol was added to the remaining material (residue) for another 24 hours. The filtered extract was then mixed with lactose to form a powder. This extraction process continued until the filtrate became colourless and the solvent had fully evaporated<sup>18</sup>.

### DPPH scavenging activity

The test was conducted by first adding 50 µL of the sample into a well of a 96-well plate, followed by 200 µL of 0.077 mmol DPPH solution. A separate blank well was filled with 250 µL of 10% DMSO, while another well, used as a control, contained 250 µL of 0.077 mmol DPPH. The samples were incubated for 30 minutes in the dark at room temperature, followed by absorbance measurements at 517 nm. The percentage of free radical scavenging was calculated using the appropriate formula<sup>19</sup>:

Scavenging activity (%) =

$$\frac{\text{Control Absorbance} - \text{Sample Absorbance}}{\text{Control Absorbance}} \times 100$$

### Antimicrobial assay by disc diffusion methods

Disc diffusion was used to assess antimicrobial effects. The process began by using a sterile cotton swab to spread bacterial suspensions of *S. mutans* onto Mueller-Hinton Agar plates. The plates were then left to settle at room temperature for 30 minutes. Paper discs (6 mm in diameter) were infused with either the test sample, positive control, or negative control solutions and placed on the agar surface. The plates were incubated for 24 hours at 37°C aerobically. The inhibition zones around each disc were measured using a vernier caliper, and the average was calculated from three measurements<sup>20,21</sup>.

### Antimicrobial assay by MIC and MBC

The antimicrobial test began by determining the minimum inhibitory concentration (MIC) level utilizing the broth microdilution strategy. For MIC and minimum bactericidal concentration (MBC) levels, each bacterial inoculum was prepared using the standard colony suspension method. Inoculum was obtained by transferring colonies of *S. mutans* that had been incubated for 24 hours on MHA media into Mueller-Hinton Broth (MHB) (Himedia, M391). Freshly cultured colonies were used to ensure bacterial viability before preparing the inoculum.

Turbidity was adjusted to the McFarland 0.5 standard, corresponding to approximately  $1.5 \times 10^8$  CFU/mL. MIC values were determined using 96-well plates containing 100 µL of red dragon fruit extract (RDFE) at concentrations of 1000, 750, 500, 250, 125, 62.5, and 31.25 µg/mL, supplemented with 100 µL of each microbe. A positive control was used 0.2% chlorhexidine, whereas the negative control was 10% DMSO. The plate was put in a incubator and kept at 37°C for 24 hours. Using spectrophotometry, we evaluated the turbidity of the plates following the incubation period (Multiskan GO Thermo Scientific 51119300) in the wavelength range 500–600 nm. After measuring the uptake, we determined the MBC level by taking 100 µL and performing graded dilutions ranging from  $10^2$  to  $10^5$  in each MIC result well. Exactly 50 µL of the aqueous solution was cultured using the pour-plate technique on MHA agar. We incubated the plates at 37°C for 24 hours. The next day, we used a Funke Gerber 8500 colony counter to assess the number of bacterial colonies. Least bactericidal concentration (MBC) is characterized as the extract concentration that produces an inhibitory impact of 99%<sup>21</sup>.

### Biofilm assay

Biofilm assay was initiated by culturing bacteria in BHIB medium for 24 hours at 37°C until it reaches McFarland 0.5 standard turbidity ( $1.5 \times 10^8$  CFU/mL). About 200 µL bacterial suspension was administered to a 96-well microplate then incubated for biofilm formation at 37°C for 24 hours. After incubation, 100 µL of RDFE treatment (1000–31.25 µg/mL), negative control (DMSO 10%), and positive control (chlorhexidine 0.2%) were added and incubated at 37°C for 24 hours, then supernatant from the wells was discarded. The wells were washed thrice with sterile PBS to remove planktonic cells, then fixed for 15 minutes with absolute methanol and dried. Staining for 5 min was performed with 0.1% crystal violet, then washed with water, and dried. Finally, 33% glacial acetic acid 200 µL was administered to dissolve the dye, and the biofilm was determined for OD at 595 nm with a microplate reader<sup>21</sup>.

### Chewy jelly preparation

Extraction of red dragon fruit was carried out via maceration with 96% ethanol as the solvent. The liquid phase was evaporated under reduced pressure using a rotary evaporator, yielding a thickened

extract. Each necessary ingredient were prepared according to the formula to be made. Aquadest was heated to a boil, gelatin is added, and allowed to stand for about 15 minutes until the gelatin expands (mixture 1). Gelatin was chosen as the main gelling agent because it can create a soft, elastic texture that is perfect for children and works well with acidic ingredients. Its clear base enhances the visual appearance of colour, an important quality for a disclosing agent. Then glycerin was heated to almost boiling (mixture 2). Mixture 2 was added to mixture 1 and heated using a water bath for 45 minutes at 40°C. Then added sucralose, mannitol, citric acid, methylparaben, BHT, and heated on medium heat. Add the active substance in the form of extract and stir until homogeneous (mixture 3). Mixture 3 was removed from heat, sprinkled with lemon oil, and allowed to stand for 5 minutes before being poured into the mold. Mixture 3 was poured into the mold and stored for 24 hours at room temperature until it formed a stable gummy mass<sup>22</sup>.

#### Moisture and Ash content assay

Ash content was analyzed by placing 2–3 g of the sample into a pre-weighed silicate crucible and gradually heated until all carbon was removed. After cooling, the sample was weighed. If carbon remained, the sample was mixed with hot water and subsequently passed through ash-free filter paper. The residue and filter paper were then returned to the same crucible, while the filtrate was added back to the crucible, evaporated, and heated until a constant weight was achieved. The percentage of ash (w/w) was determined based on the initial weight of the sample<sup>23</sup>.

The moisture content measurement began by activating the moisture balance system. First, a 3 g extract sample was prepared. An aluminum plate was placed in the meter and zeroed by pressing the tare button. The sample was then transferred onto the aluminum plate. Upon pressing the start button, then the drying process began. The system automatically heated the sample until all moisture was removed, signaled by a beep and "test over" display. The final reading showed the water content as a percentage<sup>23</sup>.

#### Texture profile analysis

A texture analyzer (Brookfield CT3) with a 50-kg load cell was used to analyze the texture of jelly cubes. With two compression cycles, jelly cubes were

compressed to 40% of their initial height using a 5 cm diameter probe. The compression speed was maintained at 0.5 mm/s, with a 5-second interval between successive compressions. All measurements were performed at ambient room temperature. Key texture attributes were obtained by analyzing the resulting force-time curve, including hardness, cohesiveness, adhesiveness, springiness, gumminess, and chewiness<sup>24</sup>.

#### Physical stability test

##### Organoleptic test and weight uniformity test

Organoleptic tests evaluated the shape, aroma, taste, and colour of each gummy formulation. The colour appeared lighter than expected, which may indicate limited retention of red pigmentation from the dragon fruit extract. This outcome is likely due to pigment degradation during the heating process or interactions with other formulation components. A weight uniformity test was carried out by weighing 20 gummies one by one randomly from each formula made.

#### Homogeneity test

Two transparent glass objects were used under light to assess the homogeneity of a chewy jelly preparation. One glass slide was evenly coated with the chewy jelly sample, then a second glass slide was placed on top of it, aligning the two slides. A homogeneous chewy jelly formulation should not exhibit any visible coarse or clumped particles when viewed between the two glass slides<sup>25</sup>.

#### pH test

The pH of the chewy jelly preparation was tested using a calibrated pH meter. The standard neutral (pH 7.00) and acidic (pH 4.00) buffer solutions were used to calibrate the pH meter. Electrodes were rinsed appropriately using distilled water then dried between measurements. The pH of the chewy jelly preparation was measured in real time until the readings stabilized at a constant value. The pH measurement was conducted in triplicate, and the average value was calculated<sup>25</sup>.

#### Viscosity test

The viscosity of the chewy jelly preparation was measured using a Brookfield viscometer. The viscometer spindle was immersed in the chewy jelly sample placed in a glass beaker, and the speed was set



to 50 rpm. The viscosity value of the chewy jelly was monitored until a stable reading was displayed<sup>25</sup>.

### Statistical analysis

Statistical analysis was conducted with SPSS software version 20.0. One-way ANOVA then Tukey's HSD post hoc test was applied for data that showed normal distribution and homogeneity. Kruskal-Wallis and Mann-Whitney tests were used for data that did not meet these assumptions (abnormal data). Results with  $p$ -values  $< 0.05$  was considered statistically significant.

## Results

### Molecular docking

Molecular docking analysis demonstrated that betanin and phyllocactin successfully interact with GbpC (Fig. 1) and SrtA (Fig. 2), as shown in Table 1. Betanin exhibited the highest binding affinity with GbpC (-12.2 kcal/mol), forming three hydrogen bonds and one hydrophobic interaction. Phyllocactin

exhibited a lower binding affinity (-8.8 kcal/mol) with one hydrogen bond and two hydrophobic interactions. For SrtA, betanin demonstrated binding affinity of -9.5 kcal/mol with three hydrogen bonds and one hydrophobic interaction. In comparison, phyllocactin showed a slightly stronger binding affinity (-9.8 kcal/mol) with three hydrogen bonds. These results indicate that betanin and phyllocactin have strong potential as inhibitors of biofilm formation through their interactions with biofilm-associated proteins.

### DPPH antioxidant activity of red dragon fruit extract

The DPPH free radical scavenging activity of red dragon fruit extract percentage as presented in Fig. 3. The results indicated that free radical scavenging activity enhanced proportionally with increasing sample concentrations. The 200  $\mu\text{g/mL}$  concentration had the highest DPPH free radical scavenging activity in red dragon fruit extract, with a significant value ( $57.73 \pm 0.57$   $\mu\text{g/mL}$ ). The red dragon fruit




Fig. 1 — Molecular docking interactions of betanin (pink) and phyllocactin (green) with the GbpC protein. (Left) 3D visualization of the protein-ligand binding. (Right) 2D interaction diagram showing hydrogen bonds (green) and hydrophobic interactions (pink).




Fig. 2 — Molecular docking interactions of betanin (pink) and phyllocactin (green) with the SrtA protein. (Left) 3D visualization of the protein-ligand binding. (Right) 2D interaction diagram showing hydrogen bonds (green) and hydrophobic interactions (pink).

Table 1 — Binding affinity and amino acid interactions of betanin and phyllocactin with GbpC and SrtA proteins				
Protein	Compound	BA (kcal/mol)	Hydrogen bonds	Hydrophobic interaction
GbpC	Betanin	-12.2	TRP351, THR449, TRP451	ILE219
	Phyllocactin	-8.8	SER347	TRP451, VAL410
SrtA	Betanin	-9.5	THR180, GLY192, ARG197	VAL193
	Phyllocactin	-9.8	GLN172, SER116, PRO163	

extract Inhibitory Concentration (IC<sub>50</sub>) value is 158.43±3.67 µg/mL. This value is included in the weak category, but it can show that red dragon fruit has the potential as an antioxidant<sup>26</sup>.

**Antibacterial activity of red dragon fruit extract by the disc diffusion method**

In antibacterial testing employing the disc diffusion method, it is known that RDFE has antimicrobial activity against *S. mutans* bacteria characterized by the presence of an inhibition zone or clear zone formed (Fig. 4). RDFE at 200 µg/mL (Treatment IX) showed the greatest antimicrobial activity among the

extract groups, with an inhibition zone diameter of 5.43 mm, whereas RDFE at 25 µg/mL (Treatment VI) produced the smallest inhibition zone (1.14 mm). Statistical analysis indicated that the inhibition zone of Treatment VI was significantly different from that of the positive control (0.2% chlorhexidine) (p < 0.05).

**Antibacterial activity by minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) methods**

The percentage of growth (viability) and inhibition of *S. mutans* bacteria with treatment using RDFE as



presented in Fig. 5. The minimum inhibitory level (MIC) was in treatment VII (RDFE 250  $\mu\text{g/mL}$ ) with a percentage of 60.36%. While the MBC is owned by treatment X (RDFE 1000  $\mu\text{g/mL}$ ), with a percentage value of 92.34%. RDFE treatment showed a concentration-dependent inhibitory effect on the growth of *S. mutans* bacteria are directly proportional to the concentration level. The higher the concentration given, the lower the growth activity of *S. mutans* and the higher the inhibition. This is in line with the results of the calculation of the number of colonies contained in Table 2.




Fig. 3 — Effect of various concentrations of red dragon fruit extract on DPPH free radical scavenging activity.

\*Data are presented as mean  $\pm$  standard deviation. For each treatment, the assay was performed in three repetitions. Different superscripts (a,ab,b,c,d,e) mark significant differences between various concentrations of red dragon fruit extract ( $P < 0.05$ , Tukey's HSD test). 6.25, 12.50, 25.00, 50.00, 100.00, and 200.00  $\mu\text{g/mL}$  numbers on the x-axis indicate the variation of red dragon fruit extract concentration.




Fig. 5 — Effect of various concentrations of red dragon fruit extract by dilution method. a) Viability of *S. mutans*, and b) Inhibition of *S. mutans*.

\*Data are presented as mean  $\pm$  standard deviation. For each treatment, the test was performed in three repetitions. Different superscripts (a,b,c,d,e,f,g) mark significant differences between various concentrations of red dragon fruit extract ( $P < 0.05$ , Dunnett T3 test). Roman numerals I–III represent the controls (I: growth control, II: negative control/DMSO, III: positive control/0.2% chlorhexidine), while IV–X indicate RDFE at concentrations of 1000, 750, 500, 250, 125, 62.5, and 31.25  $\mu\text{g/mL}$ .

### Antibacterial activity of *S. mutans* biofilm inhibition

The biofilm inhibition study in Fig. 6, RDFE has an effect on reducing biofilm. The highest antibiofilm effect was in treatment X (RDFE 1000  $\mu\text{g/mL}$ ; 91.57%), with the lowest anti-biofilm effect being treatment IV (RDFE 31.25  $\mu\text{g/mL}$ ; 17.60%) against the *S. mutans* bacteria growth. The effect of biofilm inhibition on RDFE exhibits a concentration-dependent effect; the higher the treatment concentration, the higher the biofilm inhibition produced.

### Preparation of chewy jelly

Preparation of chewy jelly with the formulation used as in Table 3. Each necessary ingredient were prepared




Fig. 4 — Effect of various concentrations of red dragon fruit extract on *S. mutans* bacteria.

\*Data are presented as mean  $\pm$  standard deviation. For each treatment, the test was performed in three repetitions. Different superscripts (a,b,c,d,e,f) mark significant differences between various concentrations of red dragon fruit extract ( $P < 0.05$ , Tukey's HSD test). Treatment I: growth control, Treatment II: negative control (DMSO), Treatment III: positive control (0.2% chlorhexidine), and Treatments IV–X: RDFE at 6.25, 12.50, 25.00, 50.00, 100.00, and 200.00  $\mu\text{g/mL}$  respectively.

Table 2 — Colony counts of *S. mutans*

Sample	CFU/mL			Average
	P1	P2	P3	
Growth Control	TNTC	TNTC	TNTC	TNTC
Negative Control	TNTC	TNTC	TNTC	TNTC
Positive Control (Chlorhexidine 0.2%)	0	0	0	0
RDFE 1000 µg/mL	57 x 10 <sup>4</sup>	56 x 10 <sup>4</sup>	51 x 10 <sup>4</sup>	54.67 x 10 <sup>4</sup>
RDFE 750 µg/mL	65 x 10 <sup>4</sup>	68 x 10 <sup>4</sup>	75 x 10 <sup>4</sup>	69.33 x 10 <sup>4</sup>
RDFE 500 µg/mL	124 x 10 <sup>4</sup>	137 x 10 <sup>4</sup>	129 x 10 <sup>4</sup>	130.00 x 10 <sup>4</sup>
RDFE 250 µg/mL	192 x 10 <sup>4</sup>	163 x 10 <sup>4</sup>	187 x 10 <sup>4</sup>	180.67 x 10 <sup>4</sup>
RDFE 125 µg/mL	243 x 10 <sup>4</sup>	248 x 10 <sup>4</sup>	201 x 10 <sup>4</sup>	230.67 x 10 <sup>4</sup>
RDFE 62.5 µg/mL	TNTC	TNTC	TNTC	TNTC
RDFE 31.25 µg/mL	TNTC	TNTC	TNTC	TNTC

Notes: \*P= Repetition; TNTC = Too Numerous Too Count >250

Table 3 — Chewy jelly formulation design

Composition	Unit	Formulation 1	Formulation 2	Formulation 3
Red dragon fruit extract	Gram	0.5	1	2
Glycerin	G	16	20	25
Gelatin	G	7	5	6
Sucralose	G	0.015	0.015	0.015
Mannitol	G	1.5	1.5	1.5
Citric acid	G	0.15	0.15	0.15
BHT	G	0.006	0.006	0.006
Aquades	G	24.829	22.329	15.329
Corn oil	As needed	As needed	As needed	As needed




Fig. 6 — Effect of various concentrations of RDFE on biofilm growth of *S. mutans*.

\*Data are presented as mean  $\pm$  standard deviation. For each treatment, the assay was performed in three repetitions. Different superscripts (a,b,c,d,e,f,g,h,i) mark significant differences between various concentrations of RDFE ( $P < 0.05$ , Independent t-test). Roman numerals I–III represent the controls (I: growth control, II: negative control/DMSO, III: positive control/0.2% chlorhexidine), while IV–X correspond to RDFE at concentrations of 31.25, 62.5, 125, 250, 500, and 1000 µg/mL.

according to the formula. Aquadest was heated to a boil, gelatin was added, and allowed to stand for about 15 minutes until the gelatin expands (mixture 1).

Glycerin was heated to almost boiling (mixture 2). Mixture 2 was added to mixture 1 and heated using a water bath at 40°C for 45 minutes. Then, sucralose, mannitol, citric acid, and BHT were added and heated on medium heat. Add the active substance in the form of extract and stir until homogeneous (mixture 3). Mixture 3 was removed from heat, sprinkled with corn oil, and allowed to stand for 5 minutes before being poured into the mold. Mixture was poured on the mold tool and stored for 24 hours at room temperature until it forms a stable chewy jelly mass.

#### Evaluation of chewy jelly based on organoleptic test

The organoleptic test results were shown in Table 4. Chewy jelly had a consistency that was quite chewy. The aroma category of chewy jelly had a distinctive aroma but no flavour. This can be caused by the very small composition of sucralose. The physical appearance was a yellow colour for formulation I, yellow slightly pink in formulation II, and pink colour in formulation III. The colour change occurred because the composition of dragon fruit extract used differs.



Table 4 — Organoleptic test results

Organoleptic category	Formula I	Formula II	Formula III
Consistency	Quite chewy	Chewy	Chewy
Shape	Semisolid	Semisolid	Semisolid
Aroma	Dragon fruit aroma	Dragon fruit aroma	Dragon fruit aroma
Taste	Flavorless	Flavorless	Flavorless
Colour	Yellow	Slightly pinkish yellow	Light red

Table 5 — Texture analysis test results

Texture Profile Analyzer	Unit Value	Formula chewy jelly		
		I	II	III
Hardness	gForce	3,157,993	7,756,027	8,795,373
Fracturability	-	Undetected	Undetected	Undetected
Adhesiveness	g.sec	-0,0320	-95,975	-185,963
Springiness	%	38,303	0,9313	0,8730
Cohesiveness	%	0,8337	10,097	0,9950
Gumminess	gForce	1.094,2410	7,833,543	8,750,223
Chewiness	gForce	3.745,2050	7,351,033	7,656,663
Resilience	%	1.076	11,183	11,367

### Evaluation of chewy jelly based on moisture and ash content tests

Evaluation of the moisture and ash content of chewy jelly was also carried out. Moisture and ash content were analyzed on all three RDFE-based chewy jelly formulations. The moisture content in formulation I was 1.79%, formulation II had the highest moisture level at 8.11%, and formulation III had the lowest at 1.08%. Moisture content is a crucial factor that affects the quality and freshness of food ingredients. Too high moisture content can cause the product to become moist, accelerate damage, and increase the risk of growth of microorganisms such as fungi and bacteria. Regarding ash content, formulation I recorded 0.10%, formulation II had 0.08%, and formulation III showed the highest ash content at 0.12%. This difference could indicate variations in the content of mineral ingredients or other inorganic components in each formulation. Ash content was analyzed because it can indicate the mineral content in a product. The higher the ash content, the poorer the food quality, as it can indicate contamination or poor-quality raw materials. From the results of the water and ash content tests carried out on all formulations, the water content values obtained are still in the safe category and meet the SNI requirements because they are not more than 20%. Additionally, the ash content obtained was not more than 3% according to the provisions of the SNI 3547.2-2008 standard for the jelly candy category. Overall, the chewy jelly formula demonstrated

potential for further analysis to ensure that chewy jelly products meet the expected quality standards and are safe for consumption.

### Evaluation of chewy jelly based on texture analysis test

The texture profile analyzer results in Table 5, Formula I had the highest hardness value (3,157,993 gForce), while Formula II (7,756,027 gForce) had a lower hardness value and tends to be softer. The adhesive characteristic, namely the tendency for particles to stick together in formula I, had a very low adhesivity (-0.0320 g.sec), meaning that it was not too sticky and comfortable when consumed, and formula III was the formulation that had the stickiest adhesivity (-185,963 g.sec). In the Springiness category, or the ability to return to its original form, formula I had the highest springiness value (38,303%) compared to formulas II and III. Cohesion or attachment between particles in formula II had the highest cohesion (10,097%) and lower gumminess value (7,833,543 gForce), compared to formula I and III, that showed higher gumminess and chewiness values, thus making them chewier and requiring more effort to chew.

Overall, to be used as a disclosing agent, Formula II was a more suitable formula, as indicated by its hardness value (7,756,027 gForce), adhesivity (-95,975 g.sec), cohesion (10,097%), and gumminess (7,833,543 gForce) which were lower than formula III making it had a texture that was easily crushed and not too sticky so that it will not interfere with users.




Fig. 7 — Visualization of the chewy jelly characteristics over four weeks of testing: a) weight uniformity, b) pH, and c) viscosity.

### Evaluation of chewy jelly based on the physical stability test

Evaluation of the physical stability of chewy jelly was carried out for 4 weeks using formulas I and II based on the results of consideration of moisture content, ash, and characteristics according to TPA. The weight uniformity of the chewy jelly analyzed is shown in Fig. 7a, where formulation II had a weight loss over time. This can be due to the water content, which was higher than that of formula I, resulting in a decrease in the water content. The pH measurement of chewy jelly (Fig. 7b) tends not to be significantly different between formulas I and II. The texture analysis of the RDPE-based chewy jelly formulations included hardness, stickiness, and elasticity measurements. Formulation I exhibited a hardness

value of 144.67 gF, stickiness of –6.00 gF, and elasticity of 2.11 mm. Formulation II demonstrated the highest hardness at 270.00 gF, with stickiness of –5.20 gF and elasticity of 2.07 mm.

Meanwhile, formulation III showed a hardness of 145.33 gF, the lowest stickiness at –10.30 gF, and the highest elasticity at 2.63 mm. These results indicated that formulation II had the firmest texture, while formulation III was the most elastic and sticky. While the viscosity results of chewy jelly formula II showed an increase over time, the viscosity, which reached 1000 cP showed the chewiness of chewy jelly (Fig. 7c). The colour of the chewy jelly produced tends not to change during the 4 weeks of testing, which remains yellow and remains homogeneous in the absence of lumps. No microbes were found in chewy jelly.

In this study, because it was expected to be a disclosing agent that was able to provide colour to dental plaque that sticks to the teeth surface, remained a challenge because the fruit of dragon used was a powder extract that was less effective in providing colour. To turn it into a disclosing agent that can colour, further research is needed related to the addition of more effective dyes in accordance with national standards related to the use of dyes in food products, for example, with Erythrosine dye (FD&C Red No. 3), Code E: E127 (SNI 01-2891-1992). Nevertheless, the chewy jelly formulation produced can be an illustration of the manufacture of chewy jelly as a disclosing agent, but only in the future will it need to be improved with more concentrated dyes, for example, by using Erythrosine.

### Discussion

Molecular docking analyses of betanin and phylloactin demonstrated their potential as an inhibitor of biofilm formation through their interactions with GbpC and SrtA, two proteins associated with biofilm development. Among the two compounds, betanin showed consistently stronger binding interactions than phylloactin. This may be attributed to its ability to form more hydrogen bonds and favourable hydrophobic contacts, which likely contribute to a more stable ligand–protein complex. Differences in binding affinity and the interactions formed are important in the design of biofilm inhibitors, as compounds with lower binding affinity are often more effective in disrupting protein function<sup>27</sup>. Similar findings from other studies emphasise the importance of binding interactions in



the therapeutic efficacy of compounds targeting protein function<sup>28</sup>. Previous docking studies have shown that chlorhexidine, a widely used antibacterial agent in oral care, exhibits a binding affinity of approximately -6.29 kcal/mol against *S. mutans* SrtA<sup>29</sup>. In comparison, betanin and phyllocactin demonstrated stronger binding affinities (-8.8 to -12.2 kcal/mol). The ability of betanin and phyllocactin to effectively interact with key proteins associated with biofilms suggests they may act as biofilm inhibitors.

Red dragon fruit extract (RDFE) shows promising potential as a natural alternative for disclosing agents, attributed to its antioxidant, antibacterial, and antibiofilm properties. The DPPH assay results demonstrated moderate antioxidant activity, with an IC<sub>50</sub> of 158.43±3.67 µg/mL. Although this falls into a lower efficacy category<sup>26</sup>, RDFE's capacity to neutralize free radicals is still noteworthy, considering the oral cavity's susceptibility to oxidative stress, contributing to periodontal disease and caries. Research by Fidrianny *et al.*<sup>30</sup> also reported antioxidant activity in red dragon fruit extract with an IC<sub>50</sub> of 94.17 µg/mL. In addition, other studies on antioxidants in dragon fruit peel extracts show more promising results. Wahdaningsih *et al.*<sup>31</sup> evaluated antioxidant activity through DPPH scavenging assay on red dragon fruit peel with two isolates and showed stronger antioxidant activity with IC<sub>50</sub> of 2,952.14 and 25,635.95 µg/mL, respectively, while Fidrianny's *et al.*<sup>32</sup> study reported a DPPH scavenging activity IC<sub>50</sub> value was 46.51±4.72 µg/mL on red dragon fruit peel extract. The higher antioxidant activity in the peel extract can be attributed to the higher concentration of phenolic compounds and betacyanin, especially in the fruit peel tissue. The antioxidant potential is likely linked to the high concentration of betacyanin compounds in RDFE, as previously established by Ramli *et al.*<sup>13</sup>, suggesting its viability for supporting oral health through oxidative balance. The findings may thus contribute to broadening the antioxidant-focused approach in caries prevention.

Our antibacterial assessments through the disc diffusion method showed RDFE's efficacy in inhibiting *Streptococcus mutans* growth, with a peak inhibition zone of 5.43 mm at a concentration of 1000 µg/mL. This effect aligns with prior studies by Rahayu *et al.*<sup>14</sup>, which confirmed RDFE's antimicrobial capabilities against oral pathogens. Importantly, RDFE's concentration-dependent

inhibition of *S. mutans* (60.36% MIC at 250 µg/mL and 92.34% MBC at 1000 µg/mL) suggest potent antimicrobial profile that can be harnessed in low concentrations for practical applications. The significance of this effect, given *S. mutans* role in cariogenic biofilms, underscores RDFE's potential as a therapeutic adjunct in dental products targeting oral microbiome modulation.

Comparing our findings with previous research, Amalia *et al.*<sup>33</sup> demonstrated that red dragon fruit extract at 40 mg/mL, produced a 12.80 mm inhibition zone against *Staphylococcus aureus*, indicating strong antibacterial potential, with inhibition percentages reaching approximately 85.1%. Similarly, Hendra *et al.*<sup>34</sup> demonstrated that extracts from red dragon fruit peel also effectively inhibited *Bacillus subtilis*, with inhibition percentages reaching 100% at 20 µg/mL concentration. While these studies focused on different bacterial strains, they support our findings regarding RDFE's broad-spectrum antimicrobial potential. The slightly lower inhibition zone observed in our study with *S. mutans* might be attributed to the different cell wall structures and resistance mechanisms between gram-negative and gram-positive bacteria. Nevertheless, the relatively small inhibition zone observed at the highest RDFE concentration (5.43 mm) compared to chlorhexidine, may suggest a need for further optimization. Future studies may consider using higher concentrations of RDFE or exploring synergistic combinations with other natural compounds to enhance its antibacterial effect against *S. mutans*.

Biofilm refers to a community of microorganisms enclosed by an extracellular polymer matrix produced by microorganisms such as *S. mutans*<sup>35</sup>. Most compelling was RDFE's antibiofilm activity, which achieved substantial inhibition (91.57%) at a concentration of 1000 µg/mL. This attribute is crucial since biofilm resilience and adherence are primary factors in caries progression<sup>3</sup>. RDFE's inhibition of biofilm formation suggests that it could effectively prevent *S. mutans* adherence to enamel surfaces, thereby reducing the initial stages of biofilm development. This finding holds significant implications for caries management, especially in pediatric populations where non-invasive, safe interventions are prioritized. Research by Yong *et al.*<sup>36</sup>, which also evaluated the red dragon fruit extract antibiofilm activity, showed the percentage of inhibition against *Staphylococcus aureus* by 30-51%

and *Pseudomonas aeruginosa* by 23–42%. This shows that red dragon fruit extract has the potential to be an effective antibiofilm. Another comparison with other natural, such as green tea extract has been reported to inhibit *S. mutans* biofilms at concentrations around 3.1–12.5 mg/mL<sup>37</sup>, while essential oils from clove, thyme, and cinnamon have shown reductions of biofilm  $\geq 85$ –100% at 1–4 $\times$  MIC<sup>38</sup>. Importantly, the 91.57% inhibition by RDFE is comparable to these agents and aligns with the efficacy of chlorhexidine in many assays.

The development of RDFE-based chewy jelly as a delivery vehicle for disclosing agents demonstrated promising physical and organoleptic properties. Among the formulations tested, Formula II exhibited optimal hardness (775.60 gForce) and adhesivity (–9.60 g.sec), making it suitable for oral application. However, the colour intensity of the jelly requires improvement, as the current formulation may not achieve sufficient visual contrast for effective plaque disclosure. Addressing this limitation could involve integrating approved food colorants, such as Erythrosine (E127) as per SNI 01-2891-1992 standards, to enhance the visual contrast without compromising safety. Future research might also explore alternative natural pigments to maintain the formulation's appeal as a natural product.

Stability tests revealed that Formula II maintained acceptable pH (5.56 $\pm$ 0.12) and viscosity (1026.42 $\pm$ 47.43) over four weeks, though minor weight loss due to moisture changes suggests a need for further optimization. Enhancements in the formulation's moisture retention could improve its long-term stability, which is critical for product viability. Improving viscosity enhances the structural integrity of the jelly matrix, minimizing syneresis and phase separation, while better moisture retention reduces evaporation-related weight loss and texture degradation during storage. This contributes significantly to product stability and shelf life. Moreover, the absence of microbial growth across the testing period indicates effective preservation, supporting RDFE's potential for stable product formulation.

In addressing broader implications, RDFE-based products could mitigate health concerns associated with synthetic disclosing agents, as noted by Silva *et al.*<sup>11</sup> and Britsun *et al.*<sup>12</sup>. RDFE's natural properties—combining colour, antioxidant, antibacterial, and antibiofilm activities—position it as

a safer alternative for pediatric applications, where adverse reactions to synthetic agents are a significant concern. The potential for RDFE as a multifunctional additive in dental hygiene products aligns with the growing consumer and regulatory preference for natural product formulations, which may enhance compliance and safety in pediatric and general populations.

A limitation of the present study lies in RDFE's relatively weak colouring ability in powder form, which may limit its efficacy as a standalone visual indicator. Future research should explore optimizing the extraction process to concentrate natural pigments or evaluate biocompatible synthetic colourants that could supplement RDFE's properties without diminishing its health benefits. Additionally, investigating RDFE's impact on diverse oral microbiota could expand its application scope and efficacy insights, contributing to a more comprehensive understanding of its role in oral health maintenance.

## Conclusion

Red dragon fruit extract exhibits promising antioxidant (IC<sub>50</sub> 158.43 $\pm$ 3.67  $\mu$ g/mL), antibacterial (92.34% inhibition at an MBC of 1000  $\mu$ g/mL), and antibiofilm activity (91.57% at 1000  $\mu$ g/mL), supporting its potential as a natural base for dental disclosing agents. The development of a chewy jelly formulation was successful, with Formula II demonstrating acceptable physical characteristics and short-term stability over four weeks. However, the colour intensity was suboptimal, which may limit its effectiveness as a visual disclosing tool. Further optimization is needed to enhance pigment stability and moisture retention, and long-term storage studies are recommended to ensure product durability. Red dragon fruit extract exhibits promising antibacterial and antibiofilm properties, but optimization of colour intensity and *in vivo* validation are needed before clinical application.

## Acknowledgement

We are also thankful to Maranatha Christian University for the internal funding research Scopus output scheme without partner collaboration with a research grant number of 023/SK/ADD/UKM/V/2024 as a source of funding. Thanks to PT FAST for providing BPOM standardized extracts RDFE, and the AMUBBRC team for facilitating and supporting this research.



## References

- Al Samadani K H, Ahmad M, Bakeer H and Elanbya M, Oral health knowledge and practice among 9–12-year-old schoolchildren in the region of Madinah, Saudi Arabia, and its impact on the prevalence of dental caries, *Eur J Gen Dent*, 2017, **6**, 54–58, doi: 10.4103/2278-9626.198615.
- Hasan F, Yuliana L T, Budi H S, Ramasamy R, Ambiya Z I, *et al.*, Prevalence of dental caries among children in Indonesia: A systematic review and meta-analysis of observational studies, *Heliyon*, 2024, **10**(11), e32102, doi: 10.1016/j.heliyon.2024.e32102.
- Vijayakumar A, Sarveswari H B, Vasudevan S, Shanmugam K, Solomon A P, *et al.*, Baicalein inhibits *Streptococcus mutans* biofilms and dental caries-related virulence phenotypes, *Antibiotics (Basel)*, 2021, **10**(2), 215, doi: 10.3390/antibiotics10020215.
- Shenkute D and Asfaw T, *Streptococcus mutans* dental caries among patients attending Debre Berhan Referral Hospital, Ethiopia, *J Bacteriol Parasitol*, 2019, **10**(1), 2, doi: 10.35248/2155-9597.1000350.
- Katsumata T, Le M, Kawada Matsuo M, Taniguchi Y, *et al.*, Comprehensive characterization of sortase A-dependent surface proteins in *Streptococcus mutans*, *Microbiol Immunol*, 2022, **66**(3), 145–156, doi: 10.1111/1348-0421.12954.
- Chen X, Liu C, Peng X, He Y, Liu H, *et al.*, Sortase A-mediated modification of the *Streptococcus mutans* transcriptome and virulence traits, *Mol Oral Microbiol*, 2019, **34**(5), 219–233, doi: 10.1111/omi.12285.
- Matsumi Y, Fujita K, Takashima Y, Yanagida K, Morikawa Y, *et al.*, Contribution of glucan-binding protein A to firm and stable biofilm formation by *Streptococcus mutans*, *Mol Oral Microbiol*, 2014, **30**(3), 217–226, doi: 10.1111/omi.12072.
- Zhou Y, The inhibitory effect of berberine chloride hydrate on *Streptococcus mutans* biofilm formation at different pH values, *Microbiol Spectrum*, 2023, **11**(5), doi: 10.1128/spectrum.04056-22.
- Asti P A, Tripalu S Z, Gunawan E, Jarigau A S and Pratiwi R, Test of anthocyanin level and evaluation of rambutan leaf extract (*Nephelium lappaceum* L.) as an alternative to disclosing agent of dental plaque detection solution, *J Dentomaxillofac Sci*, 2019, **4**(1), 3–6, doi: 10.15562/jdmfs.v4i1.717.
- Oliveira L M, Pazinato J and Zanatta F B, Are oral hygiene instructions with aid of plaque-disclosing methods effective in improving self-performed dental plaque control? A systematic review of randomized controlled trials, *Int J Dent Hyg*, 2021, **19**(3), 239–254, doi: 10.1111/idh.12475.
- Silva M M, Reboredo F H and Lidon F C, Sweetener food additives: A synoptical overview on their chemical properties, applications in food products and side effects, *Emirates J Food Agric*, 2023, **35**, 1–16, doi: 10.9755/ejfa.2023.3202.
- Britsun V M, Popova I V, Kovaleva S A and Petrenko T V, Side effects of modern chemical disinfectants and antiseptics. Part 1. The COVID-19 pandemic as a trigger of non-allergic and allergic effects on the health of the population, *Asthma Allergy*, 2024, **23**(3), 62–70, doi: 10.31655/2307-3373-2024-2-32-37.
- Ramli N S, Ismail P and Rahmat A, Influence of conventional and ultrasonic-assisted extraction on phenolic contents, betacyanin contents, and antioxidant capacity of red dragon fruit (*Hylocereus polyrhizus*), *Sci World J*, 2014, **2014**, 1–10, doi: 10.1155/2014/964731.
- Rahayu Y C, Sabir A and Setyorini D, Antibacterial activity of red dragon fruit extract (*Hylocereus polyrhizus*) on *Streptococcus mutans*, *Int J Appl Pharm*, 2019, **11**(4), 60–63, doi: 10.22159/ijap.2019.v11i4.35293.
- Suwonchoochit T, Nagasevi N, Thamprechavai P and Aroonrerk N, Efficiency of *Hylocereus undatus* extracts on biofilm formation of *Streptococcus mutans* *in vitro*, *Int J Med Sci Curr Res*, 2021, **4**(5), 834–840.
- Shulga D A and Kudryavtsev K V, Ensemble docking as a tool for the rational design of peptidomimetic *Staphylococcus aureus* sortase A inhibitors, *Int J Mol Sci*, 2024, **25**(20), 11279, doi: 10.3390/ijms252011279.
- Soni A N, Varshney M and Tale V S, Molecular docking and pharmacodynamic study of potential inhibitors of *Streptococcus mutans* biofilm, *Biomed Biotechnol Res J*, 2023, **7**(3), 471–477, doi: 10.4103/bbrj.bbrj\_176\_23.
- Widowati W, Ratnawati H, Husin W and Maesaroh M, Antioxidant properties of spice extracts, *Biomed Eng*, 2015, **1**(1), 6.
- Widowati W, Prahastuti S, Hidayat M, Hasianna S T, Wahyudianingsih R, *et al.*, Detam 1 black soybean against cisplatin-induced acute renal failure in a rat model via antioxidant, anti-inflammatory and anti-apoptosis potential, *J Tradit Complement Med*, 2022, **12**, 426–435, doi: 10.1016/j.jtcme.2021.05.006.
- Novilla A, Nawawi A, Sugihartina G and Widowati W, Antioxidative and antibacterial activities of Indonesian propolis extracts against methicillin-resistant *Staphylococcus aureus* (MRSA) *in vitro*, *Cukurova Med J*, 2014, **39**, 224–233.
- Sugiaman V K, Jeffrey J, Naliani S, Pranata N, Lelyana S, *et al.*, Brazilin cream from *Caesalpinia sappan* inhibits periodontal disease: *in vivo* study, *Peer J*, 2024, **12**, e17642, doi: 10.7717/peerj.17642.
- Azzahra F, Yulianti R and Indra, Optimasi gelatin-gliserin pada sediaan gummy candies ekstrak biji jinten hitam (*Nigella sativa* L.), *J Pharmacol Polium*, 2022, **5**(2), 135–141, doi: 10.36465/jop.v5i2.909.
- Komala A M and Haryoto, Tests of ash content, moisture content and dry shrinkage of ethanol extracts of capidada leaves (*Sonneratia alba*) and ketapang (*Terminilia cattapa*), *J Nutraceut Herb Med*, 2020, **3**(1), 10–14, doi: 10.23917/jnhm.v3i1.10549.
- Yusof N, Jaswir I, Jamal P and Jami M S, Texture profile analysis (TPA) of the jelly dessert prepared from halal gelatin extracted using high pressure processing (HPP), *Malaysian J Fund Appl Sci*, 2019, **15**(4), 604–608, doi: 10.11113/mjfas.v15n4.1583.
- Malau R C, Nasution S W, Nasution A N, Widowati W, Nindya F S, *et al.*, Antioxidant potential of temulawak (*Curcuma xanthorrhiza*) extract gel as a candidate for wound healing, *J Voc Health Stud*, 2024, **7**(3), 166–174, doi: 10.20473/jvhs.V7.I3.2024.166-174.
- Kusumawati I G A W, Putra I M W A and Yogeswara I B A, *In vitro* ACE inhibitory activity and bioactive compounds of aqueous extract of *Citrus amblycarpa*, *Majalah Obat Trad*, 2021, **26**(2), 117–122, doi: 10.22146/mot.65028.

- 27 Devgun M, Prasad S, Khokra S and Narang R, Molecular docking studies of dihydropyridazin-3(2H)-one derivatives as antifungal, antibacterial and anti-helminthic agents, *World J Adv Res Rev*, 2021, **12**(1), 186–214, doi: 10.30574/wjarr.2021.12.1.0476.
- 28 Pinzi L and Rastelli G, Molecular docking: shifting paradigms in drug discovery, *Int J Mol Sci*, 2019, **20**(18), 4331, doi: 10.3390/ijms20184331.
- 29 Salmanli M, Yilmaz G T and Tuzuner T, Investigation of the antimicrobial activities of various antimicrobial agents on *Streptococcus mutans* Sortase A through computer-aided drug design (CADD) approaches, *Comput Methods Programs Biomed*, 2021, **212**, 106454.
- 30 Fidrianny I, Ilham N and Hartati R, Antioxidant profile and phytochemical content of different parts of super red dragon fruit (*Hylocereus costaricensis*) collected from West Java-Indonesia, *Asian J Pharm Clin Res*, 2017, **10**(12), 179–183, doi: 10.22159/ajpcr.2017.v10i12.21571.
- 31 Wahdaningsih S, Wahyuono S, Riyanto S and Murwati D, Antioxidant and antibacterial activity of dragon fruit (*Hylocereus* spp.) peel extract, *Int J Pharmacol*, 2015, **11**(2), 186–192, doi: 10.5398/medpet.2017.40.1.47.
- 32 Fidrianny I, Sahar N A and Ruslan W K, Evaluation of antioxidant activities from various extracts of dragon fruit peels using DPPH, ABTS assays and correlation with phenolic, flavonoid, carotenoid content, *Int J Res Pharm Sci*, 2014, **5**(2), 105–112.
- 33 Amalia S, Wahdaningsih S and Untari E K, Antibacterial activity testing of n-hexane fraction of red dragon (*Hylocereus polyrhizus* Britton & Rose) fruit peel on *Staphylococcus aureus* ATCC 25923, *Tradit Med J*, 2014, **19**(2), 93–96.
- 34 Hendra R, Masdeatresia L, Abdulah R and Haryani Y, Antibacterial activity of red dragon peel (*Hylocereus polyrhizus*) pigment, *J Phys Conf Ser*, 2019, **1351**, 012016, doi: 10.1088/1742-6596/1351/1/012016.
- 35 Urip S K, Wardani T S and Listyani T A, Perbandingan aktivitas antibiofilm ekstrak biji kopi hijau dan sangrai kopi robusta terhadap *Staphylococcus aureus*, *Publikasi Penelit Terapan Kebijakan*, 2023, **6**(2), 172–181.
- 36 Yong Y Y, Dykes G, Lee S M and Choo W S, Biofilm inhibiting activity of betacyanins from red pitahaya (*Hylocereus polyrhizus*) and red spinach (*Amaranthus dubius*) against *Staphylococcus aureus* and *Pseudomonas aeruginosa* biofilms, *J Appl Microbiol*, 2018, **126**(1), 271–283, doi: 10.1111/jam.14115.
- 37 de Oliveira C I, Purgato G A, Piccolo M S, Pizziolo V R, Coelho R R, *et al.*, *In vitro* anticariogenic and antibiofilm activities of toothpastes formulated with essential oils, *Arch Oral Biol*, 2020, **117**, 104834, doi: 10.1016/j.archoralbio.2020.104834.
- 38 Calmarza-Chueca F, Sánchez-Gimeno A C, Raso-Pueyo J, Arbones-Mainar J M, Caverni-Muñoz A, *et al.*, Rheological properties and stability of thickeners for clinical use, *Nutrients*, 2022, **14**(17), 3455, doi: 10.3390/nu14173455.